

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau



Part 5 Paper #A1  
16

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification <sup>6</sup> : C07H 19/00, 21/00, C12N 5/00, 15/00, 1/20, C12P 21/06, C12Q 1/68, 1/00, C07K 1/00, 2/00, 4/00, 14/00, 16/00, A61K 35/14</p>	<p>A1</p>	<p>(11) International Publication Number: <b>WO 95/17412</b>  (43) International Publication Date: 29 June 1995 (29.06.95)</p>
<p>(21) International Application Number: PCT/US94/14832 (22) International Filing Date: 21 December 1994 (21.12.94)  (30) Priority Data: 08/173,497 23 December 1993 (23.12.93) US 08/286,889 5 August 1994 (05.08.94) US  (71) Applicant: ICOS CORPORATION [US/US]; 22021 20th Avenue, S.E., Bothell, WA 98021 (US).  (72) Inventors: GALLATIN, W., Michael; 8412 S.E. 33rd Place, Seattle, WA 98040 (US). VAN DER VIEREN, Monica; 7306 20th N.E., Seattle, WA 98115 (US).  (74) Agent: BORUN, Michael, F.; Marshall, O'Toole, Gerstein, Murray &amp; Borun, 233 South Wacker Drive, 6300 Sears Tower, Chicago, IL 60606 (US).</p>		<p>(81) Designated States: AU, BR, CA, CN, CZ, FI, HU, JP, NO, PL, RU, SK, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).  <b>Published</b> <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i></p>
<p>(54) Title: NOVEL HUMAN <math>\beta_2</math> INTEGRIN ALPHA SUBUNIT</p> <p>(57) Abstract</p> <p>DNA encoding a novel human <math>\beta_2</math> integrin <math>\alpha</math> subunit polypeptide, designated <math>\alpha_d</math>, is disclosed along with methods and materials for production of the same by recombinant procedures. Fusion proteins are also disclosed which include extracellular <math>\alpha_d</math> protein fragments, <math>\alpha_d</math> I domain fragments or full length <math>\alpha_d</math> polypeptides and human immunoglobulin constant regions. Binding molecules specific for <math>\alpha_d</math> are also disclosed as useful for modulating the biological activities of <math>\alpha_d</math>. DNA from other species which show homology to human <math>\alpha_d</math> encoding sequences are also disclosed.</p>		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

- 1 -

## Novel Human $\beta_2$ Integrin Alpha Subunit

This application is a continuation-in-part of U.S. Application Serial No. 08/286,889, filed August 5, 1994, which is pending, which in turn is a continuation-in-part of U.S. Application Serial No. 08/173,497, filed December 23, 1993, which is pending.

### Field of the Invention

The present invention relates to the cloning and expression of polynucleotides encoding a novel human  $\beta_2$  integrin  $\alpha$  subunit, designated  $\alpha_d$ , which is structurally related to the known human  $\beta_2$  integrin  $\alpha$  subunits, CD11a, CD11b and CD11c. The present invention also relates to polynucleotides isolated from other species which show homology to human  $\alpha_d$  encoding sequences.

### Background of the Invention

The integrins are a class of membrane-associated molecules which actively participate in cellular adhesion. Integrins are transmembrane heterodimers comprising an  $\alpha$  subunit in noncovalent association with a  $\beta$  subunit. To date, at least fourteen  $\alpha$  subunits and eight  $\beta$  subunits have been identified [reviewed in Springer, *Nature* 346:425-434 (1990)]. The  $\beta$  subunits are generally capable of association with more than one  $\alpha$  subunit and the heterodimers sharing a common  $\beta$  subunit have been classified as subfamilies within the integrin population.

One class of human integrins, restricted to expression in white blood cells, is characterized by a common  $\beta_2$  subunit. As a result of this cell-specific expression, these integrins are commonly referred to as the leukocyte integrins, Leu-CAMs or leukointegrins. Because of the common  $\beta_2$  subunit, an alternative designation of this class is the  $\beta_2$  integrins. The  $\beta_2$  subunit (CD18) has previously been isolated in association with one of three distinct  $\alpha$  subunits,

- 2 -

CD11a, CD11b or CD11c. The isolation of a cDNA encoding human CD18 is described in Kishimoto, *et al.*, *Cell* 48:681-690 (1987). In official WHO nomenclature, the heterodimeric proteins are referred to as CD11a/CD18, CD11b/CD18, and CD11c/CD18; in common nomenclature they are referred to as LFA-1, Mac-1 or Mo1 and p150,95 or LeuM5, respectively [Cobbold, *et al.*, in *Leukocyte Typing III*, McMichael (ed), Oxford Press, p.788 (1987)]. The human  $\beta_2$  integrin  $\alpha$  subunits CD11a, CD11b and CD11c have been demonstrated to migrate under reducing condition in electrophoresis with apparent molecular weights of approximately 180 kD, 155 kD and 150 kD, respectively, and DNAs encoding these subunits have been cloned [CD11a, Larson, *et al.*, *J. Cell Biol.* 108:703-712 (1989); CD11b, Corbi, *et al.*, *J. Biol. Chem.* 263:12403-12411 (1988) and CD11c, Corbi, *et al.*, *EMBO J.* 6:4023-4028 (1987)]. Putative homologs of the human  $\beta_2$  integrin  $\alpha$  and  $\beta$  chains, defined by approximate similarity in molecular weight, have been variously identified in other species including monkeys and other primates [Letvin, *et al.*, *Blood* 61:408-410 (1983)], mice [Sanchez-Madrid, *et al.*, *J. Exp. Med.* 154:1517 (1981)], and dogs [Moore, *et al.*, *Tissue Antigens* 36:211-220 (1990)].

The absolute molecular weights of presumed homologs from other species have been shown to vary significantly [see, *e.g.*, Danilenko *et al.*, *Tissue Antigens* 40:13-21 (1992)], and in the absence of sequence information, a definitive correlation between human integrin subunits and those identified in other species has not been possible. Moreover, variation in the number of members in a protein family has been observed between different species. Consider, for example, that more IgA isotypes have been isolated in rabbits than in humans [Burnett, *et al.*, *EMBO J.* 8:4041-4047 (1989) and Schneiderman, *et al.*, *Proc. Natl. Acad. Sci. (USA)* 86:7561-7565 (1989)]. Similarly, in humans, at least six variants of the metallothioneine protein have been previously identified [Karin and Richards, *Nature* 299:797-802 (1982) and Varshney, *et al.*, *Mol. Cell. Biol.* 6:26-37, (1986)], whereas in the mouse, only two such variants are

- 3 -

in evidence [Searle, *et al.*, *Mol.Cell.Biol.* 4:1221-1230 (1984)]. Therefore, existence of multiple members of a protein family in one species does not necessarily imply that corresponding family members exist in another species.

5 In the specific context of  $\beta_2$  integrins, in dogs it has been observed that the presumed canine  $\beta_2$  counterpart to the human CD18 is capable of dimer formation with as many as four potentially distinct  $\alpha$  subunits [Danilenko, *et al.*, *supra*]. Antibodies generated by immunizing mice with canine splenocytes resulted in monoclonal antibodies which immunoprecipitated proteins tentatively designated as canine homologs to human CD18, CD11a, CD11b and CD11c based  
10 mainly on similar, but not identical, molecular weights. Another anti-canine splenocyte antibody, Ca11.8H2, recognized and immunoprecipitated a fourth  $\alpha$ -like canine subunit also capable of association with the  $\beta_2$  subunit, but having a unique molecular weight and restricted in expression to a subset of differentiated tissue macrophages. Antibodies generated by immunization of hamsters with  
15 murine dendritic cells resulted in two anti-integrin antibodies [Metlay, *et al.*, *J.Exp.Med.* 171:1753-1771 (1990)]. One antibody, 2E6, immunoprecipitated a predominant heterodimer with subunits having approximate molecular weights of 180 kD and 90 kD in addition to minor bands in the molecular weight range of 150-160 kD. The second antibody, N418, precipitated another apparent  
20 heterodimer with subunits having approximate molecular weights of 150 kD and 90 Kd. Based on cellular adhesion blocking studies, it was hypothesized that antibody 2E6 recognized a murine counterpart to human CD18. While the molecular weight of the N418 antigen suggested recognition of a murine homolog to human CD11c/CD18, further analysis indicated that the murine antigen exhibited a tissue distribution pattern which was inconsistent with that observed  
25 for human CD11c/CD18.

The antigens recognized by the canine Ca11.8H2 antibody and the murine N418 antibody could represent a variant species (*e.g.*, a glycosylation or splice variant) of a previously identified canine or murine  $\alpha$  subunit.

- 4 -

Alternatively, these antigens may represent unique canine and murine integrin  $\alpha$  subunits. In the absence of specific information regarding primary structure, these alternatives cannot be distinguished.

In humans, CD11a/CD18 is expressed on all leukocytes. CD11b/CD18 and CD11c/CD18 are essentially restricted to expression on monocytes, granulocytes, macrophages and natural killer (NK) cells, but CD11c/CD18 is also detected on some B-cell types. In general, CD11a/CD18 predominates on lymphocytes, CD11b/CD18 on granulocytes and CD11c/CD18 on macrophages [see review, Arnaout, *Blood* 75:1037-1050 (1990)]. Expression of the  $\alpha$  chains, however, is variable with regard to the state of activation and differentiation of the individual cell types [See review, Larson and Springer, *Immunol.Rev.* 114:181-217 (1990).]

The involvement of the  $\beta_2$  integrins in human immune and inflammatory responses has been demonstrated using monoclonal antibodies which are capable of blocking  $\beta_2$  integrin-associated cell adhesion. For example, CD11a/CD18, CD11b/CD18 and CD11c/CD18 actively participate in natural killer (NK) cell binding to lymphoma and adenocarcinoma cells [Patarroyo, *et al.*, *Immunol.Rev.* 114:67-108 (1990)], granulocyte accumulation [Nourshargh, *et al.*, *J.Immunol.* 142:3193-3198 (1989)], granulocyte-independent plasma leakage [Arfors, *et al.*, *Blood* 69:338-340 (1987)], chemotactic response of stimulated leukocytes [Arfors, *et al.*, *supra*] and leukocyte adhesion to vascular endothelium [Price, *et al.*, *J.Immunol.* 139:4174-4177 (1987) and Smith, *et al.*, *J.Clin.Invest.* 83:2008-2017 (1989)]. The fundamental role of  $\beta_2$  integrins in immune and inflammatory responses is made apparent in the clinical syndrome referred to as leukocyte adhesion deficiency (LAD), wherein clinical manifestations include recurrent and often life threatening bacterial infections. LAD results from heterogeneous mutations in the  $\beta_2$  subunit [Kishimoto, *et al.*, *Cell* 50:193-202 (1987)] and the severity of the disease state is proportional to the degree of the

- 4 -

Alternatively, these antigens may represent unique canine and murine integrin  $\alpha$  subunits. In the absence of specific information regarding primary structure, these alternatives cannot be distinguished.

In humans, CD11a/CD18 is expressed on all leukocytes. CD11b/CD18 and CD11c/CD18 are essentially restricted to expression on monocytes, granulocytes, macrophages and natural killer (NK) cells, but CD11c/CD18 is also detected on some B-cell types. In general, CD11a/CD18 predominates on lymphocytes, CD11b/CD18 on granulocytes and CD11c/CD18 on macrophages [see review, Arnaout, *Blood* 75:1037-1050 (1990)]. Expression of the  $\alpha$  chains, however, is variable with regard to the state of activation and differentiation of the individual cell types [See review, Larson and Springer, *Immunol.Rev.* 114:181-217 (1990).]

The involvement of the  $\beta_2$  integrins in human immune and inflammatory responses has been demonstrated using monoclonal antibodies which are capable of blocking  $\beta_2$  integrin-associated cell adhesion. For example, CD11a/CD18, CD11b/CD18 and CD11c/CD18 actively participate in natural killer (NK) cell binding to lymphoma and adenocarcinoma cells [Patarroyo, *et al.*, *Immunol.Rev.* 114:67-108 (1990)], granulocyte accumulation [Nourshargh, *et al.*, *J.Immunol.* 142:3193-3198 (1989)], granulocyte-independent plasma leakage [Arfors, *et al.*, *Blood* 69:338-340 (1987)], chemotactic response of stimulated leukocytes [Arfors, *et al.*, *supra*] and leukocyte adhesion to vascular endothelium [Price, *et al.*, *J.Immunol.* 139:4174-4177 (1987) and Smith, *et al.*, *J.Clin.Invest.* 83:2008-2017 (1989)]. The fundamental role of  $\beta_2$  integrins in immune and inflammatory responses is made apparent in the clinical syndrome referred to as leukocyte adhesion deficiency (LAD), wherein clinical manifestations include recurrent and often life threatening bacterial infections. LAD results from heterogeneous mutations in the  $\beta_2$  subunit [Kishimoto, *et al.*, *Cell* 50:193-202 (1987)] and the severity of the disease state is proportional to the degree of the

- 3 -

in evidence [Searle, *et al.*, *Mol.Cell.Biol.* 4:1221-1230 (1984)]. Therefore, existence of multiple members of a protein family in one species does not necessarily imply that corresponding family members exist in another species.

5 In the specific context of  $\beta_2$  integrins, in dogs it has been observed that the presumed canine  $\beta_2$  counterpart to the human CD18 is capable of dimer formation with as many as four potentially distinct  $\alpha$  subunits [Danilenko, *et al.*, *supra*]. Antibodies generated by immunizing mice with canine splenocytes resulted in monoclonal antibodies which immunoprecipitated proteins tentatively designated as canine homologs to human CD18, CD11a, CD11b and CD11c based  
10 mainly on similar, but not identical, molecular weights. Another anti-canine splenocyte antibody, Ca11.8H2, recognized and immunoprecipitated a fourth  $\alpha$ -like canine subunit also capable of association with the  $\beta_2$  subunit, but having a unique molecular weight and restricted in expression to a subset of differentiated tissue macrophages. Antibodies generated by immunization of hamsters with  
15 murine dendritic cells resulted in two anti-integrin antibodies [Metlay, *et al.*, *J.Exp.Med.* 171:1753-1771 (1990)]. One antibody, 2E6, immunoprecipitated a predominant heterodimer with subunits having approximate molecular weights of 180 kD and 90 kD in addition to minor bands in the molecular weight range of 150-160 kD. The second antibody, N418, precipitated another apparent  
20 heterodimer with subunits having approximate molecular weights of 150 kD and 90 Kd. Based on cellular adhesion blocking studies, it was hypothesized that antibody 2E6 recognized a murine counterpart to human CD18. While the molecular weight of the N418 antigen suggested recognition of a murine homolog to human CD11c/CD18, further analysis indicated that the murine antigen  
25 exhibited a tissue distribution pattern which was inconsistent with that observed for human CD11c/CD18.

The antigens recognized by the canine Ca11.8H2 antibody and the murine N418 antibody could represent a variant species (*e.g.*, a glycosylation or splice variant) of a previously identified canine or murine  $\alpha$  subunit.



- 5 -

deficiency in  $\beta_2$  subunit expression. Formation of the complete integrin heterodimer is impaired by the  $\beta_2$  mutation [Kishimoto, *et al.*, *supra*].

Interestingly, at least one antibody specific for CD18 has been shown to inhibit human immunodeficiency virus type-1 (HIV-1) syncytia formation *in vitro*, albeit the exact mechanism of this inhibition is unclear [Hildreth and Orentas, *Science* 244:1075-1078 (1989)]. This observation is consistent with the discovery that a principal counterreceptor of CD11a/CD18, ICAM-1, is also a surface receptor for the major group of rhinovirus serotypes [Greve, *et al.*, *Cell* 56:839 (1989)].

The significance of  $\beta_2$  integrin binding activity in human immune and inflammatory responses underscores the necessity to develop a more complete understanding of this class of surface proteins. Identification of yet unknown members of this subfamily, as well as their counterreceptors, and the generation of monoclonal antibodies or other soluble factors which can alter biological activity of the  $\beta_2$  integrins will provide practical means for therapeutic intervention in  $\beta_2$  integrin-related immune and inflammatory responses.

#### **Brief Description of the Invention**

In one aspect, the present invention provides novel purified and isolated polynucleotides (*e.g.*, DNA and RNA transcripts, both sense and anti-sense strands) encoding a novel human  $\beta_2$  integrin  $\alpha$  subunit,  $\alpha_d$ , and variants thereof (*i.e.*, deletion, addition or substitution analogs) which possess binding and/or immunological properties inherent to  $\alpha_d$ . Preferred DNA molecules of the invention include cDNA, genomic DNA and wholly or partially chemically synthesized DNA molecules. A presently preferred polynucleotide is the DNA as set forth in SEQ ID NO: 1, encoding the polypeptide of SEQ ID NO: 2. Also provided are recombinant plasmid and viral DNA constructions (expression constructs) which include  $\alpha_d$  encoding sequences, wherein the  $\alpha_d$  encoding

- 6 -

sequence is operatively linked to a homologous or heterologous transcriptional regulatory element or elements.

Also provided by the present invention are isolated and purified mouse and rat polynucleotides which exhibit homology to polynucleotides encoding human  $\alpha_d$ . A preferred mouse polynucleotide is set forth in SEQ ID NO: 52; a preferred rat polynucleotide is set forth in SEQ ID NO: 54.

As another aspect of the invention, prokaryotic or eukaryotic host cells transformed or transfected with DNA sequences of the invention are provided which express  $\alpha_d$  polypeptide or variants thereof. Host cells of the invention are particularly useful for large scale production of  $\alpha_d$  polypeptide, which can be isolated from either the host cell itself or from the medium in which the host cell is grown. Host cells which express  $\alpha_d$  polypeptide on their extracellular membrane surface are also useful as immunogens in the production of  $\alpha_d$ -specific antibodies. Preferably, host cells transfected with  $\alpha_d$  will be co-transfected to express a  $\beta_2$  integrin subunit in order to allow surface expression of the heterodimer.

Also provided by the present invention are purified and isolated  $\alpha_d$  polypeptides, fragments and variants thereof. Preferred  $\alpha_d$  polypeptides are as set forth in SEQ ID NO: 2. Novel  $\alpha_d$  products of the invention may be obtained as isolates from natural sources, but, along with  $\alpha_d$  variant products, are preferably produced by recombinant procedures involving host cells of the invention. Completely glycosylated, partially glycosylated and wholly deglycosylated forms of the  $\alpha_d$  polypeptide may be generated by varying the host cell selected for recombinant production and/or post-isolation processing. Variant  $\alpha_d$  polypeptides of the invention may comprise water soluble and insoluble  $\alpha_d$  polypeptides including analogs wherein one or more of the amino acids are deleted or replaced: (1) without loss, and preferably with enhancement, of one or more biological activities or immunological characteristics specific for  $\alpha_d$ ; or (2) with specific disablement of a particular ligand/receptor binding or signalling

- 10 -

library. Alternatively, polymerase chain reaction (PCR) using oligonucleotide primers that are designed based on the known cDNA sequence can be used to amplify and identify genomic  $\alpha_d$  DNA sequences. Synthetic DNAs encoding the  $\alpha_d$  polypeptide, including fragments and other variants thereof, may be produced by conventional synthesis methods.

DNA sequence information of the invention also makes possible the development, by homologous recombination or "knockout" strategies [see, e.g., Kapecchi, *Science* 244:1288-1292 (1989)], to produce rodents that fail to express a functional  $\alpha_d$  polypeptide or that express a variant  $\alpha_d$  polypeptide. Such rodents are useful as models for studying the activities of  $\alpha_d$  and  $\alpha_d$  modulators *in vivo*.

DNA and amino acid sequences of the invention also make possible the analysis of  $\alpha_d$  epitopes which actively participate in counterreceptor binding as well as epitopes which may regulate, rather than actively participate in, binding. Identification of epitopes which may participate in transmembrane signal transduction is also comprehended by the invention.

DNA of the invention is also useful for the detection of cell types which express  $\alpha_d$  polypeptide. Standard DNA/RNA hybridization techniques which utilize  $\alpha_d$  DNA to detect  $\alpha_d$  RNA may be used to determine the constitutive level of  $\alpha_d$  transcription within a cell, as well as changes in the level of transcription in response to internal or external agents. Identification of agents which modify transcription and/or translation of  $\alpha_d$  can, in turn, be assessed for potential therapeutic or prophylactic value. DNA of the invention also makes possible *in situ* hybridization of  $\alpha_d$  DNA to cellular RNA to determine the cellular localization of  $\alpha_d$  specific messages within complex cell populations and tissues.

DNA of the invention is also useful for identification of non-human polynucleotide sequences which display homology to human  $\alpha_d$  sequences. Possession of non-human  $\alpha_d$  DNA sequences permits development of animal models (including, for example, transgenic models) of the human system.

- 9 -

transfecting appropriate host cells with a DNA construct comprising a reporter gene under the control of a promoter regulated by a transcription factor having a DNA-binding domain and an activating domain, expressing in the host cells a first hybrid DNA sequence encoding a first fusion of part or all of  $\alpha_d$  and either the DNA binding domain or the activating domain of the transcription factor, expressing in the host cells a library of second hybrid DNA sequences encoding second fusions of part or all of putative  $\alpha_d$  binding proteins and the DNA binding domain or activating domain of the transcription factor which is not incorporated in the first fusion, detecting binding of an  $\alpha_d$  binding protein to  $\alpha_d$  in a particular host cell by detecting the production of reporter gene product in the host cell, and isolating second hybrid DNA sequences encoding  $\alpha_d$  binding protein from the particular host cell.

Hybridoma cell lines which produce antibodies specific for  $\alpha_d$  are also comprehended by the invention. Techniques for producing hybridomas which secrete monoclonal antibodies are well known in the art. Hybridoma cell lines may be generated after immunizing an animal with purified  $\alpha_d$ , variants of  $\alpha_d$  or cells which express  $\alpha_d$  or a variant thereof on the extracellular membrane surface. Immunogen cell types include cells which express  $\alpha_d$  *in vivo*, or transfected prokaryotic or eukaryotic cell lines which normally do not normally express  $\alpha_d$  *in vivo*.

The value of the information contributed through the disclosure of the DNA and amino acid sequences of  $\alpha_d$  is manifest. In one series of examples, the disclosed  $\alpha_d$  cDNA sequence makes possible the isolation of the human  $\alpha_d$  genomic DNA sequence, including transcriptional control elements for the genomic sequence. Identification of  $\alpha_d$  allelic variants and heterologous species (e.g., rat or mouse) DNAs is also comprehended. Isolation of the human  $\alpha_d$  genomic DNA and heterologous species DNAs can be accomplished by standard DNA/DNA hybridization techniques, under appropriately stringent conditions, using all or part of the  $\alpha_d$  cDNA sequence as a probe to screen an appropriate

- 8 -

thereof on a solid support coated (or impregnated with) a fluorescent agent, labelling the ligand with a compound capable of exciting the fluorescent agent, contacting the immobilized  $\alpha_d$  with the labelled ligand in the presence and absence of a putative modulator compound, detecting light emission by the fluorescent agent, and identifying modulating compounds as those compounds that affect the emission of light by the fluorescent agent in comparison to the emission of light by the fluorescent agent in the absence of a modulating compound. Alternatively, the  $\alpha_d$  ligand may be immobilized and  $\alpha_d$  may be labelled in the assay.

Yet another method contemplated by the invention for identifying compounds that modulate the interaction between  $\alpha_d$  and a ligand involves transforming or transfecting appropriate host cells with a DNA construct comprising a reporter gene under the control of a promoter regulated by a transcription factor having a DNA-binding domain and an activating domain, expressing in the host cells a first hybrid DNA sequence encoding a first fusion of part or all of  $\alpha_d$  and either the DNA binding domain or the activating domain of the transcription factor, expressing in the host cells a second hybrid DNA sequence encoding part or all of the ligand and the DNA binding domain or activating domain of the transcription factor which is not incorporated in the first fusion, evaluating the effect of a putative modulating compound on the interaction between  $\alpha_d$  and the ligand by detecting binding of the ligand to  $\alpha_d$  in a particular host cell by measuring the production of reporter gene product in the host cell in the presence or absence of the putative modulator, and identifying modulating compounds as those compounds altering production of the reported gene product in comparison to production of the reporter gene product in the absence of the modulating compound. Presently preferred for use in the assay are the *lexA* promoter, the *lexA* DNA binding domain, the GAL4 transactivation domain, the *lacZ* reporter gene, and a yeast host cell.

he  
ent  
A modified version of the foregoing assay may be used in isolating nucleotide encoding a protein that binds to  $\alpha_d$  by transforming or

- 7 -

function. Fusion polypeptides are also provided, wherein  $\alpha_d$  amino acid sequences are expressed contiguously with amino acid sequences from other polypeptides. Such fusion polypeptides may possess modified biological, biochemical, and/or immunological properties in comparison to wild-type  $\alpha_d$ .  
5 Analog polypeptides including additional amino acid (*e.g.*, lysine or cysteine) residues that facilitate multimer formation are contemplated.

Also comprehended by the present invention are polypeptides and other non-peptide molecules which specifically bind to  $\alpha_d$ . Preferred binding molecules include antibodies (*e.g.*, monoclonal and polyclonal antibodies),  
10 counterreceptors (*e.g.*, membrane-associated and soluble forms) and other ligands (*e.g.*, naturally occurring or synthetic molecules), including those which competitively bind  $\alpha_d$  in the presence of  $\alpha_d$  monoclonal antibodies and/or specific counterreceptors. Binding molecules are useful for purification of  $\alpha_d$  polypeptides and identifying cell types which express  $\alpha_d$ . Binding molecules are also useful for  
15 modulating (*i.e.*, inhibiting, blocking or stimulating) of *in vivo* binding and/or signal transduction activities of  $\alpha_d$ .

Assays to identify  $\alpha_d$  binding molecules are also provided, including immobilized ligand binding assays, solution binding assays, scintillation proximity assays, di-hybrid screening assays, and the like.

20 *In vitro* assays for identifying antibodies or other compounds that modulate the activity of  $\alpha_d$  may involve, for example, immobilizing  $\alpha_d$  or a natural ligand to which  $\alpha_d$  binds, detectably labelling the nonimmobilized binding partner, incubating the binding partners together and determining the effect of a test compound on the amount of label bound wherein a reduction in the label  
25 bound in the presence of the test compound compared to the amount of label bound in the absence of the test compound indicates that the test agent is an inhibitor of  $\alpha_d$  binding.

Another type of assay for identifying compounds that modulate the interaction between  $\alpha_d$  and a ligand involves immobilizing  $\alpha_d$  or a fragment

- 11 -

As another aspect of the invention, monoclonal or polyclonal antibodies specific for  $\alpha_d$  may be employed in immunohistochemical analysis to localize  $\alpha_d$  to subcellular compartments or individual cells within tissues. Immunohistochemical analyses of this type are particularly useful when used in combination with *in situ* hybridization to localize both  $\alpha_d$  mRNA and polypeptide products of the  $\alpha_d$  gene.

Identification of cell types which express  $\alpha_d$  may have significant ramifications for development of therapeutic and prophylactic agents. It is anticipated that the products of the invention related to  $\alpha_d$  can be employed in the treatment of diseases wherein macrophages are an essential element of the disease process. Animal models for many pathological conditions associated with macrophage activity have been described in the art. For example, in mice, macrophage recruitment to sites of both chronic and acute inflammation is reported by Jutila, *et al.*, *J. Leukocyte Biol.* 54:30-39 (1993). In rats, Adams, *et al.*, [*Transplantation* 53:1115-1119(1992) and *Transplantation* 56:794-799 (1993)] describe a model for graft arteriosclerosis following heterotropic abdominal cardiac allograft transplantation. Rosenfeld, *et al.*, [*Arteriosclerosis* 7:9-23 (1987) and *Arteriosclerosis* 7:24-34 (1987)] describe induced atherosclerosis in rabbits fed a cholesterol supplemented diet. Hanenberg, *et al.*, [*Diabetologia* 32:126-134 (1989)] report the spontaneous development of insulin-dependent diabetes in BB rats. Yamada *et al.*, [*Gastroenterology* 104:759-771 (1993)] describe an induced inflammatory bowel disease, chronic granulomatous colitis, in rats following injections of streptococcal peptidoglycan-polysaccharide polymers. Cromartie, *et al.*, [*J. Exp. Med.* 146:1585-1602 (1977)] and Schwab, *et al.*, [*Infection and Immunity* 59:4436-4442 (1991)] report that injection of streptococcal cell wall protein into rats results in an arthritic condition characterized by inflammation of peripheral joints and subsequent joint destruction. Finally, Huitinga, *et al.*, [*Eur. J. Immunol* 23:709-715 (1993)] describe experimental allergic encephalomyelitis, a model for multiple sclerosis, in Lewis rats. In each of these

- 12 -

models,  $\alpha_d$  antibodies, other  $\alpha_d$  binding proteins, or soluble forms of  $\alpha_d$  are utilized to attenuate the disease state, presumably through inactivation of macrophage activity.

5        Pharmaceutical compositions for treatment of these and other disease states are provided by the invention. Pharmaceutical compositions are designed for the purpose of inhibiting interaction between  $\alpha_d$  and its ligand(s) and include various soluble and membrane-associated forms of  $\alpha_d$  (comprising the entire  $\alpha_d$  polypeptide, or fragments thereof which actively participate in  $\alpha_d$  binding), soluble and membrane-associated forms of  $\alpha_d$  binding proteins  
10        (including antibodies, ligands, and the like), intracellular or extracellular modulators of  $\alpha_d$  binding activity, and/or modulators of  $\alpha_d$  and/or  $\alpha_d$ -ligand polypeptide expression, including modulators of transcription, translation, post-translational processing and/or intracellular transport. The invention also comprehends methods for treatment of disease states in which  $\alpha_d$  binding is  
15        implicated, wherein a patient suffering from said disease state is provided an amount of a pharmaceutical composition of the invention sufficient to modulate levels of  $\alpha_d$  binding. The method of treatment of the invention is applicable to disease states such as, but not limited to, Type I diabetes, atherosclerosis, multiple sclerosis, asthma, psoriasis, and rheumatoid arthritis.

20

#### Brief Description of the Drawing

Numerous other aspects and advantages of the present invention will be apparent upon consideration of the following description thereof, reference being made to the drawing wherein:

25        Figure 1A through 1D comprises an alignment of the human amino acid sequences of CD11b (SEQ ID NO: 3), CD11c (SEQ ID NO: 4) and  $\alpha_d$  (SEQ ID NO: 2).



- 13 -

### Detailed Description of the Invention

The present invention is illustrated by the following examples relating to the isolation of a cDNA clone encoding  $\alpha_d$  from a human spleen cDNA library. More particularly, Example 1 illustrates the use of anti-canine  $\alpha_{TM1}$  antibody in an attempt to detect a homologous human protein. Example 2 details purification of canine  $\alpha_{TM1}$  and N-terminal sequencing of the polypeptide to design oligonucleotide primers for PCR amplification of the canine  $\alpha_{TM1}$  gene. Example 3 addresses large scale purification of canine  $\alpha_{TM1}$  for internal sequencing in order to design additional PCR primers. Example 4 describes use of the PCR and internal sequence primers to amplify a fragment of the canine  $\alpha_{TM1}$  gene. Example 5 addresses cloning of the human  $\alpha_d$ -encoding cDNA sequence. Example 6 describes Northern blot hybridization analysis of human tissues and cells for expression of  $\alpha_d$  mRNA. Example 7 details the construction of human  $\alpha_d$  expression plasmids and transfection of COS cells with the resulting plasmids. Example 8 addresses ELISA analysis of  $\alpha_d$  expression in transfected COS cells. Example 9 describes FACS analysis of COS cells transfected with human  $\alpha_d$  expression plasmids. Example 10 addresses immunoprecipitation of CD18 in association with  $\alpha_d$  in co-transfected COS cells. Example 11 relates to stable transfection of  $\alpha_d$  expression constructs in Chinese hamster ovary cells. Example 12 addresses CD18-dependent binding of  $\alpha_d$  to the intercellular adhesion molecule, ICAM-R. Example 13 describes scintillation proximity screening assays to identify inhibitors of  $\alpha_d$  ligand/anti-ligand binding interactions. Example 14 addresses construction of expression plasmids which encode soluble forms of  $\alpha_d$ . Example 15 relates to production of  $\alpha_d$ -specific monoclonal antibodies. Example 16 describes analysis of  $\alpha_d$  tissue distribution using polyclonal antiserum. Example 17 describes isolation of rat cDNA sequences which show homology to human  $\alpha_d$  gene sequences. Example 18 relates to construction of rat  $\alpha_d$  I domain expression plasmids, including I domain/IgG fusion proteins, and production of monoclonal antibodies to I domain fusion

- 14 -

proteins. Example 19 addresses isolation of mouse cDNA sequences which show homology to human  $\alpha_d$  gene sequences. Example 20 describes isolation of additional mouse  $\alpha_d$  cDNA clones used for conformational sequence analysis. Example 21 relates to *in situ* hybridization analysis of various mouse tissues to determine tissue and cell specific expression of the putative mouse homolog to human  $\alpha_d$ . Example 22 describes generation of expression constructs which encode the putative mouse homolog of human  $\alpha_d$ . Example 23 addresses design of a "knock-out" mouse wherein the gene encoding the putative mouse homolog of human  $\alpha_d$  is disrupted. Example 24 describes isolation of rabbit cDNA clones which show homology to human  $\alpha_d$  encoding sequences. Example 25 describes animal models which resemble human disease states wherein modulation of  $\alpha_d$  is assayed for therapeutic capabilities.

### Example 1

#### Attempt to Detect a Human Homolog of Canine $\alpha_{TM1}$

The monoclonal antibody Ca11.8H2 [Moore, et al., *supra*] specific for canine  $\alpha_{TM1}$  was tested for cross-reactivity on human peripheral blood leukocytes in an attempt to identify a human homolog of canine  $\alpha_{TM1}$ . Cell preparations (typically  $1 \times 10^6$  cells) were incubated with undiluted hybridoma supernatant or a purified mouse IgG-negative control antibody (10  $\mu$ g/ml) on ice in the presence of 0.1 % sodium azide. Monoclonal antibody binding was detected by subsequent incubation with FITC-conjugated horse anti-mouse IgG (Vector Laboratories, Burlingame, CA) at 6  $\mu$ g/ml. Stained cells were fixed with 2 % w/v paraformaldehyde in phosphate buffered saline (PBS) and were analyzed with a Facstar Plus fluorescence-activated cell sorter (Becton Dickinson, Mountain View, CA). Typically, 10,000 cells were analyzed using logarithmic amplification for fluorescence intensity.

The results indicated that Ca11.8H2 did not cross-react with surface proteins expressed on human peripheral blood leukocytes, while the control cells,

- 15 -

neoplastic canine peripheral blood lymphocytes, were essentially all positive for  $\alpha_{TM1}$ .

Because the monoclonal antibody Ca11.8H2 specific for the canine  $\alpha$  subunit did not cross react with a human homolog, isolation of canine  $\alpha_{TM1}$  DNA was deemed a necessary prerequisite to isolate a counterpart human gene if one existed.

### Example 2

#### Affinity Purification Of Canine $\alpha_{TM1}$ For N-Terminal Sequencing

Canine  $\alpha_{TM1}$  was affinity purified in order to determine N-terminal amino acid sequences for oligonucleotide probe/primer design. Briefly, anti- $\alpha_{TM1}$  monoclonal antibody Ca11.8H2 was coupled to Affigel 10 chromatographic resin (BioRad, Hercules, CA) and protein was isolated by specific antibody-protein interaction. Antibody was conjugated to the resin, according to the BioRad suggested protocol, at a concentration of approximately 5 mg antibody per ml of resin. Following the conjugation reaction, excess antibody was removed and the resin blocked with three volumes of 0.1 M ethanolamine. The resin was then washed with thirty column volumes of phosphate buffered saline (PBS).

Twenty-five grams of a single dog spleen were homogenized in 250 ml of buffer containing 0.32 M sucrose in 25 mM Tris-HCl, Ph 8.0, with protease inhibitors. Nuclei and cellular debris were pelleted with centrifugation at 1000 g for 15 minutes. Membranes were pelleted from the supernatant with centrifugation at 100,000 g for 30 minutes. The membrane pellet was resuspended in 200 ml lysis buffer (50 mM NaCl, 50 mM borate, pH 8.0, with 2% NP-40) and incubated for 1 hour on ice. Insoluble material was then pelleted by centrifugation at 100,000 g for 60 minutes. Ten milliliters of the cleared lysate were transferred to a 15 ml polypropylene tube with 0.5 ml Ca11.8H2-conjugated Affigel 10 resin described above. The tube was incubated overnight at 4°C with rotation and the resin subsequently washed with 50 column volumes

- 16 -

D-PBS. The resin was then transferred to a microfuge tube and boiled for ten minutes in 1 ml Laemmli (non-reducing) sample buffer containing 0.1 M Tris-HCl, pH 6.8, 2% SDS, 20% glycerol and 0.002% bromophenol blue. The resin was pelleted by centrifugation and discarded; the supernatant was treated with  
5 1/15 volume  $\beta$ -mercaptoethanol (Sigma, St. Louis, MO) and run on a 7% polyacrylamide gel. The separated proteins were transferred to Immobilon PVDF membrane (Millipore, Bedford, MA) as follows.

The gels were washed once in deionized, Millipore-filtered water and equilibrated for 15-45 minutes in 10 mM 3-[cyclohexylamino]-1-propanesulfonic acid (CAPS) transfer buffer, pH 10.5, with 10% methanol.  
10 Immobilon membranes were moistened with methanol, rinsed with filtered water, and equilibrated for 15-30 minutes in CAPS transfer buffer. The initial transfer was carried out using a Biorad transfer apparatus at 70 volts for 3 hours. The Immobilon membrane was removed after transfer and stained in filtered 0.1%  
15 R250 Coomassie stain for 10 minutes. Membranes were destained in 50% methanol/10% acetic acid three times, ten minutes each time. After destaining, the membranes were washed in filtered water and air-dried.

Protein bands of approximately 150 kD, 95 kD, 50 kD and 30 kD were detected. Presumably the 50 kD and 30 kD bands resulted from antibody  
20 contamination. N-terminal sequencing was then attempted on both the 150 kD and 95 kD bands, but the 95 kD protein was blocked, preventing sequencing. The protein band of 150 kD was excised from the membrane and directly sequenced with an Applied Biosystems (Foster City, CA) Model 473A protein sequencer according to the manufacturer's instructions. The resulting amino acid  
25 sequence is set in SEQ ID NO: 5 using single letter amino acid designations.

FNLDVEEPMVFQ

(SEQ ID NO: 5)

- 17 -

The identified sequence included the FNLD sequence characteristic of  $\alpha$  subunits of the integrin family [Tamura, *et al.*, *J. Cell. Biol.* 111:1593-1604 (1990)].

Primer Design and Attempt to Amplify Canine  $\alpha_{TM1}$  Sequences

5 From the N-terminal sequence information, three oligonucleotide probes were designed for hybridization: a) "Tommer," a fully degenerate oligonucleotide; b) "Patmer," a partially degenerate oligonucleotide; and c) "Guessmer," a nondegenerate oligonucleotide based on mammalian codon usage. These probes are set out below as SEQ ID NOS: 6, 7 and 8, respectively. Nucleic acid symbols are in accordance with 37 C.F.R. §1.882 for these and all other nucleotide sequences herein.

5'-TTYAAYYTGGAYGTNGARGARCCNATGGTNTTYCA-3' (SEQ ID NO: 6)  
5'-TTCAACCTGGACGTGGAGGAGCCCATGGTGTTCCTCAA-3' (SEQ ID NO: 7)  
5'-TTCAACCTGGACGTNGAASANCCCATGGTCTTCCAA-3' (SEQ ID NO: 8)

15 Based on sequencing data, no relevant clones were detected using these oligonucleotides in several low stringency hybridizations to a canine spleen/peripheral blood macrophage cDNA library cloned into  $\lambda$ ZAP (Stratagene, La Jolla, CA).

20 Four other oligonucleotide primers, designated 5'Deg, 5'Spec, 3'Deg and 3'Spec (as set out in SEQ ID NOS: 9, 10, 11 and 12, respectively, wherein Deg indicates degenerate and Spec indicates non-degenerate) were subsequently designed based on the deduced N-terminal sequence for attempts to amplify canine  $\alpha_{TM1}$  sequences by PCR from phage library DNA purified from plate lysates of the Stratagene library described above.

25 5'-TTYAAYYTNGAYGTNGARGARCC-3' (SEQ ID NO: 9)  
5'-TTYAAYYTGGACGTNGAAGA-3' (SEQ ID NO: 10)

- 18 -

5'-TGRAANACCATNGGYTC-3' (SEQ ID NO: 11)

5'-TTGGAAGACCATNGGYTC-3' (SEQ ID NO: 12)

5 The  $\alpha_{TM1}$  oligonucleotide primers were paired with T3 or T7 vector primers, as set out in SEQ ID NOS: 13 and 14, respectively, which hybridize to sequences flanking the polylinker region in the Bluescript phagemid found in  $\lambda$ ZAP.

5'-ATTAACCCTCACTAAAG-3' (SEQ ID NO: 13)

5'-AATACGACTCACTATAG-3' (SEQ ID NO: 14)

10 The PCR amplification was carried out in *Taq* buffer (Boehringer Mannheim, Indianapolis, IN) containing magnesium with 150 ng of library DNA, 1  $\mu$ g of each primer, 200  $\mu$ M dNTPs and 2.5 units *Taq* polymerase (Boehringer Mannheim) and the products were separated by electrophoresis on a 1% agarose gel in Tris-Acetate-EDTA (TAE) buffer with 0.25  $\mu$ g/ml ethidium bromide. DNA was transferred to a Hybond (Amersham, Arlington Heights, IL) membrane by wicking overnight in 10X SSPE. After transfer, the immobilized DNA was denatured with 0.5 M NaOH with 0.6 M NaCl, neutralized with 1.0 M Tris-HCl, pH 8.0, in 1.5 M NaCl, and washed with 2X SSPE before UV crosslinking with a Stratalinker (Stratagene) crosslinking apparatus. The membrane was incubated in prehybridization buffer (5X SSPE, 4X Denhardts, 0.8% SDS, 30% formamide) for 2 hr at 50°C with agitation.

25 Oligonucleotide probes 5'Deg, 5'Spec, 3'Deg and 3'Spec (SEQ ID NOS: 9, 10, 11 and 12, respectively) were labeled using a Boehringer Mannheim kinase buffer with 100-300  $\mu$ Ci  $\gamma$ P<sup>32</sup>-dATP and 1-3 units of polynucleotide kinase for 1-3 hr at 37°C. Unincorporated label was removed with Sephadex G-25 fine (Pharmacia, Piscataway, NJ) chromatography using 10 mM Tris-HCl, pH 8.0, 1 mM EDTA (TE) buffer and the flow-through added directly to the prehybridization solution. Membranes were probed for 16 hr at 42°C with

- 19 -

agitation and washed repeatedly, with a final stringency wash of 1X SSPE/0.1% SDS at 50° for 15 min. The blot was then exposed to Kodak X-Omat AR film for 1-4 hours at -80°C.

5 The oligonucleotides 5'Deg, 5'Spec, 3'Deg and 3'Spec only hybridized to PCR products from the reactions in which they were used as primers and failed to hybridize as expected to PCR products from the reactions in which they were not used as primers. Thus, it was concluded that none of the PCR products were specific for  $\alpha_{TM1}$  because no product hybridized with all of the appropriate probes.

10

### Example 3

#### Large Scale Affinity Purification Of Canine $\alpha_{TM1}$ For Internal Sequencing

15 In order to provide additional amino acid sequence for primer design, canine  $\alpha_{TM1}$  was purified for internal sequencing. Three sections of frozen spleen (approximately 50 g each) and frozen cells from two partial spleens from adult dogs were used to generate protein for internal sequencing. Fifty grams of spleen were homogenized in 200-300 ml borate buffer with a Waring blender. The homogenized material was diluted with 1 volume of buffer containing 4% NP-40, and the mixture then gently agitated for at least one hour. The resulting lysate was cleared of large debris by centrifugation at 2000 g for 20 min, and then filtered through either a Corning (Corning, NY) prefilter or a  
20 Corning 0.8 micron filter. The lysate was further clarified by filtration through the Corning 0.4 micron filter system.

25 Splenic lysate and the antibody-conjugated Affigel 10 resin described in Example 2 were combined at a 150:1 volume ratio in 100 ml aliquots and incubated overnight at 4°C with rocking. The lysate was removed after centrifugation at 1000 g for 5 minutes, combined with more antibody-conjugated Affigel 10 resin and incubated overnight as above. The absorbed resin aliquots were then combined and washed with 50 volumes D-PBS/0.1% Tween 20 and the

- 20 -

resin transferred to a 50 ml Biorad column. Adsorbed protein was eluted from the resin with 3-5 volumes of 0.1 M glycine (pH 2.5); fractions of approximately 900  $\mu$ l were collected and neutralized with 100  $\mu$ l 1 M Tris buffer, pH 8.0. Aliquots of 15  $\mu$ l were removed from each fraction and boiled in an equal volume of 2X Laemmli sample buffer with 1/15 volume 1 M dithiothreitol (DTT). These samples were electrophoresed on 8% Novex (San Diego, CA) polyacrylamide gels and visualized either by Coomassie stain or by silver stain using a Daiichi kit (Enprotech, Natick, MA) according to the manufacturer's suggested protocol. Fractions which contained the largest amounts of protein were combined and concentrated by vacuum. The remaining solution was diluted by 50% with reducing Laemmli sample buffer and run on 1.5 mm 7% polyacrylamide gels in Tris-glycine/SDS buffer. Protein was transferred from the gels to Immobilon membrane by the procedure described in Example 2 using the Hoefer transfer apparatus.

The protein bands corresponding to canine  $\alpha_{TM1}$  were excised from 10 PVDF membranes and resulted in approximately 47  $\mu$ g total protein. The bands were destained in 4 ml 50% methanol for 5 minutes, air dried and cut into 1 x 2 mm pieces. The membrane pieces were submerged in 2 ml 95% acetone at 4°C for 30 minutes with occasional vortexing and then air dried.

Prior to proteolytic cleavage of the membrane bound protein, 3 mg of cyanogen bromide (CNBr) (Pierce, Rockford, IL) were dissolved in 1.25 ml 70% formic acid. This solution was then added to a tube containing the PVDF membrane pieces and the tube incubated in the dark at room temperature for 24 hours. The supernatant (S1) was then removed to another tube and the membrane pieces washed with 0.25 ml 70% formic acid. This supernatant (S2) was removed and added to the previous supernatant (S1). Two milliliters of Milli Q water were added to the combined supernatants (S1 and S2) and the solution lyophilized. The PVDF membrane pieces were dried under nitrogen and extracted again with 1.25 ml 60% acetonitrile, 0.1% tetrafluoroacetic acid (TFA) at 42°C for 17 hours.



- 21 -

This supernatant (S3) was removed and the membrane pieces extracted again with 1.0 ml 80% acetonitrile with 0.08% TFA at 42°C for 1 hour. This supernatant (S4) was combined with the previous supernatants (S1, S2 and S3) and vacuum dried.

5                   The dried CNBr fragments were then dissolved in 63  $\mu$ l 8 M urea, 0.4 M  $\text{NH}_4\text{HCO}_3$ . The fragments were reduced in 5  $\mu$ l 45 mM dithiothreitol (DTT) and subsequently incubated at 50°C for 15 minutes. The solution was then cooled to room temperature and the fragments alkylated by adding 5  $\mu$ l 100 mM iodoacetamide (Sigma, St. Louis, MO). Following a 15 minute incubation at  
10                   room temperature, the sample was diluted with 187  $\mu$ l Milli Q water to a final urea concentration of 2.0 M. Trypsin (Worthington, Freehold, NJ) was then added at a ratio of 1:25 (w:w) of enzyme to protein and the protein digested for 24 hours at 37°C. Digestion was terminated with addition of 30  $\mu$ l TFA.

                  The protein fragments were then separated with high performance  
15                   liquid chromatography (HPLC) on a Waters 625 LC system (Millipore, Milford, MA) using a 2.1 x 250 mm, 5 micron Vydac C-18 column (Vydac, Hesperia, CA) equilibrated in 0.05% TFA and HPLC water (buffer A). The peptides were eluted with increasing concentration of 80% acetonitrile in 0.04% TFA (buffer B) with a gradient of 38-75% buffer B for 65-95 minutes and 75-98% buffer B for  
20                   95-105 minutes. Peptides were fractionated at a flow rate of 0.2 ml/minute and detected at 210 nm.

                  Following fractionation, the amino acid sequence of the peptides was analyzed by automated Edman degradation performed on an Applied Biosystems Model 437A protein sequencer using the manufacturer's standard  
25                   cycles and the Model 610A Data Analysis software program, Version 1.2.1. All sequencing reagents were supplied by Applied Biosystems. The amino acid sequences of seven of the eight internal fragments are set out below wherein "X" indicates the identity of the amino acid was not certain.

- 22 -

	VFQEXGAGFGQ	(SEQ ID NO: 15)
	LYDXVAATGLXQPI	(SEQ ID NO: 16)
	PLEYXDVIPQAE	(SEQ ID NO: 17)
	FQEGFSXVLX	(SEQ ID NO: 18)
5	TSPTFIXMSQENVD	(SEQ ID NO: 19)
	LVVGAPLEVAVXQTGR	(SEQ ID NO: 20)
	LDXKPXDTA	(SEQ ID NO: 21)

Primer Design

One internal amino acid sequence (set out in SEQ ID NO: 22) obtained was then used to design a fully degenerate oligonucleotide primer, designated p4(R) as set out in SEQ ID NO: 23.

	FGEQFSE	(SEQ ID NO: 22)
	5'-RAANCCYTCYTGRAAACTYTC-3'	(SEQ ID NO: 23)

Example 415 PCR Cloning Of A Canine  $\alpha_{TM1}$  Fragment

The 5' portion of the canine  $\alpha_{TM1}$  gene was amplified from double-stranded canine splenic cDNA by PCR.

## A. Generation of Double Stranded Canine Spleen cDNA

One gram of frozen material from a juvenile dog spleen was ground in liquid nitrogen on dry ice and homogenized in 20 ml RNA-Stat 60 buffer (Tel-Test B, Inc, Friendswood, TX). Four ml chloroform were added, and the solution extracted by centrifugation at 12,000 g for 15 minutes. RNA was precipitated from the aqueous layer with 10 ml ethanol. Poly A<sup>+</sup> RNA was then selected on Dynal Oligo dT Dynabeads (Dynal, Oslo, Norway). Five aliquots of 100  $\mu$ g total RNA were combined and diluted with an equal volume of 2X binding

- 23 -

buffer (20 mM Tris-HCl, pH 7.5, 1.0 M LiCl, 1 mM EDTA, 0.1% SDS). RNA was then incubated 5 minutes with the Oligo dT Dynabeads (1.0 ml or 5 mg beads for all the samples). Beads were washed with buffer containing 10 mM Tris-HCl, pH 7.5, 0.15 M LiCl, 1 mM EDTA and 0.1% SDS, according to the manufacturer's suggested protocol prior to elution of poly A<sup>+</sup> mRNA with 2 mM EDTA, pH 7.5. Double-stranded cDNA was then generated using the eluted poly A<sup>+</sup> mRNA and the Boehringer Mannheim cDNA Synthesis Kit according to the manufacturer's suggested protocol.

#### B. Isolation of a Partial Canine $\alpha_{TM1}$ cDNA

Oligonucleotide primers 5'Deg (SEQ ID NO: 9) and p4(R) (SEQ ID NO: 23) were employed in a standard PCR reaction using 150 ng double-stranded cDNA, 500 ng of each primer, 200  $\mu$ M dNTPs and 1.5 units *Taq* polymerase (Boehringer Mannheim) in *Taq* buffer (Boehringer Mannheim) with magnesium. The resulting products (1  $\mu$ l of the original reaction) were subjected to a second round of PCR with the same primers to increase product yield. This band was eluted from a 1% agarose gel onto Schleicher & Schuell (Keene, NH) NA45 paper in a buffer containing 10 mM Tris-HCl, pH 8, 1 mM EDTA, 1.5 M NaCl at 65°C, precipitated, and ligated into the pCR<sup>tm</sup>II vector (Invitrogen, San Diego, CA) using the TA cloning kit (Invitrogen) and the manufacturer's suggested protocol. The ligation mixture was transformed by electroporation into XL-1 Blue bacteria (Stratagene). One clone, 2.7, was determined to contain sequences corresponding to  $\alpha_{TM1}$  peptide sequences which were not utilized in design of the primers.

Sequencing was performed with an Applied Biosystems 373A DNA sequencer (Foster City, CA) with a Dye-deoxy terminator cycle sequence kit (ABI) in which fluorescent-labeled dNTPs were incorporated in an asymmetric PCR reaction [McCabe, "Production of Single Stranded DNA by Asymmetric PCR," in PCR Protocols: A Guide to Methods and Applications, Innis, *et al.*

- 24 -

(eds.) pp. 76-83 Academic Press: New York (1990)] as follows. Samples were held at 96°C for 4 minutes and subjected to 25 cycles of the step sequence: 96°C, for 15 seconds; 50°C for 1 second; 60°C for 4 minutes. Sequence data was automatically down-loaded into sample files on the computer that included chromatogram and text files. The sequence of the entire insert of clone 2.7 is set out in SEQ ID NO: 24.

Attempts to isolate the full length canine  $\alpha_{TM1}$  cDNA from the Stratagene library (as described in Example 2) were unsuccessful. Approximately  $1 \times 10^6$  phage plaques were screened by hybridization under low stringency conditions using 30% formamide with clone 2.7 as a probe, but no positive clones resulted. Attempts to amplify relevant sequences downstream from those represented in clone 2.7 using specific oligonucleotides derived from clone 2.7 or degenerate primers based on amino acid sequence from other peptide fragments paired with a degenerate oligonucleotide based on the conserved  $\alpha$  subunit amino acid motif GFFKR [Tamura, *et al.*, *supra*] were also unsuccessful.

### Example 5

#### Cloning Of A Putative Human Homolog Of Canine $\alpha_{TM1}$

To attempt the isolation of a human sequence homologous to canine  $\alpha_{TM1}$  the approximately 1 kb canine  $\alpha_{TM1}$  fragment from clone 2.7 was used as a probe. The probe was generated by PCR under conditions described in Example 2 using NT2 (as set out in SEQ ID NO: 25) and p4(R) (SEQ ID NO: 23) primers.

5'-GTNTTYCARGARGAYGG-3' (SEQ ID NO: 25)

The PCR product was purified using the Qiagen (Chatsworth, GA) Quick Spin kit and the manufacturer's suggested protocol. The purified DNA (200 ng) was labeled with 200  $\mu$ Ci  $\alpha^{32}$ PdCTP using the Boehringer Mannheim Random Prime

- 25 -

Labelling kit and the manufacturer's suggested protocol. Unincorporated isotope was removed with Sephadex G25 (fine) gravity chromatography. The probe was denatured with 0.2 N NaOH and neutralized with 0.4 M Tris-HCl, pH 8.0, before use.

5 Colony lifts on Hybond filters (Amersham) of a human spleen cDNA library in pCDNA/Amp (Invitrogen, San Diego, CA) were prepared. The filters were initially denatured and neutralized as described in Example 2 and subsequently incubated in a prehybridization solution (8 ml/filter) with 30% formamide at 50°C with gentle agitation for 2 hours. Labeled probe as described  
10 above was added to this solution and incubated with the filters for 14 hours at 42°C. The filters were washed twice in 2X SSC/0.1% SDS at 37°C and twice in 2X SSC/0.1% SDS at 50°C. Final stringency washes were 1X SSC/0.1% SDS, twice at 65°C (1X SSC is 150 mM NaCl, 15 mM sodium citrate, pH 7.0). Filters were exposed to Kodak X-Omat AR film for six hours with an intensifying  
15 screen. Colonies giving signals on duplicate lifts were streaked on LB medium with magnesium (LBM)/carbenicillin plates and incubated overnight at 37°C. Resulting streaked colonies were lifted with Hybond filters and these filters were treated as above. The filters were hybridized under more stringent conditions with the 1 kb probe from clone 2.7, labeled as previously described, in a 50%  
20 formamide hybridization solution at 50°C for 3 hours. Probed filters were washed with a final stringency of 0.1 X SSC/0.1% SDS at 65°C and exposed to Kodak X-Omat AR film for 2.5 hours at -80°C with an intensifying screen. Positive colonies were identified and cultured in LBM/carbenicillin medium overnight. DNA from the cultures was prepared using the Promega Wizard  
25 miniprep kit according to the manufacturer's suggested protocol and the resulting DNA was sequenced.

The initial screening resulted in 18 positive clones, while the secondary screening under more stringent hybridization conditions produced one positive clone which was designated 19A2. The DNA and deduced amino acid

- 26 -

sequences of the human  $\alpha_d$  clone 19A2 are set out in SEQ ID NOS: 1 and 2, respectively.

#### Characteristics Of The Human $\alpha_d$ cDNA and Predicted Polypeptide

5 Clone 19A2 encompasses the entire coding region for the mature protein, plus 48 bases (16 amino acid residues) of the 5' upstream signal sequence and 241 bases of 3' untranslated sequence which do not terminate in a polyadenylation sequence. The core molecular weight of the mature protein is predicted to be around 125 kD. The extracellular domain is predicted to encompass approximately amino acid residues 17 through 1108 of SEQ ID NO:  
10 2. This extracellular region is contiguous with about a 20 amino acid region homologous to the human CD11c transmembrane region (residues 1109 through 1128 of SEQ ID NO: 2). The cytoplasmic domain comprises approximately 30 amino acids (about residues 1129 through 1161 of SEQ ID NO: 2). The protein also contains a region (around residues 150 through 352) of approximately 202  
15 amino acids homologous to the I (insertion) domain common to CD11a, CD11b and CD11c [Larson and Springer, *supra*],  $\alpha_E$  [Shaw, *et al.*, *J.Biol.Chem.* 269:6016-6025 (1994)] and in VLA-1 and VLA-2, [Tamura, *et al.*, *supra*]. The I domain in other integrins has been shown to participate in ICAM binding [Landis, *et al.*, *J.Cell.Biol.* 120:1519-1527 (1993); Diamond, *et al.*, *J.Cell.Biol.*  
20 120:1031-1043 (1993)], suggesting that  $\alpha_d$  may also bind members of the ICAM family of surface molecules. This region has not been demonstrated to exist in any other integrin subunits.

The deduced amino acid sequence of  $\alpha_d$  shows approximately 36% identity to that of CD11a, approximately 60% identity to CD11b and  
25 approximately 66% identity to CD11c. An alignment of amino acid sequences for (CD11b SEQ ID NO: 3), CD11c (SEQ ID NO: 4) and  $\alpha_d$  (SEQ ID NO: 2) is presented in Figure 1.

- 27 -

The cytoplasmic domains of  $\alpha$  subunits in  $\beta_2$  integrins are typically distinct from one another within the same species, while individual  $\alpha$  subunits show high degrees of homology across species boundaries. Consistent with these observations, the cytoplasmic region of  $\alpha_d$  differs markedly from CD11a, CD11b, and CD11c except for a membrane proximal GFFKR amino acid sequence which has been shown to be conserved among all  $\alpha$  integrins [Rojiani, *et al.*, *Biochemistry* 30: 9859-9866 (1991)]. Since the cytoplasmic tail region of integrins has been implicated in "inside out" signaling and in avidity regulation [Landis *et al.*, *supra*], it is possible that  $\alpha_d$  interacts with cytosolic molecules distinct from those interacting with CD11a, CD11b, and CD11c, and, as a result, participates in signaling pathways distinct from those involving other  $\beta_2$  integrins.

The extracellular domain of  $\alpha_d$  contains a conserved DGSGS amino acid sequence adjacent the I-domain; in CD11b, the DGSGS sequence is a metal-binding region required for ligand interaction [Michishita, *et al.* *Cell* 72:857-867 (1993)]. Three additional putative cation binding sites in CD11b and CD11c are conserved in the  $\alpha_d$  sequence at amino acids 465-474, 518-527, and 592-600 in clone 19A2 (SEQ ID NO: 1). The  $\alpha_d$  I-domain is 36%, 62%, and 57% identical to the corresponding regions in CD11a, CD11b, and CD11c, respectively, and the relatively low sequence homology in this region suggests that  $\alpha_d$  may interact with a set of extracellular proteins distinct from proteins with which other known  $\beta_2$  integrins interact. Alternatively, the affinity of  $\alpha_d$  for known  $\beta_2$  integrin ligands, for example, ICAM-1, ICAM-2 and/or ICAM-R, may be distinct from that demonstrated for the other  $\beta_2$  integrin/ICAM interactions. [See Example 12.]

- 28 -

Example 6Northern Analysis of Human  $\alpha_d$  Expression in Tissues

In order to determine the relative level of expression and tissue specificity of  $\alpha_d$ , Northern analysis was performed using fragments from clone 19A2 as probes. Approximately 10  $\mu$ g of total RNA from each of several human tissues or cultured cell lines were loaded on a formaldehyde agarose gel in the presence of 1  $\mu$ g of ethidium bromide. After electrophoresis at 100 V for 4 hr, the RNA was transferred to a nitrocellulose membrane (Schleicher & Schuell) by wicking in 10X SSC overnight. The membrane was baked 1.5 hr at 80°C under vacuum. Prehybridization solution containing 50% formamide in 3-(N-morpholino)propane sulfonic acid (MOPS) buffer was used to block the membrane for 3 hr at 42°C. Fragments of clone 19A2 were labeled with the Boehringer Mannheim Random Prime kit according to the manufacturer's instructions including both  $\alpha$ P<sup>32</sup>dCTP and  $\alpha$ P<sup>32</sup>dTTP. Unincorporated label was removed on a Sephadex G25 column in TE buffer. The membrane was probed with 1.5 x 10<sup>6</sup> counts per ml of prehybridization buffer. The blot was then washed successively with 2X SSC/0.1% SDS at room temperature, 2X SSC/0.1% SDS at 42°C, 2X SSC/0.1% SDS at 50°C, 1X SSC/0.1% SDS at 50°C, 0.5X SSC/0.1% SDS at 50°C and 0.1X SSC/0.1% SDS at 50°C. The blot was then exposed to film for 19 hr.

Hybridization using a *Bst*XI fragment from clone 19A2 (corresponding to nucleotides 2011 to 3388 in SEQ ID NO: 1) revealed a weak signal in the approximately 5 kb range in liver, placenta, thymus, and tonsil total RNA. No signal was detected in kidney, brain or heart samples. The amount of RNA present in the kidney lane was minimal, as determined with ethidium bromide staining.

When using a second fragment of clone 19A2 (encompassing the region from bases 500 to 2100 in SEQ ID NO: 1), RNA transcripts of two different sizes were detected in a human multi-tissue Northern (MTN) blot using



- 29 -

polyA<sup>+</sup> RNA (Clontech). An approximately 6.5 kb band was observed in spleen and skeletal muscle, while a 4.5 kb band was detected in lung and peripheral blood leukocytes. The variation in sizes observed could be caused by tissue specific polyadenylation, cross reactivity of the probe with other integrin family members, or hybridization with alternatively spliced mRNAs.

Northern analysis using a third fragment from clone 19A2, spanning nucleotides 2000 to 3100 in SEQ ID NO: 1, gave results consistent with those using the other clone 19A2 fragments.

RNA from three myeloid lineage cell lines was also probed using the fragments corresponding to nucleotides 500 to 2100 and 2000 to 3100 in SEQ ID NO:1. A THP-1 cell line, previously stimulated with PMA, gave a diffuse signal in the same size range (approximately 5.0 kb), with a slightly stronger intensity than the tissue signals. RNA from unstimulated and DMSO-stimulated HL-60 cells hybridized with the  $\alpha_d$  probe at the same intensity as the tissue samples, however, PMA treatment seemed to increase the signal intensity. Since PMA and DMSO drive HL-60 cell differentiation toward monocyte/macrophage and granulocyte pathways, respectively, this result suggests enhanced  $\alpha_d$  expression in monocyte/macrophage cell types. U937 cells expressed the  $\alpha_d$  message and this signal did not increase with PMA stimulation. No band was detected in Molt, Daudi, H9, JY, or Jurkat cells.

### Example 7

#### Transient Expression of Human $\alpha_d$ Constructs

##### A. Generation of expression constructs

The human clone 19A2 lacks an initiating methionine codon and possibly some of the 5' signal sequence. Therefore, in order to generate a human expression plasmid containing 19A2 sequences, two different strategies were used. In the first, two plasmids were constructed in which signal peptide sequences derived from genes encoding either CD11b or CD11c were spliced into clone

- 30 -

19A2 to generate a chimeric  $\alpha_d$  sequence. In the second approach, a third plasmid was constructed in which an adenosine base was added at position 0 in clone 19A2 to encode an initiating methionine.

The three plasmids contained different regions which encoded the 5' portion of the  $\alpha_d$  sequence or the chimeric  $\alpha_d$  sequence. The  $\alpha_d$  region was PCR amplified (see conditions in Example 2) with a specific 3' primer BamRev (set out below in SEQ ID NO: 26) and one of three 5' primers. The three 5' primers contained in sequence: (1) identical nonspecific bases at positions 1-6 allowing for digestion, an *Eco*RI site from positions 7-12 and a consensus Kozak sequence from positions 13-18; (2) a portion of the CD11b (primer ER1B) or CD11c (primer ER1C) signal sequence, or an adenosine (primer ER1D); and (3) an additional 15-17 bases specifically overlapping 5' sequences from clone 19A2 to allow primer annealing. Primers ER1B, ER1C or ER1D are set out in SEQ ID NOS: 27, 28 or 29, respectively, where the initiating methionine codon is underlined and the *Eco*RI site is double underlined.

5'-CCACTGTCAGGATGCCCCGTG-3' (SEQ ID NO: 26)

5'-AGTTACGAATTCGCCACCATGGCTCTACGGGTGCTT (SEQ ID NO: 27)

20 5'-AGTTACGAATTCGCCACCATGACTCGGACTGTGCTT (SEQ ID NO: 28)

5'-AGTTACGAATTCGCCACCATGACCTTCGGCACTGTG (SEQ ID NO: 29)

The resulting PCR product was digested with *Eco*RI and *Bam*HI.

25 All three plasmids contained a common second  $\alpha_d$  region (to be inserted immediately downstream from the 5' region described in the previous paragraph) including the 3' end of the  $\alpha_d$  clone. The second  $\alpha_d$  region, which

- 31 -

extended from nucleotide 625 into the *Xba*I site in the vector 3' polylinker region of clone 19A2, was isolated by digestion of clone 19A2 with *Bam*HI and *Xba*I.

Three ligation reactions were prepared in which the 3'  $\alpha_d$  *Bam*HI/*Xba*I fragment was ligated to one of the three 5'  $\alpha_d$  *Eco*RI/*Bam*HI fragments using Boehringer Mannheim ligase buffer and T4 ligase (1 unit per reaction). After a 4 hour incubation at 14°C, an appropriate amount of vector pcDNA.3 (Invitrogen) digested with *Eco*RI and *Xba*I was added to each reaction with an additional unit of ligase. Reactions were allowed to continue for another 14 hours. One tenth of the reaction mixture was then transformed into competent XL-1 Blue cells. The resulting colonies were cultured and the DNA isolated as in Example 5. Digestion with *Eco*RI identified three clones which were positive for that restriction site, and thus, the engineered signal sequences. The clones were designated pATM.B1 (CD11b/ $\alpha_d$ , from primer ER1B), pATM.C10 (CD11c/ $\alpha_d$ , from primer ER1C) and pATM.D12 (adenosine/ $\alpha_d$  from primer ER1d). The presence of the appropriate signal sequences in each clone was verified by nucleic acid sequencing.

#### B. Transfection of COS Cells

Expression from the  $\alpha_d$  plasmids discussed above was effected by cotransfection of COS cells with the individual plasmids and a CD18 expression plasmid, pRC.CD18. As a positive control, COS cells were also co-transfected with the plasmid pRC.CD18 and a CD11a expression plasmid, pDC.CD11A.

Cells were passaged in culture medium (DMEM/10%FBS/pen-strep) into 10 cm Corning tissue culture-treated petri dishes at 50% confluency 16 hours prior to transfection. Cells were removed from the plates with Versene buffer (0.5 mM NaEDTA in PBS) without trypsin for all procedures. Before transfection, the plates were washed once with serum-free DMEM. Fifteen micrograms of each plasmid were added to 5 ml transfection buffer (DMEM with 20  $\mu$ g/ml DEAE-Dextran and 0.5 mM chloroquine) on each plate. After 1.5

- 32 -

hours incubation at 37°C, the cells were shocked for 1 minute with 5 ml DMEM/10% DMSO. This DMSO solution was then replaced with 10 ml/plate culture medium.

Resulting transfectants were analyzed by ELISA, FACS, and immunoprecipitation as described in Examples 8, 9, and 10.

### Example 8

#### ELISA Analysis of COS Transfectants

In order to determine if the COS cells co-transfected with CD18 expression plasmid pRC.CD18 and an  $\alpha_d$  plasmid expressed  $\alpha_d$  on the cell surface in association with CD18, ELISAs were performed using primary antibodies raised against CD18 (*e.g.*, TS1/18 purified from ATCC HB203). As a positive control, ELISAs were also performed on cells co-transfected with the CD18 expression plasmid and a CD11a expression plasmid, pDC.CD11A. The primary antibodies in this control included CD18 antibodies and anti-CD11a antibodies (*e.g.*, TS1/22 purified from ATCC HB202).

For ELISA, cells from each plate were removed with Versene buffer and transferred to a single 96-well flat-bottomed Corning tissue culture plate. Cells were allowed to incubate in culture media 2 days prior to assay. The plates were then washed twice with 150  $\mu$ l/well D-PBS/0.5% teleost skin gelatin (Sigma) solution. This buffer was used in all steps except during the development. All washes and incubations were performed at room temperature. The wells were blocked with gelatin solution for 1 hour. Primary antibodies were diluted to 10  $\mu$ g/ml in gelatin solution and 50  $\mu$ l were then added to each well. Triplicate wells were set up for each primary antibody. After 1 hour incubation, plates were washed 3X with 150  $\mu$ l/well gelatin solution. Secondary antibody (goat anti-mouse Ig/HRP-Fc specific [Jackson, West Grove, PA]) at a 1:3500 dilution was added at 50  $\mu$ l/well and plates were incubated for 1 hour. After three washes, plates were developed for 20 minutes with 100  $\mu$ l/well o-

- 33 -

phenyldiamine (OPD) (Sigma) solution (1 mg/ml OPD in citrate buffer) before addition of 50  $\mu$ l/well 15% sulfuric acid.

Analysis of transfectants in the ELISA format with anti-CD18 specific antibodies revealed no significant expression above background in cells transfected only with the plasmid encoding CD18. Cells co-transfected with plasmid containing CD11a and CD18 showed an increase in expression over background when analyzed with CD18 specific antibodies or with reagents specific for CD11a. Further analysis of cells co-transfected with plasmids encoding CD18 and one of the  $\alpha_d$  expression constructs (pATM.C10 or pATM.D12) revealed that cell surface expression of CD18 was rescued by concomitant expression of  $\alpha_d$ . The increase in detectable CD18 expression in COS cells transfected with pATM.C10 or pATM.D12 was comparable to that observed in co-transfected CD11a/CD18 positive control cells.

#### Example 9

##### FACS Analysis of COS Transfectants

For FACS analysis, cells in petri dishes were fed with fresh culture medium the day after transfection and allowed to incubate 2 days prior to the assay. Transfectant cells were removed from the plates with 3 ml Versene, washed once with 5 ml FACS buffer (DMEM/2% FBS/0.2% sodium azide) and diluted to 500,000 cells/sample in 0.1 ml FACS buffer. Ten microliters of either 1 mg/ml FITC-conjugated CD18, CD11a, or CD11b specific antibodies (Becton Dickinson) or 800  $\mu$ g/ml CFSE-conjugated murine 23F2G (anti-CD18) (ATCC HB11081) were added to each sample. Samples were then incubated on ice for 45 minutes, washed 3X with 5 ml/wash FACS buffer and resuspended in 0.2 ml FACS buffer. Samples were processed on a Becton Dickinson FACscan and the data analyzed using Lysys II software (Becton Dickinson).

COS cells transfected with CD18 sequences only did not stain for CD18, CD11a or CD11b. When co-transfected with CD11a/CD18, about 15%

- 34 -

of the cells stained with antibodies to CD11a or CD18. All cells transfected with CD18 and any  $\alpha_d$  construct resulted in no detectable staining for CD11a and CD11b. The pATM.B1, pATM.C10 and pATM.D12 groups stained 4%, 13% and 8% positive for CD18, respectively. Fluorescence of the positive population in the CD11a/CD18 group was 4-fold higher than background. In comparison, the co-transfection of  $\alpha_d$  constructs with the CD18 construct produced a positive population that showed a 4- to 7-fold increase in fluorescence intensity over background.

#### Example 10

##### Biotin-Labeled Immunoprecipitation of Human $\alpha_d$ /CD18 Complexes from Co-transfected COS Cells

Immunoprecipitation was attempted on cells co-transfected with CD18 and each of the  $\alpha_d$  expression plasmids separately described in Example 7 in order to determine if  $\alpha_d$  could be isolated as part of the  $\alpha\beta$  heterodimer complex characteristic of integrins.

Transfected cells ( $1-3 \times 10^8$  cells/group) were removed from petri dishes with Versene buffer and washed 3 times in 50 ml/group D-PBS. Each sample was labeled with 2 mg Sulpho-NHS Biotin (Pierce, Rockford, IL) for 15 minutes at room temperature. The reaction was quenched by washing 3 times in 50 ml/sample cold D-PBS. Washed cells were resuspended in 1 ml lysis buffer (1% NP40, 50 mM Tris-HCl, pH 8.0, 0.2 M NaCl, 2 mM  $\text{Ca}^{++}$ , 2 mM  $\text{Mg}^{++}$ , and protease inhibitors) and incubated 15 minutes on ice. Insoluble material was pelleted by centrifugation at 10,000 g for 5 minutes, and the supernatant removed to fresh tubes. In order to remove material non-specifically reactive with mouse immunoglobulin, a pre-clearance step was initially performed. Twenty-five micrograms of mouse immunoglobulin (Cappel, West Chester, PA) was incubated with supernatants at 4°C. After 2.5 hr, 100  $\mu\text{l}$  (25  $\mu\text{g}$ ) rabbit anti-mouse Ig conjugated Sepharose (prepared from Protein A Sepharose 4B and rabbit anti-mouse IgG, both from Zymed, San Francisco, CA) was added to each sample;

- 35 -

incubation was continued at 4°C with rocking for 16 hours. Sepharose beads were removed from the supernatants by centrifugation. After pre-clearance, the supernatants were then treated with 20 µg anti-CD18 antibody (TS1.18) for 2 hours at 4°C. Antibody/antigen complexes were isolated from supernatants by incubation with 100 µl/sample rabbit anti-mouse/Protein A-sepharose preparation described above. Beads were washed 4 times with 10 mM HEPES, 0.2 M NaCl, and 1% Triton-X 100. Washed beads were pelleted and boiled for 10 minutes in 20 µl 2X Laemmli sample buffer with 2% β-mercaptoethanol. Samples were centrifuged and run on an 8% prepoured Novex polyacrylamide gel (Novex) at 100 V for 30 minutes. Protein was transferred to nitrocellulose membranes (Schleicher & Schuell) in TBS-T buffer at 200 mAmps for 1 hour. Membranes were blocked for 2 hr with 3% BSA in TBS-T. Membranes were treated with 1:6000 dilution of Strep-avidin horse radish peroxidase (POD) (Boehringer Mannheim) for 1 hour, followed by 3 washes in TBS-T. The Amersham Enhanced Chemiluminescence kit was then used according to the manufacturer's instructions to develop the blot. The membrane was exposed to Hyperfilm MP (Amersham) for 0.5 to 2 minutes.

Immunoprecipitation of CD18 complexes from cells transfected with pRC.CD18 and either pATM.B1, pATM.C10 or pATM.D12 revealed surface expression of a heterodimeric species consisting of approximately 100 kD β chain, consistent with the predicted size of CD18, and an α chain of approximately 150 kD, corresponding to α<sub>d</sub>.

### Example 11

#### Stable Transfection of Human α<sub>d</sub> in Chinese Hamster Ovary Cells

To determine whether α<sub>d</sub> is expressed on the cell surface as a heterodimer in association with CD18, cDNAs encoding each chain were both transiently and stably transfected into a cell line lacking both α<sub>d</sub> and CD18.

- 36 -

For these experiments,  $\alpha_d$  cDNA was augmented with additional leader sequences and a Kozak consensus sequence, as described in Example 7, and subcloned into expression vector pcDNA3. The final construct, designated pATM.D12, was co-transfected with a modified commercial vector, pDC1.CD18 encoding human CD18 into dihydrofolate reductase (DHFR)<sup>-</sup> Chinese hamster ovary (CHO) cells. The plasmid pDC1.CD18 encodes a DHFR<sup>+</sup> marker and transfectants can be selected using an appropriate nucleoside-deficient medium. The modifications which resulted in pDC1.CD18 are as follows.

The plasmid pRC/CMV (Invitrogen) is a mammalian expression vector with a cytomegalovirus promoter and ampicillin resistance marker gene. A DHFR gene from the plasmid pSC1190-DHFR was inserted into pRC/CMV 5' of the SV40 origin of replication. In addition, a polylinker from the 5' region of the plasmid pHF2G-DHF was ligated into the pRC/CMV/DHFR construct, 3' to the DHFR gene. CD18 encoding sequences are subsequently cloned into the resulting plasmid between the 5' flanking polylinker region and the bovine growth hormone poly A encoding region.

Surface expression of CD18 was analyzed by flow cytometry using the monoclonal antibody TS1/18. Heterodimer formation detected between  $\alpha_d$  and CD18 in this cell line was consistent with the immunoprecipitation described in Example 10 with transient expression in COS cells.

### Example 12

#### Human $\alpha_d$ binds to ICAM-R in a CD18-dependent fashion

In view of reports that demonstrate interactions between the leukocyte integrins and intercellular adhesion molecules (ICAMs) which mediate cell-cell contact [Hynes, *Cell* 69:11-25 (1992)], the ability of CHO cells expressing  $\alpha_d$ /CD18 to bind ICAM-1, ICAM-R, or VCAM-1 was assessed by two methods.



- 37 -

In replicate assays, soluble ICAM-1, ICAM-R, or VCAM-1 IgG1 fusion proteins were immobilized on plastic and the ability of  $\alpha_d$ /CD18 CHO transfected cells to bind the immobilized ligand was determined. Transfected cells were labeled internally with calcein, washed in binding buffer (RPMI with 1% BSA), and incubated in either buffer only (with or without 10 ng/ml PMA) or buffer with anti-CD18 monoclonal antibodies at 10  $\mu$ g/ml. Transfected cells were added to 96-well Immulon 4 microtiter plates previously coated with soluble ICAM-1/IgG1, ICAM-R/IgG1 or VCAM-1/IgG1 fusion protein, or bovine serum albumin (BSA) as a negative control. Design of the soluble forms of these adhesion molecules is described and fully disclosed in co-pending and co-owned U.S. Patent Application Serial No. 08/102,852, filed August 5, 1993. Wells were blocked with 1% BSA in PBS prior to addition of labeled cells. After washing the plates by immersion in PBS with 0.1% BSA for 20 minutes, total fluorescence remaining in each well was measured using a Cytofluor 2300 (Millipore, Milford, MA).

In experiments with immobilized ICAMs,  $\alpha_d$ /CD18 co-transfectants consistently showed a 3-5 fold increase in binding to ICAM-R/IgG1 wells over BSA coated wells. The specificity and CD18-dependence of this binding was demonstrated by the inhibitory effects of anti-CD18 antibody TS1/18. The binding of cells transfected with CD11a/CD18 to ICAM-1/IgG1 wells was comparable to the binding observed with BSA coated wells. CD11a/CD18 transfected cells showed a 2-3 fold increase in binding to ICAM-1/IgG1 wells only following pretreatment with PMA. PMA treatment of  $\alpha_d$ /CD18 transfectants did not affect binding to ICAM-1/IgG1 or ICAM-R/IgG1 wells. No detectable binding of  $\alpha_d$ /CD18 transfectants to VCAM-1/IgG1 wells was observed.

Binding of  $\alpha_d$ /CD18-transfected cells to soluble ICAM-1/IgG1, ICAM-R/IgG1, or VCAM-1/IgG1 fusion proteins was determined by flow cytometry. Approximately one million  $\alpha_d$ /CD18-transfected CHO cells (grown in spinner flasks for higher expression) per measurement were suspended in 100  $\mu$ l

- 38 -

binding buffer (RPMI and 1% BSA) with or without 10  $\mu$ g/ml anti-CD18 antibody. After a 20 minute incubation at room temperature, the cells were washed in binding buffer and soluble ICAM-1/IgG1 or ICAM-R/IgG1 fusion protein was added to a final concentration of 5  $\mu$ g/ml. Binding was allowed to  
5 proceed for 30 minute at 37°C, after which the cells were washed three times and resuspended in 100  $\mu$ l binding buffer containing FITC-conjugated sheep anti-human IgG1 at a 1:100 dilution. After a 30 minute incubation, samples were washed three times and suspended in 200  $\mu$ l binding buffer for analysis with a Becton Dickinson FACScan.

10 Approximately 40-50% of the  $\alpha_d$ /CD18 transfectants indicated binding to ICAM-R/IgG1, but no binding to ICAM-1/IgG1 or VCAM-1/IgG1 proteins. Pretreatment of transfected cells with PMA has no effect on  $\alpha_d$ /CD18 binding to either ICAM-1/IgG1, ICAM-R/IgG1 or VCAM-1/IgG1, which was consistent with the immobilized adhesion assay. Binding by ICAM-R was  
15 reduced to background levels after treatment of  $\alpha_d$ /CD18 transfectants with anti-CD18 antibody TS1/18.

The collective data from these two binding assays illustrate that  $\alpha_d$ /CD18 binds to ICAM-R and does so preferentially as compared to ICAM-1 and VCAM-1. The  $\alpha_d$ /CD18 binding preference for ICAM-R over ICAM-1 is  
20 opposite that observed with CD11a/CD18 and CD11b/CD18. Thus modulation of  $\alpha_d$ /CD18 binding may be expected to selectively affect normal and pathologic immune function where ICAM-R plays a prominent role. Moreover, results of similar assays, in which antibodies immunospecific for various extracellular domains of ICAM-R were tested for their ability to inhibit binding of ICAM-R  
25 to  $\alpha_d$ /CD18 transfectants, indicated that  $\alpha_d$ /CD18 and CD11a/CD18 interact with different domains of ICAM-R.

The failure of CD11a/CD18 to bind ICAM-1/IgG1 or ICAM-R/IgG1 in solution suggests that the affinity of binding between CD11a/CD18 and ICAM-1 or ICAM-R is too low to permit binding in solution. Detection of

- 39 -

$\alpha_d$ /CD18 binding to ICAM-R/IgG1, however, suggests an unusually high binding affinity.

#### $\alpha_d$ Binding to iC3b

Complement component C3 can be proteolytically cleaved to form the complex iC3b, which initiates the alternative pathway of complement activation and leads ultimately to cell-mediated destruction of a target. Both CD11b and CD11c have been implicated in iC3b binding and subsequent phagocytosis of iC3b-coated particles. A peptide fragment in the CD11b I domain has recently been identified as the site of iC3b interaction [Ueda, *et al.*, *Proc.Natl.Acad.Sci.(USA)* 91:10680-10684 (1994)]. The region of iC3b binding is highly conserved in CD11b, CD11c, and  $\alpha_d$ , suggesting an  $\alpha_d$ /iC3b binding interaction.

Binding of  $\alpha_d$  to iC3b is performed using transfectants or cell lines naturally expressing  $\alpha_d$  (for example, PMA-stimulated HL60 cells) and iC3b-coated sheep red blood cells (sRBC) in a rosette assay [Dana, *et al.*, *J.Clin.Invest.* 73:153-159 (1984)]. The abilities of  $\alpha_d$ /CD18 CHO transfectants, VLA4-CHO transfectants (negative control) and PMA-stimulated HL60 cells (positive control) to form rosettes are compared in the presence and absence of an anti-CD18 monoclonal antibody (for example TS1/18.1).

#### 20 Example 13

##### Screening by Scintillation Proximity Assay

Specific inhibitors of binding between the  $\alpha_d$  ligands of the present invention and their binding partners ( $\alpha_d$  ligand/anti-ligand pair) may be determined by a variety of means, such as scintillation proximity assay techniques as generally described in U.S. Patent No. 4,271,139, Hart and Greenwald, *Mol.Immunol.* 12:265-267 (1979), and Hart and Greenwald, *J.Nuc.Med.* 20:1062-1065 (1979), each of which is incorporated herein by reference.

- 40 -

Briefly, one member of the  $\alpha_d$  ligand/anti-ligand pair is bound to a solid support. A fluorescent agent is also bound to the support. Alternatively, the fluorescent agent may be integrated into the solid support as described in U.S. Patent No. 4,568,649, incorporated herein by reference. The non-support bound member of the  $\alpha_d$  ligand/anti-ligand pair is labeled with a radioactive compound that emits radiation capable of exciting the fluorescent agent. When the ligand binds the radiolabeled anti-ligand, the label is brought sufficiently close to the support-bound fluorescer to excite the fluorescer and cause emission of light. When not bound, the label is generally too distant from the solid support to excite the fluorescent agent, and light emissions are low. The emitted light is measured and correlated with binding between the ligand and the anti-ligand. Addition of a binding inhibitor to the sample will decrease the fluorescent emission by keeping the radioactive label from being captured in the proximity of the solid support. Therefore, binding inhibitors may be identified by their effect on fluorescent emissions from the samples. Potential anti-ligands to  $\alpha_d$  may also be identified by similar means.

#### Example 14

##### Soluble Human $\alpha_d$ Expression Constructs

The expression of full-length, soluble human  $\alpha_d$ /CD18 heterodimeric protein provides easily purified material for immunization and binding assays. The advantage of generating soluble protein is that it can be purified from supernatants rather than from cell lysates (as with full-length membrane-bound  $\alpha_d$ /CD18); recovery is therefore improved and impurities reduced.

The soluble  $\alpha_d$  expression plasmid was constructed as follows. A nucleotide fragment corresponding to the region from bases 0 to 3161 in SEQ ID NO: 1, cloned into plasmid pATM.D12, was isolated by digestion with *HindIII* and *AatII*. A PCR fragment corresponding to bases 3130 to 3390 in SEQ ID NO:

- 41 -

1, overlapping the *HindIII/AatII* fragment and containing an addition *MluI* restriction site at the 3' terminus, was amplified from pATM.D12 with primers sHAD.5 and sHAD.3 set out in SEQ ID NOS: 30 and 31, respectively.

5'-TTGCTGACTGCCTGCAGTTC-3' (SEQ ID NO: 30)

5'-GTTCTGACGCGTAATGGCATTGTAGACCTCGTCTTC (SEQ ID NO: 31)

The PCR amplification product was digested with *AatII* and *MluI* and ligated to the *HindIII/AatII* fragment. The resulting product was ligated into *HindIII/MluI*-digested plasmid pDC1.s.

This construct is co-expressed with soluble CD18 in stably transfected CHO cells, and expression is detected by autoradiographic visualization of immunoprecipitated CD18 complexes derived from <sup>35</sup>S-methionine labeled cells. The construct is also co-expressed with CD18 in 293 cells [Berman, *et al.*, *J. Cell. Biochem.* 52:183-195 (1993)].

#### Soluble Human $\alpha_d$ I Domain Expression Constructs

It has previously been reported that the I domain in CD11a can be expressed as an independent structural unit that maintains ligand binding capabilities and antibody recognition [Randi and Hogg, *J. Biol. Chem.* 269:12395-12398 (1994); Zhout, *et al.*, *J. Biol. Chem.* 269:17075-17079 (1994); Michishita, *et al.*, *Cell* 72:857-867 (1993)]. To generate a soluble fusion protein comprising the  $\alpha_d$  I domain and human IgG4, the  $\alpha_d$  I domain is amplified by PCR using primers designed to add flanking *Bam*HI and *Xho*I restriction sites to facilitate subcloning. These primers are set out in SEQ ID NOS: 32 and 33 with restriction sites underlined.

5'-ACGTATGCAGGATCCCATCAAGAGATGGACATCGCT (SEQ ID NO: 32)

5'-ACTGCATGTCTCGAGGCTGAAGCCTTCTTGGGACAT (SEQ ID NO: 33)

- 42 -

The C nucleotide immediately 3' to the *Bam*HI site in SEQ ID NO: 32 corresponds to nucleotide 435 in SEQ ID NO: 1; the G nucleotide 3' to the *Xho*I site in SEQ ID NO: 33 is complementary to nucleotide 1067 in SEQ ID NO: 1. The amplified I domain is digested with the appropriate enzymes, the purified  
5 fragment ligated into the mammalian expression vector pDCs and the prokaryotic expression vector pGEX-4T-3 (Pharmacia) and the I domain fragment sequenced. The fusion protein is then expressed in COS, CHO or *E.coli* cells transfected or transformed with an appropriate expression construct.

Given the affinity of  $\alpha_d$  for ICAM-R, expression of the  $\alpha_d$  I  
10 domain may be of sufficient affinity to be a useful inhibitor of cell adhesion in which  $\alpha_d$  participates.

#### Analysis of Human $\alpha_d$ I Domain/IgG4 Fusion Proteins

Protein was resolved by SDS-PAGE under reducing and non-reducing conditions and visualized by either silver staining or Coomassie staining.  
15 Protein was then transferred to Immobilon PVDF membranes and subjected to Western blot analysis using anti-human IgG monoclonal antibodies or anti-bovine Ig monoclonal antibodies.

Protein detected was determined to migrate at about 120 kD under non-reducing conditions and at about 45 kD under reducing conditions. Minor  
20 bands were also detected on non-reducing gels at approximately 40-50 kD which were reactive with the anti-human, but not anti-bovine, antibodies. A 200 kD minor band was determined to be bovine Ig by Western blot.

#### Binding Assays Using I Domain Expression Products

The ability of the I domain to specifically recognize ICAM-R/IgG  
25 chimeric protein was tested in an ELISA format. Serial dilutions of  $\alpha_d$  I domain IgG4 fusion protein ( $I\alpha_d$ /IgG4) in TBS were incubated with ICAM-1/IgG, ICAM-R/IgG, VCAM-1/IgG, or an irrelevant IgG1 myeloma protein

- 43 -

immobilized on Immulon IV RIA/EIA plates. CD11a I domain/IgG chimeric protein and human IgG4/kappa myeloma protein were used as negative controls. Bound IgG4 was detected with the biotinylated anti-IgG4 monoclonal antibody HP6023 followed by addition of streptavidin-peroxidase conjugate and development with substrate *o*-phenyldiamine.

In repeated assays, no binding of the CD11a/IgG4 protein or the IgG4 myeloma protein was detected with any of the immobilized proteins. The  $I\alpha_d$ /IgG4 protein did not bind to fish skin gelatin or bovine serum albumin blocking agents, human IgG1, or ICAM-1/IgG. A two to three fold increase in binding signal over background was detected in ICAM-R/IgG protein coated wells using 1-5  $\mu$ g/ml concentrations of  $I\alpha_d$ /IgG4 protein. The signal in VCAM-1/IgG protein coated wells was 7-10 fold higher than background. In previous assays,  $\alpha_d$ /CD18 transfected CHO cells did not bind VCAM-1/IgG protein, suggesting that VCAM-1 binding may be characteristic of isolated I domain amino acid sequences.

#### Additional $\alpha_d$ I domain constructs

Additional  $\alpha_d$  I domain constructs are generated in the same fashion as the previous construct, but incorporating more amino acids around the  $\alpha_d$  I domain. Specific constructs include: i) sequences from exon 5 (amino acids 127-353 in SEQ ID NO: 2), preceding the current construct, ii) the EF-hand repeats (amino acids 17-603 in SEQ ID NO: 2) following the I domain, and iii) the alpha chain truncated at the transmembrane region (amino acids 17-1029 in SEQ ID NO: 2), with an IgG4 tail for purification and detection purposes. These constructs are ligated into either the mammalian expression vector pDCS1 or the prokaryotic expression vector pGEX-4T-3 (Pharmacia) and the I domain sequenced. The fusion proteins are then be expressed in COS, CHO, or *E.coli* cells transformed or transfected with an appropriate expression construct. Protein are purified on a ProSepA column (Bioprocessing Limited, Durham, England),

- 44 -

tested for reactivity with the anti-IgG4 monoclonal antibody HP6023 and visualized on polyacrylamide gels with Coomassie staining.

5 In order to construct an expression plasmid for the entire  $\alpha_d$  polypeptide, pATM.D12, described *supra*, is modified to express an  $\alpha_d$ -IgG4 fusion protein by the following method. IgG4 encoding DNA is isolated from the vector pDCS1 by PCR using primers which individually incorporate a 5' *Aat*II restriction site (SEQ ID NO: 89) and a 3' *Xba*I restriction site (SEQ ID NO: 90).

5'-CGCTGTGACGTCAGAGTTGAGTCCAAATATGG-3' (SEQ ID NO: 89)

5'-GGTGACACTATAGAATAGGGC-3' (SEQ ID NO: 90)

10 Plasmid pATM.D12 is digested with *Aat*II and *Xba*I, and the appropriately digested and purified IgG4 PCR product ligated into the linear vector.



- 45 -

**Example 15****Production of Human  $\alpha_d$ -Specific Monoclonal Antibodies**

Transiently transfected cells from Example 7 were washed three times in Dulbecco's phosphate buffered saline (D-PBS) and injected at  $5 \times 10^6$  cells/mouse into Balb/c mice with 50  $\mu$ g/mouse muramyl dipeptidase (Sigma) in PBS. Mice were injected two more times in the same fashion at two week intervals. The pre-bleed and immunized serum from the mice were screened by FACS analysis as outlined in Example 9 and the spleen from the mouse with the highest reactivity to cells transfected with  $\alpha_d$ /CD18 was fused. Hybridoma culture supernatants were then screened separately for lack of reactivity against COS cells transfected with CD11a/CD18 and for reactivity with cells co-transfected with an  $\alpha_d$  expression plasmid and CD18.

This method resulted in no monoclonal antibodies.

As an alternative for production of monoclonal antibodies, soluble  $\alpha_d$  I domain IgG4 fusion protein was affinity purified from supernatant of stably transfected CHO cells and used to immunize Balb/c mice as described above. Hybridomas were established and supernatants from these hybridomas were screened by ELISA for reactivity against  $\alpha_d$  I domain fusion protein. Positive cultures were then analyzed for reactivity with full length  $\alpha_d$ /CD18 complexes expressed on CHO transfectants.

Mouse 1908 received three initial immunizations of  $\alpha_d$ /CD18 transfected CHO cells and two subsequent boosts with soluble  $\alpha_d$ /CD18 heterodimer. Two final immunizations included 50  $\mu$ g/mouse I $\alpha_d$ /IgG4 fusion protein. The fusion produced 270 IgG-producing wells. Supernatant from 45 wells showed at least 7-fold higher binding to I $\alpha_d$ /IgG4 fusion protein than to human IgG4 by ELISA. None of the supernatants reacted to  $\alpha_d$ /CD18 transfected CHO cells as determined by FACS analysis.

To determine whether the supernatants were able to recognize integrin alpha subunit proteins in another context, fresh frozen splenic sections

- 46 -

were stained with supernatants from 24 of the 45 wells. Three supernatants were determined to be positive: one stained large cells in the red pulp, while two others stained scattered cells in the red pulp and also trabeculae.

5           These supernatants were further analyzed by their ability to immunoprecipitate biotinylated CD18 complexes from either  $\alpha_d$ /CD18 transfected CHO cells or PMA-stimulated HL60 cells. Fusion wells with supernatants that recognized protein in detergent lysates (which should not be as conformationally constrained as protein expressed as heterodimers) were selected for further subcloning. Monoclonal antibodies which recognize protein in detergent may be  
10       more useful in immunoprecipitation of heterodimeric complexes from transfectants, tissues, and cell lines.

          As another alternative, monoclonal antibodies are generated as follows. Affinity purified  $\alpha_d$ /CD18 heterodimeric protein from detergent lysates of stably transfected CHO cells is used with 50  $\mu$ g/ml muramyl dipeptidase to  
15       immunize Balb/c mice as described above. Mice receive three immunizations before serum reactivity against  $\alpha_d$ /CD18 is determined by immunoprecipitation of biotinylated complexes in the CHO transfectants. Hybridomas from positive animals are established according to standard protocols, after which hybridoma cultures are selected by flow cytometry using  $\alpha_d$ /CD18 transfectants.  
20       CD11a/CD18 transfectants are utilized to control for CD18-only reactivity.

          As another alternative for monoclonal antibody production, Balb/c mice undergo an immunization/immunosuppression protocol designed to reduce reactivity to CHO cell determinants on transfectants used for immunization. This protocol involves immunization with untransfected CHO cells and subsequent  
25       killing of CHO-reactive B-cell blasts with cyclophosphamide treatment. After three rounds of immunization and cyclophosphamide treatment are performed, the mice are immunized with  $\alpha_d$ /CD18 CHO transfected cells as described above.

          As still another alternative, heterodimeric CD18 complexes are immunoprecipitated from detergent lysates of whole spleen using an anti-CD18

- 47 -

monoclonal antibody, following preclearance of CD11a/CD18 and CD11b/CD18. CD11a/CD18 and CD11b/CD18 complexes are precleared by affinity chromatography using monoclonal antibodies TS2/4 and Mo1, respectively, coupled to a chromatographic resin. The remaining CD18 complexes are used as an immunogen in Balb/c mice for the first immunization. Three immunizations are given at three week intervals, the initial immunization administered in conjunction with Freund's Complete Adjuvant and the subsequent immunizations with Freund's Incomplete Adjuvant. Serum is assayed for  $\alpha_d$ -specific reactivity by immunoprecipitation. Resulting hybridomas are screened by flow cytometry with  $\alpha_d$ /CD18 CHO transfectants.

As another alternative, CD18 complexes from detergent lysates of PMA stimulated HL60 cells are enriched by preclearance as described above. Other  $\beta 2$  integrins are cleared on the same columns. Immunization with the resulting complexes, hybridoma production, and screening protocols are performed as described *supra*.

#### Example 16

##### Analysis of $\alpha_d$ distribution with polyclonal serum

Tissue distribution of  $\alpha_d$ /CD18 was determined using polyclonal antiserum. Antiserum used to stain tissue was obtained from a mouse immunized 3 times with  $\alpha_d$  transfected CHO cells (D6.CHO,  $\alpha_d$ /CD18) with adjuvant peptide and once with purified  $\alpha_d$ /CD18 heterodimer. A final boost included only  $\alpha_d$ /CD18 heterodimer. Approximately 100  $\mu$ l immunized serum was precleared by addition of approximately  $10^8$  LFA-1-transfected CHO cells for 2 hours at 4°C. The resulting serum was assayed for  $\alpha_d$  reactivity at dilutions of 1/5000, 1/10000, 1/20000 and 1/40000 on normal human spleen. The polyclonal antibody was reactive at a dilution of 1/20000, while a 1/40000 dilution stained very weakly.

- 48 -

Once serum was determined to have specific  $\alpha_d$  reactivity, it was used to stain various lymphoid and non-lymphoid tissues. Monoclonal antibodies recognizing CD18, CD11a, CD11b, and CD11c were used in the same experiment as controls. Staining of normal spleen sections with  $\alpha_d$  polyclonal sera, and monoclonal antibodies to CD11a, CD11b, CD11c, and CD18 revealed the following results. The pattern observed with  $\alpha_d$  polyclonal sera did not display the same pattern of labeling as CD11a, CD11b, CD11c, or CD18. There is a distinct pattern of labeling with some cells located in the marginal zone of the white pulp and a distinct labeling of cells peripheral to the marginal zone. This pattern was not observed with the other antibodies. Individual cells scattered throughout the red pulp were also labeled which may or may not be the same population or subset seen with CD11a and CD18.

Labeling with CD11c did display some cells staining in the marginal zone, but the antibody did not show the distinct ring pattern around the white pulp when compared to  $\alpha_d$  polyclonal sera, nor did labeling in the red pulp give the same pattern of staining as  $\alpha_d$  polyclonal sera.

Therefore, the labeling pattern seen with  $\alpha_d$  polyclonal serum was unique compared to that seen using antibodies to the other  $\beta_2$  integrins (CD11a, CD11b, CD11c, and CD18), and suggests that the *in vivo* distribution of  $\alpha_d$  in man is distinct from that of other  $\beta_2$  integrins.

### Example 17

#### Isolation of Rat cDNA Clones

In view of the existence of both canine and human  $\alpha_d$  subunits, attempts were made to isolate homologous genes in other species, including rat (this example) and mouse (Example 17, *infra*).

A partial sequence of a rat cDNA showing homology to the human  $\alpha_d$  gene was obtained from a rat splenic  $\lambda$ gt10 library (Clontech). The library was plated at  $2 \times 10^4$  pfu/plate onto 150 mm LBM/agar plates. The library was

- 49 -

lifted onto Hybond membranes (Amersham), denatured 3 minutes, neutralized 3 minutes and washed 5 minutes with buffers as described in standard protocols [Sambrook, *et al.*, Molecular Cloning: a laboratory manual, p.2.110]. The membranes were placed immediately into a Stratalinker (Stratagene) and the DNA crosslinked using the autocrosslinking setting. The membranes were prehybridized and hybridized in 30% or 50% formamide, for low and high stringency conditions, respectively. Membranes were initially screened with a <sup>32</sup>P-labeled probe generated from the human  $\alpha_d$  cDNA, corresponding to bases 500 to 2100 in clone 19A2 (SEQ ID NO: 1). The probe was labeled using Boehringer Mannheim's Random Prime Kit according to manufacturer's suggested protocol. Filters were washed with 2X SSC at 55°C.

Two clones, designated 684.3 and 705.1, were identified which showed sequence homology to human  $\alpha_d$ , human CD11b, and human CD11c. Both clones aligned to the human  $\alpha_d$  gene in the 3' region of the gene, starting at base 1871 and extending to base 3012 for clone 684.3, and bases 1551 to 3367 for clone 705.1.

In order to isolate a more complete rat sequence which included the 5' region, the same library was rescreened using the same protocol as employed for the initial screening, but using a mouse probe generated from clone A1160 (See Example 17, *infra*). Single, isolated plaques were selected from the second screening and maintained as single clones on LBM/agar plates. Sequencing primers 434FL and 434FR (SEQ ID NOS: 34 and 35, respectively) were used in a standard PCR protocol to generate DNA for sequencing.

5'-TATAGACTGCTGGGTAGTCCCCAC-3' (SEQ ID NO: 34)

25 5'-TGAAGATTGGGGGTAAATAACAGA-3' (SEQ ID NO: 35)

DNA from the PCR was purified using a Quick Spin Column (Qiagen) according to manufacturer's suggested protocol.

- 50 -

Two clones, designated 741.4 and 741.11, were identified which overlapped clones 684.3 and 705.1; in the overlapping regions, clones 741.1 and 741.11 were 100% homologous to clones 684.3 and 705.1. A composite rat cDNA having homology to the human  $\alpha_d$  gene is set out in SEQ ID NO: 36; the predicted amino acid sequence is set forth in SEQ ID NO: 37.

Cloning of the 5' end of Rat  $\alpha_d$

A 5' cDNA fragment for the rat  $\alpha_d$  gene was obtained using a Clontech rat spleen RACE cloning kit according to manufacturer's suggested protocol. The gene specific oligonucleotides used were designated 741.11#2R and 741.2#1R (SEQ ID NOS: 59 and 58, respectively).

5'-CCAAAGCTGGCTGCATCCTCTC-3' (SEQ ID NO: 59)  
5'-GGCCTTGCAGCTGGACAATG-3' (SEQ ID NO: 58)

Oligo 741.11#2R encompasses base pairs 131-152 in SEQ ID NO: 36, in the reverse orientation and 741.2#1R encompasses bases pairs 696-715 in SEQ ID NO: 36, also in the reverse orientation. A primary PCR was carried out using the 3'-most oligo, 741.2#1R. A second PCR followed using oligo 741.11#2R and DNA generated from the primary reaction. A band of approximately 300 base pairs was detected on a 1% agarose gel.

The secondary PCR product was ligated into plasmid pCRTAI (Invitrogen) according to manufacturer's suggested protocol. White (positive) colonies were picked and added to 100  $\mu$ l LBM containing 1  $\mu$ l of a 50 mg/ml carbenicillin stock solution and 1  $\mu$ l M13 K07 phage culture in individual wells in a round bottom 96 well tissue culture plate. The mixture was incubated at 37°C for 30 minutes to one hour. Following the initial incubation period, 100  $\mu$ l of LBM (containing 1  $\mu$ l of 50 mg/ml carbenicillin and a 1:250 dilution of a 10

- 51 -

mg/ml kanamycin stock solution) were added and the incubation was continued overnight at 37°C.

Using a sterile 96 well metal transfer prong, supernatant from the 96 well plate was transferred to four Amersham Hybond nylon filters. The filters were denatured, neutralized and cross linked by standard protocols. The filters were prehybridized in 20 mls of prehybridization buffer (5X SSPE; 5X Denhardt's; 1% SDS; 50 ugs/ml denatured salmon sperm DNA) at 50°C for several hours while shaking.

Oligo probes 741.11#1 and 741.11#1R (SEQ ID NOS: 56 and 57, respectively), encompassing base pairs 86-105 (SEQ ID NO: 36) in the forward and reverse orientation respectively, were labeled as follows.

5'-CCTGTCATGGGTCTAACCTG-3' (SEQ ID NO: 56)

5'-AGGTTAGACCCATGACAGG-3' (SEQ ID NO: 57)

Approximately 65 ng oligo DNA in 12  $\mu$ l dH<sub>2</sub>O was heated to 65°C for two minutes. Three  $\mu$ l of 10 mCi/ml  $\gamma$ -<sup>32</sup>P-ATP were added to the tube along with 4  $\mu$ l 5x Kinase Buffer (Gibco) and 1  $\mu$ l T4 DNA Kinase (Gibco). The mixture was incubated at 37°C for 30 minutes. Following incubation, 16  $\mu$ l of each labeled oligo probe were added to the prehybridization buffer and filters and hybridization was continued overnight at 42°C. The filters were washed three times in 5X SSPE; 0.1% SDS for 5 minutes per wash at room temperature, and autoradiographed for 6 hours. Positive clones were expanded and DNA purified using the Magic Mini Prep Kit (Promega) according to manufacturer's suggested protocol. Clone 2F7 was selected for sequencing and showed 100% homology to clone 741.11 in the overlapping region. The complete rat  $\alpha_d$  nucleic acid sequence is set out in SEQ ID NO: 54; the amino acid sequence is set out in SEQ ID NO: 55.

- 52 -

### Characteristics of the Rat cDNA and Amino Acid Sequences

Neither nucleic acid nor amino acid sequences have previously been reported for rat  $\alpha$  subunits in  $\beta_2$  integrins. However sequence comparisons to reported human  $\beta_2$  integrin  $\alpha$  subunits suggests that the isolated rat clone and its predicted amino acid sequence are most closely related to  $\alpha_d$  nucleotide and amino acid sequences.

At the nucleic acid level, the isolated rat cDNA clone shows 80% identity in comparison to the human  $\alpha_d$  cDNA; 68% identity in comparison to human CD11b; 70% identity in comparison to human CD11c; and 65% identity in comparison to mouse CD11b. No significant identity is found in comparison to human CD11a and to mouse CD11a.

At the amino acid level, the predicted rat polypeptide encoded by the isolated cDNA shows 70% identity in comparison to human  $\alpha_d$  polypeptide; 28% identity in comparison to human CD11a; 58% identity in comparison to human CD11b; 61% identity in comparison to human CD11c; 28% identity in comparison to mouse CD11a; and 55% identity in comparison to mouse CD11b.

### Example 18

#### Monoclonal Antibodies against Rat $\alpha_d$ I domain/Hu IgG4 Fusion Proteins

In view of the fact that the I domain of human  $\beta_2$  integrins has been demonstrated to participate in ligand binding, it was assumed that the same would be true for rat  $\alpha_d$  protein. Monoclonal antibodies immunospecific for the rat  $\alpha_d$  I domain may therefore be useful in rat models of human disease states wherein  $\alpha_d$  binding is implicated.

Oligos "rat alpha-DI5" (SEQ ID NO: 87) and "rat alpha-DI3" (SEQ ID NO: 88) were generated from the rat  $\alpha_d$  sequence corresponding to base pairs 469-493 and base pairs 1101-1125 (in the reverse orientation), respectively, in SEQ ID NO: 54. The oligos were used in a standard PCR reaction to generate a rat  $\alpha_d$  DNA fragment containing the I domain spanning base pairs 459-1125 in



- 53 -

SEQ ID NO: 54. The PCR product was ligated into vector pCRTAII (Invitrogen) according to manufacturer's suggested protocol. A positive colony was selected and expanded for DNA purification using a Qiagen (Chatswoth, GA) Midi Prep kit according to manufacturer's protocol. The DNA was digested with *Xho*I and *Bgl*III in a standard restriction enzyme digest and a 600 base pair band was gel purified which was subsequently ligated into pDCS1/HuIgG4 expression vector. A positive colony was selected, expanded and DNA purified with a Quiagen Maxi Prep Kit.

COS cells were plated at half confluence on 100mm culture dishes and grown overnight at 37°C in 7% CO<sub>2</sub>. Cells were rinsed once with 5 ml DMEM. To 5 ml DMEM, 50 µl DEAE-Dextran, 2 µl chloroquine and 15 µg rat α<sub>d</sub> I domain/HuIgG4 DNA described above was added. The mixture was added to the COS cells and incubated at 37°C for 3 hours. Media was then removed and 5 ml 10% DMSO in CMF-PBS was added for exactly one minute. The cells were gently rinsed once with DMEM. Ten ml DMEM containing 10% FBS was added to the cells and incubation continued overnight at 37°C in 7% CO<sub>2</sub>. The next day, media was replaced with fresh media and incubation continued for three additional days. The media was harvested and fresh media was added to the plate. After three days, the media was collected again and the plates discarded. The procedure was repeated until 2 liters of culture supernatant were collected.

Supernatant collected as described above was loaded onto a Prosep-A column (Bioprocessing Limited) and protein purified as described below.

The column was initially washed with 15 column volumes of Wash Buffer containing 35 mM Tris and 150 mM NaCl, pH 7.5. Supernatant was loaded at a slow rate of less than approximately 60 column volumes per hour. After loading, the column was washed with 15 column volumes of Wash Buffer, 15 column volumes of 0.55 M diethanolamine, pH 8.5, and 15 column volumes

- 54 -

50 mM citric acid, pH 5.0. Protein was eluted with 50 mM citric acid, pH 3.0. Protein was neutralized with 1.0 M Tris, pH 8.0, and dialyzed in sterile PBS.

The rat  $\alpha_d$  I domain protein was analyzed as described in Example 14. The detected protein migrated in the same manner as observed with human I domain protein.

#### Immunization Protocol

Mice were individually immunized with 50  $\mu$ g purified rat  $\alpha_d$  I domain/HuIgG4 fusion protein previously emulsified in an equal volume of Freund's Complete Adjuvant (FCA) (Sigma). Approximately 200  $\mu$ l of the antigen/adjuvant preparation was injected at 4 sites in the back and flanks of each of the mice. Two weeks later the mice were boosted with an injection of 100  $\mu$ l rat  $\alpha_d$  I domain/HuIgG4 antigen (50  $\mu$ g/mouse) previously emulsified in an equal volume of Freund's Incomplete Adjuvant (FIA). After two additional weeks, the mice were boosted with 50  $\mu$ g antigen in 200  $\mu$ l PBS injected intravenously.

To evaluate serum titers in the immunized mice, retro-orbital bleeds were performed on the animals ten days following the third immunization. The blood was allowed to clot and serum isolated by centrifugation. The serum was used in an immunoprecipitation on biotinylated (BIP) rat splenocytes. Serum from each mouse immunoprecipitated protein bands of expected molecular weight for rat  $\alpha_d$  and rat CD18. One mouse was selected for the fusion and was boosted a fourth time as described above for the third boost.

The hybridoma supernatants were screened by antibody capture, described as follows. Immulon 4 plates (Dynatech, Cambridge, Massachusetts) were coated at 4°C with 50  $\mu$ l/well goat anti-mouse IgA, IgG or IgM (Organon Teknika) diluted 1:5000 in 50 mM carbonate buffer, pH 9.6. Plates were washed 3X with PBS containing 0.05% Tween 20 (PBST) and 50  $\mu$ l culture supernatant was added. After incubation at 37°C for 30 minutes, and washing as described above, 50  $\mu$ l horseradish peroxidase-conjugated goat anti-mouse IgG9(Fc)

- 55 -

(Jackson ImmunoResearch, West Grove, Pennsylvania) diluted 1:3500 in PBST was added. Plates were incubated as described above and washed 4X with PBST. Immediately thereafter, 100 $\mu$ l substrate, containing 1 mg/ml *o*-phenylene diamine (Sigma) and 0.1  $\mu$ l/ml 30% H<sub>2</sub>O<sub>2</sub> in 100 mM citrate, pH4.5, was added. The color reaction was stopped after 5 minutes with the addition of 50  $\mu$ l 15% H<sub>2</sub>SO<sub>4</sub>. Absorbance at 490 nm was read on a Dynatech plate reader.

Supernatant from antibody-containing wells was also analyzed by ELISA with immobilized rat  $\alpha_d$  I domain/HuIgG4 fusion protein. An ELISA with HuIgG4 antibody coated plates served as a control for reactivity against the IgG fusion partner. Positive wells were selected for further screening by BIP on rat splenocyte lysates using techniques described below.

#### Biotinylation of Cell Surface Antigens

Rats were sacrificed by asphyxiation with CO<sub>2</sub> and spleens were removed using standard surgical techniques. Splenocytes were harvested by gently pushing the spleen through a wire mesh with a 3 cc syringe plunger in 20 mls RPMI. Cells were collected into a 50 ml conical tube and washed in the appropriate buffer.

Cells were washed three times in cold D-PBS and resuspended at a density of 10<sup>8</sup> to 10<sup>9</sup> cells in 40 ml PBS. Four mg of NHS-Biotin (Pierce) was added to the cell suspension and the reaction was allowed to continue for exactly 15 minutes at room temperature. The cells were pelleted and washed three times in cold D-PBS.

#### Cell Lysates

Cells were resuspended at a density of 10<sup>8</sup> cells/ml in cold lysis Buffer (1% NP40; 50 mM Tris-HCl, pH 8.0; 150 mM NaCl; 2 mM CaCl<sub>2</sub>; 2 mM MgCl<sub>2</sub>; 1:100 solution of pepstain, leupeptine, and aprotinin, added just before adding to cells; and 0.0001 g PMSF crystals, added just before adding to cells).

- 56 -

Lysates were vortexed for approximately 30 seconds, incubated for 5 minute at room temperature, and further incubated for 15 minutes on ice. Lysates were centrifuged for 10 minutes at 10,000 xg to pellet the insoluble material. Supernatant was collected into a new tube and stored at between 4°C and -20°C.

5     Immunoprecipitation

One ml cell lysate was precleared by incubation with 200  $\mu$ l of a protein A sepharose slurry (Zymed) overnight at 4°C. Precleared lysate was aliquoted into Eppendorf tubes at 50  $\mu$ l/tube for each antibody to be tested. Twenty-five  $\mu$ l of polyclonal serum or 100 to 500  $\mu$ l of monoclonal antibody supernatant were added to the precleared lysates and the resulting mixture incubated for 2 hours at 4°C with rotation. One hundred  $\mu$ l rabbit anti-mouse IgG (Jackson) bound to protein A sepharose beads in a PBS slurry was then added and incubation continued for 30 minutes at room temperature with rotation. Beads were pelleted with gentle centrifugation, and washed three times with cold Wash Buffer (10 mM HEPES; 0.2 M NaCl; 1% Triton X-100). Supernatant was removed by aspiration, and 20  $\mu$ l 2X SDS sample buffer containing 10%  $\beta$ -mercaptoethanol was added. The sample was boiled for 2 minutes in a water bath, and the sample loaded onto a 5% SDS PAGE gel. Following separation, the proteins were transferred to nitrocellulose at constant current overnight. The nitrocellulose filters were blocked with 3% BSA in TBS-T for 1 hour at room temperature and the blocking buffer was removed. A 1:6000 dilution of Streptavidin-HRP conjugate (Jackson) in 0.1% BSA TBS-T was added and incubation continued for 30 minutes at room temperature. Filters were washed three times for 15 minutes each with TBS-T and autoradiographed using Amersham's ECL kit according to manufacturer's suggested protocol.

- 57 -

**Example 19****Isolation of Mouse cDNA Clones**

Isolation of a mouse  $\alpha_d$  homolog was attempted.

5 Cross-species hybridization was performed using two PCR-generated probes: a 1.5 kb fragment corresponding to bases 522 to 2047 from human clone 19A2 (SEQ ID NO: 1), and a 1.0 kb rat fragment which corresponds to bases 1900 to 2900 in human clone 19A2 (SEQ ID NO: 1). The human probe was generated by PCR using primer pairs designated ATM-2 and 9-10.1 set out in SEQ ID NOS: 38 and 39, respectively; the rat probe was generated using  
10 primer pairs 434L and 434R, set out in SEQ ID NOS: 34 and 35, respectively. Samples were incubated at 94°C for 4 minutes and subjected to 30 cycles of the temperature step sequence: 94°C; 50°C 2 minutes; 72°C, 4 minutes.

5'-GTCCAAGCTGTCATGGGCCAG-3' (SEQ ID NO: 38)

5'-GTCCAGCAGACTGAAGAGCACGG-3' (SEQ ID NO: 39)

15 The PCR products were purified using the Qiagen Quick Spin kit according to manufacturer's suggested protocol, and approximately 180 ng DNA was labeled with 200  $\mu$ Ci [ $^{32}$ P]-dCTP using a Boehringer Mannheim Random Primer Labeling kit according to manufacturer's suggested protocol. Unincorporated isotope was removed using a Centri-sep Spin Column (Princeton  
20 Separations, Adelphia, NJ) according to manufacturer's suggested protocol. The probes were denatured with 0.2 N NaOH and neutralized with 0.4 M Tris-HCl, pH 8.0, before use.

25 A mouse thymic oligo dT-primed cDNA library in lambda ZAP II (Stratagene) was plated at approximately 30,000 plaques per 15 cm plate. Plaque lifts on nitrocellulose filters (Schleicher & Schuell, Keene, NH) were incubated at 50°C with agitation for 1 hour in a prehybridization solution (8 ml/lift) containing 30% formamide. Labeled human and rat probes were added to the

- 58 -

prehybridization solution and incubation continued overnight at 50°C. Filters were washed twice in 2X SSC/0.1% at room temperature, once in 2X SSC/0.1% SDS at 37°C, and once in 2X SSC/0.1% SDS at 42°C. Filters were exposed on Kodak X-Omat AR film at -80°C for 27 hours with an intensifying screen.

5 Four plaques giving positive signals on duplicate lifts were restreaked on LB medium with magnesium (LBM)/carbenicillin (100 mg/ml) plates and incubated overnight at 37°C. The phage plaques were lifted with Hybond filters (Amersham), probed as in the initial screen, and exposed on Kodak X-Omat AR film for 24 hours at -80°C with an intensifying screen.

10 Twelve plaques giving positive signals were transferred into low  $Mg^{++}$  phage diluent containing 10 mM Tris-HCl and 1 mM  $MgCl_2$ . Insert size was determined by PCR amplification using T3 and T7 primers (SEQ ID NOS: 13 and 14, respectively) and the following reaction conditions. Samples were incubated at 94°C for 4 minutes and subjected to 30 cycles of the temperature  
15 step sequence: 94°C, for 15 seconds; 50°C, for 30 seconds; and 72°C for 1 minute.

Six samples produced distinct bands that ranged in size from 300 bases to 1 kb. Phagemids were released via co-infection with helper phage and recircularized to generate Bluescript SK<sup>-</sup> (Stratagene). The resulting colonies  
20 were cultured in LBM/carbenicillin (100 mg/ml) overnight. DNA was isolated with a Promega Wizard miniprep kit (Madison, WI) according to manufacturer's suggested protocol. *EcoRI* restriction analysis of purified DNA confirmed the molecular weights which were detected using PCR. Insert DNA was sequenced with M13 and M13 reverse.1 primers set out in SEQ ID NOS: 40 and 41,  
25 respectively.

5'-TGTAACGACGGCCAGT-3'

(SEQ ID NO: 40)

5'-GGAAACAGCTATGACCATG-3'

(SEQ ID NO: 41)

- 59 -

Sequencing was performed as described in Example 4.

Of the six clones, only two, designated 10.3-1 and 10.5-2, provided sequence information and were identical 600 bp fragments. The 600 bp sequence was 68% identical to a corresponding region of human  $\alpha_d$ , 40% identical to human CD11a, 58% identical to human CD11c, and 54% identical to mouse CD11b. This 600 bp fragment was then utilized to isolate a more complete cDNA encoding a putative mouse  $\alpha_d$  homolog .

A mouse splenic cDNA library (oligo dT<sup>-</sup> and random-primed) in lambda Zap II (Stratagene) was plated at  $2.5 \times 10^4$  phage/15 cm LBM plate. Plaques were lifted on Hybond nylon transfer membranes (Amersham), denatured with 0.5 M NaOH/1.5 M NaCl, neutralized with 0.5 M Tris Base/1.5 M NaCl/11.6 HCl, and washed in 2X SSC. The DNA was cross-linked to filters by ultraviolet irradiation.

Approximately 500,000 plaques were screened using probes 10.3-1 and 10.5-2 previously labeled as described *supra*. Probes were added to a prehybridization solution and incubated overnight at 50°C. The filters were washed twice in 2X SSC/0.1% SDS at room temperature, once in 2X SSC/0.1% SDS at 37°C, and once in 2X SSC/0.1% SDS at 42°C. Filters were exposed on Kodak X-Omat AR film for 24 hours at -80°C with an intensifying screen. Fourteen plaques giving positive signals on duplicate lifts were subjected to a secondary screen identical to that for the initial screen except for additional final high stringency washes in 2X SSC/ 0.1% SDS at 50°C, in 0.5X SSC/0.1% SDS at 50°C, and at 55°C in 0.2X SSC/0.1% SDS. The filters were exposed on Kodak X-Omat AR film at -80°C for 13 hours with an intensifying screen.

Eighteen positive plaques were transferred into low Mg<sup>++</sup> phage diluent and insert size determined by PCR amplification as described above. Seven of the samples gave single bands that ranged in size from 600 bp to 4 kb. *EcoRI* restriction analysis of purified DNA confirmed the sizes observed from

- 60 -

PCR and the DNA was sequenced with primers M13 and M13 reverse.1 (SEQ ID NOS: 40 and 41, respectively).

One clone designated B3800 contained a 4 kb insert which corresponded to a region 200 bases downstream of the 5' end of the human  $\alpha_d$  19A2 clone and includes 553 bases of a 3' untranslated region. Clone B3800 showed 77% identity to a corresponding region of human  $\alpha_d$ , 44% identity to a corresponding region of human CD11a, 59% identity to a corresponding region of human CD11c, and 51% identity to a corresponding region of mouse CD11b. The second clone A1160 was a 1.2 kb insert which aligned to the 5' end of the coding region of human  $\alpha_d$  approximately 12 nucleic acids downstream of the initiating methionine. Clone A1160 showed 75% identity to a corresponding region of human  $\alpha_d$ , 46% identity to a corresponding region of human CD11a, 62% identity to a corresponding region of human CD11c, and 66% identity to a corresponding region of mouse CD11b.

Clone A1160, the fragment closer to the 5' end of human clone 19A2, is 1160 bases in length, and shares a region of overlap with clone B3800 starting at base 205 and continuing to base 1134. Clone A1160 has a 110-base insertion (bases 704-814 of clone A1160) not present in the overlapping region of clone B3800. This insertion occurs at a probable exon-intron boundary [Fleming, *et al.*, *J.Immunol.* 150:480-490 (1993)] and was removed before subsequent ligation of clones A1160 and B3800.

#### Rapid Amplification of 5' cDNA End of the Putative Mouse $\alpha_d$ Clone

RACE PCR [Frohman, "RACE: Rapid Amplification of cDNA Ends," in PCR Protocols: A Guide to Methods and Applications, Innis, *et al.* (eds.) pp. 28-38, Academic Press:New York (1990)] was used to obtain missing 5' sequences of the putative mouse  $\alpha_d$  clone, including 5' untranslated sequence and initiating methionine. A mouse splenic RACE-Ready kit (Clontech, Palo Alto, CA) was used according to the manufacturer's suggested protocol. Two



- 61 -

antisense, gene-specific primers, A1160 RACE1-primary and A1160 RACE2-nested (SEQ ID NOS: 42 and 43), were designed to perform primary and nested PCR.

5'-GGACATGTTCACTGCCTCTAGG-3' (SEQ ID NO: 42)

5 5'-GGCGGACAGTCAGACGACTGTCCTG-3' (SEQ ID NO: 43)

The primers, SEQ ID NOS: 42 and 43, correspond to regions starting 302 and 247 bases from the 5' end, respectively. PCR was performed as described, *supra*, using the 5' anchor primer (SEQ ID NO: 44) and mouse spleen cDNA supplied with the kit.

10 5'-CTGGTTCGGCCACCTCTGAAGGTTCCAGAATCGAT (SEQ ID NO: 44)

Electrophoresis of the PCR product revealed a band approximately 280 bases in size, which was subcloned using a TA cloning kit (Invitrogen) according to manufacturer's suggested protocol. Ten resulting colonies were cultured, and the DNA isolated and sequenced. An additional 60 bases of 5' sequence were identified by this method, which correspond to bases 1 to 60 in SEQ ID NO: 45.

#### Characteristics of the Mouse cDNA and Predicted Amino Acid Sequence

A composite sequence of the mouse cDNA encoding a putative homolog of human  $\alpha_d$  is set out in SEQ ID NO: 45. Although homology between the external domains of the human and mouse clones is high, homology between the cytoplasmic domains is only 30%. The observed variation may indicate C-terminal functional differences between the human and mouse proteins. Alternatively, the variation in the cytoplasmic domains may result from splice variation, or may indicate the existence of an additional  $\beta_2$  integrin gene(s).

- 62 -

At the amino acid level, the mouse cDNA predicts a protein (SEQ ID NO: 46) with 28% identity to mouse CD11a, 53% identity to mouse CD11b, 28% identity to human CD11a, 55% identity to human CD11b, 59% identity to human CD11c, and 70% identity to human  $\alpha_d$ . Comparison of the amino acid sequences of the cytoplasmic domains of human  $\alpha_d$  and the putative mouse homolog indicates regions of the same length, but having divergent primary structure. Similar sequence length in these regions suggests species variation rather than splice variant forms. When compared to the predicted rat polypeptide, Example 16, *supra*, mouse and rat cytoplasmic domains show greater than 60% identity.

#### Example 20

##### Isolation of additional mouse $\alpha_d$ cDNA clones for sequence verification

In order to verify the nucleic and amino acids sequences describe in Example 19 for mouse  $\alpha_d$ , additional mouse sequences were isolated for the purposes of confirmation.

Isolation of mouse cDNA by hybridization with two homologous  $\alpha_d$  probes (3' and 5') was performed using both a mouse splenic random primed library and an oligo dT-primed cDNA library in lambda ZAP II (Stratagene). The library was plated at  $5 \times 10^5$  phage per 15 cm LBM plate. Plaques were lifted on Hybond nylon membranes (Amersham), and the membranes were denatured (0.5 M NaOH/1.5 M NaCl), neutralized (0.5 M Tris Base/1.5 M NaCl / 11.6 M HCl) and washed (2X SSC salt solution). DNA was cross-lined to filters by ultraviolet irradiation.

Probes were generated using primers described below in a PCR reaction under the following conditions. Samples were held at 94°C for 4 minutes and then run through 30 cycles of the temperature step sequence (94°C for 15 seconds; 50°C for 30 seconds; 72°C for 1 minute in a Perkin-Elmer 9600 thermocycler).

- 63 -

The 3' probe was approximately 900 bases long and spanned a region from nucleotides 2752 to 3651 (in SEQ ID NO: 1) (5' → 3') and was produced with primers 11.b-1/2FOR11 and 11.b-1/2REV2 as shown in SEQ ID NOS: 69 and 74, respectively. This probe was used in a first set of lifts.

5           The 5' probe was approximately 800 bases long and spanned a region from nucleotides 149 to 946 (in SEQ ID NO: 1) (5' → 3') and was produced with primers 11.b-1/2FOR1 and 11.a-1/1REV1 as shown in SEQ ID NOS: 50 and 85, respectively). This probe was used in a second set of lifts.

10           In a third set of lifts, both probes described above were used together on the same plates.

15           Approximately 500,000 plaques were screened using the two probes from above which were labeled in the same way as described in Example 17. Labeled probes were added to a prehybridization solution, containing 45% formamide, and incubated overnight at 50°C. Filters were washed twice in 2X SSC/0.1% SDS at room temperature (22°C). A final wash was carried out in 2X SSC/0.1% SDS at 50°C. Autoradiography was for 19 hours at -80°C on Kodak X-Omat AR film with an intensifying screen.

20           Thirteen plaques giving positive signals on at least duplicate lifts were subjected to a secondary screen performed as described for the initial screen except that both the 3' and 5' labeled probes were used for hybridization and an additional final wash was incorporated using 2X SSC/0.1% SDS at 65°C. Autoradiography was performed as described above for 2.5 hours.

25           Thirteen plaques (designated MS2P1 through MS2P13) giving positive signals were transferred into low Mg<sup>++</sup> phage diluent. Insert size was determined by PCR amplification (Perkin-Elmer 9600 thermocycler) using T3 and T7 primers which anneal to Bluescript phagemid in ZAP II (sequence previously described) under the same conditions shown above. Band sizes ranged from 500 bases to 4Kb. Phagemids were isolated, prepared, and sequenced with M13 and M13 reverse.1 primers (SEQ ID NOS: 40 and 41, respectively). Five of the

- 64 -

thirteen clones; MS2P-3, MS2P-6, MS2P-9, MS2P-12, and MS2P-13, were sequenced, and together, represented a region from approximately base 200 at the 5' end to about 300 bases past a first stop codon at the 3' end.

Automated sequencing was performed as described in Example 4 by first using M13 and M13 reverse.1 primers (SEQ ID NOS: 40 and 41, respectively) to sequence the ends of each clone and to determine its position relative to construct #17 (SEQ ID NO: 45). Each clone was then completely sequenced using the appropriate primers (listed below) for that particular region.

	11.b-1/2FOR1	5'-GCAGCCAGCTTCGGACAGAC-3'	(SEQ ID NO: 50)
10	11.a-1/1FOR2	5'-CCGCCTGCCACTGGCGTGTGC-3'	(SEQ ID NO: 60)
	11.a-1/1FOR3	5'-CCCAGATGAAGGACTTCGTCAA-3'	(SEQ ID NO: 61)
	11.b-1/2FOR4	5'-GCTGGGATCATTCGCTATGC-3'	(SEQ ID NO: 62)
	11.b-1/2FOR5	5'-CAATGGATGGACCAGTTCTGG-3'	(SEQ ID NO: 63)
	11.b-1/2FOR6	5'-CAGATCGGCTCCTACTTTGG-3'	(SEQ ID NO: 64)
15	11.b-1/2FOR7	5'-CATGGAGCCTCGAGACAGG-3'	(SEQ ID NO: 65)
	11.b-1/2FOR8	5'-CCACTGTCCTCGAAGCTGGAG-3'	(SEQ ID NO: 66)
	11.b-1/2FOR9	5'-CTTCGTCCTGTGCTGGCTGTGGGCTC-3'	(SEQ ID NO: 67)
	11.b-1/2FOR10	5'-CGCCTGGCATGTGAGGCTGAG-3'	(SEQ ID NO: 68)
20	11.b-1/2FOR11	5'-CCGTGATCAGTAGGCAGGAAG-3'	(SEQ ID NO: 69)
	11.b-1/2FOR12	5'-GTCACAGAGGGAACCTCC-3'	(SEQ ID NO: 70)
	11.b-1/2FOR13	5'-GCTCCTGAGTGAGGCTGAAATCA-3'	(SEQ ID NO: 71)
	11.b-1/2FOR14	5'-GAGATGCTGGATCTACCATCTGC-3'	(SEQ ID NO: 72)
	11.b-1/2FOR15	5'-CTGAGCTGGGAGATTTTATGG-3'	(SEQ ID NO: 73)
25	11.b-1/2REV2	5'-GTGGATCAGCACTGAAATCTG-3'	(SEQ ID NO: 74)
	11.b-1/2REV3	5'-CGTTTGAAGAAGCCAAGCTTG-3'	(SEQ ID NO: 75)
	11.b-1/2REV4	5'-CACAGCGGAGGTGCAGGCAG-3'	(SEQ ID NO: 76)
	11.b-1/2REV5	5'-CTCACTGCTTGCGCTGGC-3'	(SEQ ID NO: 77)

- 65 -

11.b-1/2REV6 5'-CGGTAAGATAGCTCTGCTGG-3' (SEQ ID NO: 78)  
 11.b-1/2REV7 5'-GAGCCCACAGCCAGCACAGG-3' (SEQ ID NO: 79)  
 11.b-1/2REV8 5'-GATCCAACGCCAGATCATACC-3' (SEQ ID NO: 80)  
 11.b-1/2REV9 5'-CACGGCCAGGTCCACCAGGC-3' (SEQ ID NO: 81)  
 5 11.b-1/2REV10 5'-CACGTCCCCTAGCACTGTCAG-3' (SEQ ID NO: 82)  
 11.b-1/2REV11 5'-CCATGTCCACAGAACAGAGAG-3' (SEQ ID NO: 51)  
 11.b-1/2REV12 5'-TTGACGAAGTCCTTCATCTGGG-3' (SEQ ID NO: 83)  
 11.b-1/2REV13 5'-GAACTGCAAGCTGGAGCCCAG-3' (SEQ ID NO: 84)  
 11.a-1/1REV1 5'-CTGGATGCTGCGAAGTGCTAC-3' (SEQ ID NO: 85)  
 10 11.a-1/1REV2 5'-GCCTTGGAGCTGGACGATGGC-3' (SEQ ID NO: 86)

Sequences were edited, aligned, and compared to a previously isolated mouse  $\alpha_d$  sequence (construct #17, SEQ ID NO: 45).

Alignment of the new sequences revealed an 18 base deletion in  
 construct #17 beginning at nucleotide 2308; the deletion did not cause a shift in  
 15 the reading frame. Clone MS2P-9, sequenced as described above, also revealed  
 the same 18 base deletion. The deletion has been observed to occur in 50% of  
 mouse clones that include the region but has not been detected in rat or human  $\alpha_d$   
 clones. The eighteen base deletion is characterized by a 12 base palindromic  
 sequence AAGCAGGAGCTCCTGTGT (SEQ ID NO: 91). This inverted repeat  
 20 in the nucleic acid sequence is self-complementary and may form a loop out,  
 causing cleavage during reverse transcription. The mouse  $\alpha_d$  sequence which  
 includes the additional 18 bases is set forth in SEQ ID NO: 52; the deduced  
 amino acid sequence is set forth in SEQ ID NO: 53.

- 66 -

Example 21In situ hybridizations in Mouse

Tissue distribution was then determined for mouse  $\alpha_d$  in order to provide a comparison to that in humans, described in Example 6.

5           A single stranded 200 bp mRNA probe was generated from a DNA template, corresponding to nucleotides 3460 to 3707 in the cytoplasmic tail region of the murine cDNA, by *in vitro* RNA transcription incorporating  $^{35}\text{S}$ -UTP (Amersham).

10           Whole mouse embryos (harvested at days 11-18 after fertilization) and various mouse tissues, including spleen, kidney, liver, intestine, and thymus, were hybridized *in situ* with the radiolabeled single-stranded mRNA probe.

15           Tissues were sectioned at 6  $\mu\text{m}$  thickness, adhered to Vectabond (Vector Laboratories, Inc., Burlingame, CA) coated slides, and stored at  $-70^\circ\text{C}$ . Prior to use, slides were removed from  $-70^\circ\text{C}$  and placed at  $50^\circ\text{C}$  for approximately 5 minutes. Sections were fixed in 4% paraformaldehyde for 20 minutes at  $4^\circ\text{C}$ , dehydrated with an increasing ethanol gradient (70-95-100%) for 1 minute at  $4^\circ\text{C}$  at each concentration, and air dried for 30 minutes at room temperature. Sections were denatured for 2 minutes at  $70^\circ\text{C}$  in 70% formamide/2X SSC, rinsed twice in 2X SSC, dehydrated with the ethanol gradient described *supra* and air dried for 30 minutes. Hybridization was carried out  
20           overnight (12-16 hours) at  $55^\circ\text{C}$  in a solution containing  $^{35}\text{S}$ -labeled riboprobes at  $6 \times 10^5$  cpm/section and diethylpyrocarbonate (DEPC)-treated water to give a final concentration of 50% formamide, 0.3 M NaCl, 20 mM Tris-HCl, pH 7.5, 10% dextran sulfate, 1X Denhardt's solution, 100 mM dithiothreitol (DTT) and  
25           5 mM EDTA. After hybridization, sections were washed for 1 hour at room temperature in 4X SSC/10 mM DTT, 40 minutes at  $60^\circ\text{C}$  in 50% formamide/2X SSC/10 mM DTT, 30 minutes at room temperature in 2X SSC, and 30 minutes at room temperature in 0.1X SSC. The sections were dehydrated, air dried for 2 hours, coated with Kodak NTB2 photographic emulsion, air dried for 2 hours,

- 67 -

developed (after storage at 4°C in complete darkness) and counterstained with hematoxylin/eosin.

Spleen tissue showed a strong signal primarily in the red pulp. This pattern is consistent with that of tissue macrophage distribution in the spleen, but does not exclude other cell types.

### Example 22

#### Generation of Mouse Expression Constructs

In order to construct an expression plasmid including mouse cDNA sequences exhibiting homology to human  $\alpha_d$ , inserts from clones A1160 and B3800 were ligated. Prior to this ligation, however, a 5' leader sequence, including an initiating methionine, was added to clone A1160. A primer designated "5' PCR leader" (SEQ ID NO: 47) was designed to contain: (1) identical nonspecific bases at positions 1-6 allowing for digestion; (2) a *Bam*HI site (underlined in SEQ ID NO: 47) from positions 7-12 to facilitate subcloning into an expression vector; (3) a consensus Kozak sequence from positions 13-18, (4) a signal sequence including a codon for an initiating methionine (bold in SEQ ID NO: 47), and (5) an additional 31 bases of specifically overlapping 5' sequence from clone A1160 to allow primer annealing. A second primer designated "3' end frag" (SEQ ID NO: 48) was used with primer "5' PCR leader" to amplify the insert from clone A1160.

5'-AGTTACGGATCCGGCACCATGAC-  
-CTTCGGCACTGTGATCCTCCTGTGTG-3' (SEQ ID NO: 47)

5'-GCTGGACGATGGCATCCAC-3' (SEQ ID NO: 48)

The resulting PCR product did not digest with *Bam*HI, suggesting that an insufficient number of bases preceded the restriction site, prohibiting

- 68 -

recognition by the enzyme. The length of the "tail" sequence preceding the *Bam*HI site in the 5' primer (SEQ ID NO: 47) was increased and PCR was repeated on the amplification product from the first PCR. A 5' primer, designated mAD.5'.2 (SEQ ID NO: 49), was designed with additional nonspecific bases at positions 1-4 and an additional 20 bases specifically overlapping the previously employed "5' PCR leader" primer sequences.

5'-GTAGAGTTACGGATCCGGCACCAT-3' (SEQ ID NO: 49)

Primers "mAD.5'.2" and "3' end frag" were used together in PCR with the product from the first amplification as template. A resulting secondary PCR product was subcloned into plasmid pCRtmII (Invitrogen) according to manufacturer's suggested protocol and transformed into competent One shot cells (Invitrogen). One clone containing the PCR product was identified by restriction enzyme analysis using *Bam*HI and *Eco*RI and sequenced. After the sequence was verified, the insert was isolated by digestion with *Bam*HI and *Eco*RI and gel purified.

The insert from clone B3800 was isolated by digestion with *Eco*RI and *Not*I, gel purified, and added to a ligation reaction which included the augmented A1160 *Bam*HI/*Eco*RI fragment. Ligation was allowed to proceed for 14 hours at 14°C. Vector pcDNA.3 (Invitrogen), digested with *Bam*HI and *Not*I, was added to the ligation reaction with additional ligase and the reaction was continued for another 12 hours. An aliquot of the reaction mixture was transformed into competent *E. coli* cells, the resulting colonies cultured, and one positive clone identified by PCR analysis with the primers 11.b-1/2FOR1 and 11.b-1/2REV11 (SEQ ID NOS: 50 and 51, respectively). These primers bridge the A1160 and B3800 fragments, therefore detection of an amplification product indicates the two fragments were ligated. The sequence of the positive clone was



- 70 -

thymidine kinase encoding cassettes. Further analysis of this clone with an I domain probe (corresponding to nucleotides 454-1064 in SEQ ID NO: 45) indicated that the clone did not contain I domain encoding sequences.

5 Using the same I domain probe, the  $\lambda$ FIXII genomic library was rescreened. Initially, six positive clones were detected, one of which remained positive upon secondary screening. DNA isolated from this clone reacted strongly in Southern analysis with an I domain probe. No reactivity was detected using the original 750 bp probe, however, indicating that this clone included regions 5' to nucleotides 1985-2773 of SEQ ID NO: 45..

10 Alternatively, the lack of hybridization to the 750 bp probe may have suggested that the clone was another member of the integrin family of proteins. To determine if this explanation was plausible, the 13 kb insert was subcloned into pBluescript SKII<sup>+</sup>. Purified DNA was sequenced using primers corresponding to  $\alpha_d$  I domain nucleic acid sequences 441-461, 591-612, 717-739, 15 and reverse 898-918 in SEQ ID NO: 52. Sequence information was obtained using only the first 4441-4461 primer, and only the 5'-most exon of the I domain was efficiently amplified. The remainder of the I domain was not amplified. The resulting clone therefore comprised exon 6 of the mouse  $\alpha_d$  gene, and intronic sequences to the 3' and 5' end of the exon. Exon 7 was not represented in the 20 clone. After sequencing, a construct is generated containing neomycin resistance and thymidine kinase genes.

25 The neomycin resistance ( $neo^r$ ) gene is inserted into the resulting plasmid in a manner that interrupts the protein coding sequence of the genomic mouse DNA. The resulting plasmid therefore contains a  $neo^r$  gene within the mouse genomic DNA sequences, all of which are positioned within a thymidine kinase encoding region. Plasmid construction in this manner is required to favor homologous recombination over random recombination [Chisaka, *et al.*, *Nature* 355:516-520 (1992)].

- 69 -

verified with the primers set out in SEQ ID NOS: 50 and 51, which amplify from base 100 to 1405 after the initiating methionine.

### Example 23

#### Construction of a Knock-out Mouse

5 In order to more accurately assess the immunological role of the protein encoded by the putative mouse  $\alpha_d$  cDNA, a "knock-out" mouse is designed wherein the genomic DNA sequence encoding the putative  $\alpha_d$  homolog is disrupted by homologous recombination. The significance of the protein encoded by the disrupted gene is thereby assessed by the absence of the encoded  
10 protein. Generation of "knock-out" mice is described in Deng, *et al.*, *Mol.Cell.Biol.* 13:2134-2140 (1993).

Design of such a mouse begins with construction of a plasmid containing sequences to be "knocked out" by homologous recombination events. A 750 base pair fragment of the mouse cDNA (corresponding to nucleotides 1985  
15 to 2733 in SEQ ID NO: 45) was used to identify a mouse genomic sequence encoding the putative mouse  $\alpha_d$  homolog from a  $\lambda$ FIXII genomic library. Primary screening resulted in 14 positive plaques, seven of which were confirmed by secondary screening. Liquid lysates were obtained from two of the plaques giving the strongest signal and the  $\lambda$  DNA was isolated by conventional methods.  
20 Restriction mapping and Southern analysis confirmed the authenticity of one clone, designated 14-1, and the insert DNA was isolated by digestion with *NorI*. This fragment was cloned into Bluescript SKII<sup>+</sup>.

In order to identify a restriction fragment of approximately 9 to 14 kb, a length reported to optimize the probability of homologous recombination events, Southern hybridization was performed with the 750 bp cDNA probe.  
25 Prior to hybridization, a restriction map was constructed for clone 14-1. A 12 kb fragment was identified as a possible candidate and this fragment was subcloned into pBluescript SKII<sup>+</sup> in a position wherein the mouse DNA is flanked by

- 72 -

$10^6$  dpm/ml. Hybridization was carried out at 42°C for 16-18 hours. Filters were washed extensively in 2X SSPE/0.1% SDS at room temperature and exposed to X-ray film to visualize any hybridizing plaques.

Two clones with significant sequence homology to human  $\alpha_d$  were identified. Clone #2 was approximately 800 bp in length and mapped to the 5' end of human  $\alpha_d$ . Clone #2 includes an initiating methionine and complete leader sequence. Clone #7 was approximately 1.5 kb and includes an initiating methionine. The 5' end of clone #7 overlapped that of clone #2, while the 3' sequences terminated at a point beyond the I domain sequences. Internal sequencing of clone #7 is performed using the nested deletions sequencing technique.

The predicted N terminal amino acid sequence for rabbit  $\alpha_d$  as determined from clones #2 and #7 indicated a protein with 73% identity with human  $\alpha_d$ , 65% identity with mouse  $\alpha_d$ , and 58% identity with mouse CD11b, human CD11b, and human CD11c. The nucleic acid sequence for clone #2 is set out in SEQ ID NO: 92; the predicted amino acid sequence is set out in SEQ ID NO: 93

Isolation of a full length rabbit  $\alpha_d$  cDNA is carried out using labeled rabbit fragment, clone # 7, and rescreening the cDNA library from which the fragment was derived.

Isolation of a rabbit  $\alpha_d$  clone allows expression of the protein, either on the surface of transfectants or as a soluble full length or truncated form. This protein is then used as an immunogen for the production of monoclonal antibodies for use in rabbit models of human disease states.

- 71 -

**Example 24****Cloning of Rabbit  $\alpha_d$  - Construction and Screening of the Rabbit cDNA Library**

Identification of human  $\alpha_d$  homologs in rats and mice led to the investigation of the existence of a rabbit homolog which would be useful in rabbit models of human disease states described *infra*.

Poly A<sup>+</sup> RNA was prepared from a whole rabbit spleen using an Invitrogen FastTrack kit (San Diego, CA) according to manufacturer's suggested protocol and reagents supplied with the kit. From 1.65 g tissue, 73  $\mu$ g poly A<sup>+</sup>RNA were isolated. The rabbit spleen RNA was used to construct a ZAP Express cDNA library using a kit from Stratagene (La Jolla, CA). Resulting cDNA was directionally cloned into *Eco*RI and *Xho*I sites in the lambda arms of a pBK-CMV phagemid vector. Gigapack II Gold (Stratagene) was used to package the lambda arms into phage particles. The resulting library titer was estimated to be approximately  $8 \times 10^5$  particles, with an average insert size of 1.2 kb.

The library was amplified once by plating for confluent plaque growth and cell lysate was collected. The amplified library was plated at approximately 30,000 plaque forming units (pfu) per 150 mm plate with *E. coli* and the resulting mixture incubated for 12-16 hrs at 37°C to allow plaque formation. Phage DNA was transferred onto Hybond N<sup>+</sup> nylon membranes (Amersham, Arlington Heights, Illinois). The membranes were hybridized with a mixture of two random primed radiolabeled mouse  $\alpha_d$  PCR DNA probes. The first probe was generated from a PCR product spanning nucleotides 149-946 in SEQ ID NO: 52. The second probe was from a PCR product spanning nucleotides 2752-3651 in SEQ ID NO: 52. Probes were labeled by random priming (Boehringer Mannheim Random Primed DNA Labeling Kit) and the reaction mixture was passed over a Sephadex G-50 column to remove unincorporated nucleotides. The hybridization solution was composed of 5X SSPE, 5X Denhardts, 1% SDS, 40% Formamide and the labeled probes at 1 x

- 74 -

functions which damage normal host tissue through either specific autoimmune responses or as a result of bystander cell damage.

5 Disease states in which there is evidence of macrophages playing a significant role in the disease process include multiple sclerosis, arthritis, graft atherosclerosis, some forms of diabetes and inflammatory bowel disease. Animal models, discussed below, have been shown to reproduce many of the aspects of these human disorders. Inhibitors of  $\alpha_d$  function are tested in these model systems to determine if the potential exists for treating the corresponding human diseases.

10 A. Graft Arteriosclerosis

Cardiac transplantation is now the accepted form of therapeutic intervention for some types of end-state heart disease. As the use of cyclosporin A has increased one year survival rates to 80%, the development of progressive graft arteriosclerosis has emerged as the leading cause of death in cardiac transplants surviving beyond the first year. Recent studies have found that the incidence of significant graft arteriosclerosis 3 years following a cardiac transplant is in the range of 36-44% [Adams, *et al.*, *Transplantation* 53:1115-1119 (1992); Adams, *et al.*, *Transplantation* 56:794-799 (1993)].

20 Graft arteriosclerosis typically consists of diffuse, occlusive, intimal lesions which affect the entire coronary vessel wall, and are often accompanied by lipid deposition. While the pathogenesis of graft arteriosclerosis remains unknown, it is presumably linked to histocompatibility differences between donor and recipient, and is immunologic in nature. Histologically, the areas of intimal thickening are composed primarily of macrophages, although T cells are occasionally seen. It is therefore possible that macrophages expressing  $\alpha_d$  may play a significant role in the induction and/or development of graft arteriosclerosis. In such a case, monoclonal antibodies or small molecule inhibitors (for example, soluble ICAM-R) of  $\alpha_d$  function could be given

25

- 73 -

**Example 25****Animal Models For Determining  $\alpha_d$  Therapeutic Utility**

Immunohistologic data in dog and *in situ* hybridization in rats and mice has determined that in spleen  $\alpha_d$  is expressed primarily by macrophages present in red pulp and in lymph nodes,  $\alpha_d$  is found in medullary cords and sinuses. The expression pattern is remarkably similar to what has been reported for two murine antigens defined by the monoclonal antibodies F4/80 and SK39. While biochemical characterization of these murine antigens has demonstrated that they are distinct from  $\alpha_d$ , it is highly probably that  $\alpha_d$  defines the same macrophage subset as the murine F4/80 and SK39 antigens.

In mouse, SK39-positive macrophages have been identified in splenic red pulp where they may participate in the clearance of foreign materials from circulation, and in medulla of lymph nodes [Jutila, *et al.*, *J. Leukocyte Biol.* 54:30-39 (1993)]. SK39-positive macrophages have also been reported at sites of both acute and chronic inflammation. Furthermore, monocytes recruited to thioglycolate-inflamed peritoneal cavities also express the SK39 antigen. Collectively, these findings suggest that, if SK39<sup>+</sup> cells are also  $\alpha_d$ <sup>+</sup>, then these cells are responsible for the clearance of foreign materials in the spleen and participate in inflammation where macrophages play a significant role.

While the function of  $\alpha_d$  remains unclear, other more well characterized  $\beta_2$  integrins have been shown to participate in a wide variety of adhesion events that facilitate cell migration, enhance phagocytosis, and promote cell-cell interactions, events which all lead to upregulation of inflammatory processes. Therefore, it is highly plausible that interfering with the normal  $\alpha_d$  function may also interfere with inflammation where macrophages play a significant role. Such an anti-inflammatory effect could result from: i) blocking macrophage recruitment to sites of inflammation, ii) preventing macrophage activation at the site of inflammation or iii) interfering with macrophage effector

- 76 -

luminal surface of the ascending aorta [Rosenfeld, *et al.*, *Arteriosclerosis* 7:9-23 (1987); Rosenfeld, *et al.*, *Arteriosclerosis* 7:24-34 (1987)]. The atherosclerotic lesions seen in these rabbits are similar to those in humans. Lesions contain large numbers of T cells, most of which express CD45RO, a marker associated with memory T cells. Approximately half of the infiltrating T cells also express MHC class II antigen and some express the IL-2 receptor suggesting that many of the cells are in an activated state.

One feature of the atherosclerotic lesions found in cholesterol fed rabbits, but apparently absent in rodent models, is the accumulation of foam cell-rich lesions. Foam cell macrophages are believed to result from the uptake of oxidized low-density lipoprotein (LDL) by specific receptors. Oxidized LDL particles have been found to be toxic for some cell types including endothelial cells and smooth muscle cells. The uptake of potentially toxic, oxidized LDL particles by macrophages serves as an irritant and drives macrophage activation, contributing to the inflammation associated with atherosclerotic lesions.

Once monoclonal antibodies have been generated to rabbit  $\alpha_d$ , cholesterol fed rabbits are treated. Treatments include prophylactic administration of  $\alpha_d$  monoclonal antibodies or small molecule inhibitors, to demonstrate that  $\alpha_d^+$  macrophages are involved in the disease process. Additional studies would demonstrate that monoclonal antibodies to  $\alpha_d$  or small molecule inhibitors are capable of reversing vessel damage detected in rabbits fed an atherogenic diet.

### C. Insulin-dependent Diabetes

BB rats spontaneously develop insulin-dependent diabetes at 70-150 days of age. Using immunohistochemistry, MHC class II<sup>+</sup>, ED1<sup>+</sup> macrophages can be detected infiltrating the islets early in the disease. Many of the macrophages appear to be engaged in phagocytosis of cell debris or normal cells. As the disease progresses, larger numbers of macrophages are found infiltrating

- 75 -

prophylactically to individuals who received heart transplants and are at risk of developing graft arteriosclerosis.

Although atherosclerosis in heart transplants presents the greatest threat to life, graft arteriosclerosis is also seen in other solid organ transplants, including kidneys and livers. Therapeutic use of  $\alpha_d$  blocking agents could prevent graft arteriosclerosis in other organ transplants and reduce complications resulting from graft failure.

One model for graft arteriosclerosis in the rat involves heterotopic cardiac allografts transplanted across minor histocompatibility barriers. When Lewis cardiac allografts are transplanted into MHC class I and II compatible F-344 recipients, 80% of the allografts survive at least 3 weeks, while 25% of the grafts survive indefinitely. During this low-grade graft rejection, arteriosclerosis lesions form in the donor heart. Arterial lesions in 120 day old allografts typically have diffuse fibrotic intimal thickening indistinguishable in appearance from graft arteriosclerosis lesions found in rejecting human cardiac allografts.

Rats are transplanted with hearts mismatched at minor histocompatibility antigens, for example Lewis into F-344. Monoclonal antibodies specific for rat  $\alpha_d$  or small molecule inhibitors of  $\alpha_d$  are given periodically to transplant recipients. Treatment is expected to reduce the incidence of graft arteriosclerosis in non-rejecting donor hearts. Treatment of rats with  $\alpha_d$  monoclonal antibodies or small molecule inhibitors may not be limited to prophylactic treatments. Blocking  $\alpha_d$  function is also expected to reduce macrophage mediated inflammation and allow reversal of arterial damage in the graft.

25

#### B. Atherosclerosis in Rabbits Fed Cholesterol

Rabbits fed an atherogenic diet containing a cholesterol supplement for approximately 12-16 weeks develop intimal lesions that cover most of the



- 77 -

the islets, although significant numbers of T cells, and later B cells, also appear to be recruited to the site [Hanenberg, *et al.*, *Diabetologia* 32:126-134 (1989)].

Development of diabetes in BB rats appears to depend on both early macrophage infiltration and subsequent T cells recruitment. Treatment of BB rats with silica particles, which are toxic to macrophages, has been effective in blocking the early macrophage infiltration of the islets. In the absence of early macrophage infiltration, subsequent tissue damage by an autoaggressive lymphocyte population fails to occur. Administration of monoclonal antibody OX-19 (specific for rat CD5) or monoclonal antibody OX-8 (specific for rat CD8), which block the T cell-associated phase of the disease, is also effective in suppressing the development of diabetes.

The central role of macrophages in the pathology of this model makes it attractive for testing inhibitors of  $\alpha_d$  function. Rats genetically predisposed to the development of insulin-dependent diabetes are treated with monoclonal antibodies to  $\alpha_d$  or small molecule inhibitors and evaluated for the development of the disease. Preventing or delaying clinical onset is evidence that  $\alpha_d$  plays a pivotal role in macrophage damage to the islet cells.

#### D. Inflammatory Bowel Disease (Crohn's Disease, Ulcerative Colitis)

Animal models used in the study of inflammatory bowel disease (IBD) are generally elicited by intrarectal administration of noxious irritants (e.g. acetic acid or trinitrobenzene sulfonic acid/ethanol). Colonic inflammation induced by these agents is the result of chemical or metabolic injury and lacks the chronic and spontaneously relapsing inflammation associated with human IBD. However, a recently described model using subserosal injections of purified peptidoglycan-polysaccharide (PG-PS) polymers from either group A or group D streptococci appears to be a more physiologically relevant model for human IBD [Yamada, *et al.*, *Gastroenterology* 104:759-771 (1993)].

- 78 -

In this model PG-PS is injected into the subserosal layer of the distal colon. The resulting inflammatory response is biphasic with an initial acute episode three days after injection, which is followed by a spontaneous chronic phase three to four weeks later. The late phase response is granulomatous in nature, and results in colonic thickening, adhesions, colonic nodules and mucosal lesions. In addition to mucosal injury, PG-PS colitis frequently leads to arthritis anemia and granulomatous hepatitis. The extraintestinal manifestations of the disease make the model attractive for studying Crohn's colitis in that a significant number of patients with active Crohn's disease suffer from arthritic joint disease and hepatobiliary inflammation.

Granulomatous lesions are the result of chronic inflammation which leads to the recruitment and subsequent activation of cells of the monocyte/macrophage lineage. Presence of granulomatous lesions in Crohn's disease and the above animal model make this an attractive clinical target for  $\alpha_d$  monoclonal antibodies or other inhibitors of  $\alpha_d$  function. Inhibitors of  $\alpha_d$  function are expected to block the formation of lesions associated with IBD or even reverse tissue damage seen in the disease.

#### E. Arthritis

Arthritis appears to be a multi-factorial disease process involving a variety of inflammatory cell types including neutrophils, T lymphocytes and phagocytic macrophages. Although a variety of arthritis models exist, preparations of streptococcal cell wall proteoglycan produce a disorder most similar to the human disease.

In rats, streptococcal cell wall induces inflammation of peripheral joints characterized by repeated episodes of disease progression followed by remission and eventually resulting in joint destruction over a period of several months [Cromartie, *et al.*, *J.Exp.Med.* 146:1585-1602 (1977); Schwab *et al.*, *Infection and Immunity* 59:4436-4442 (1991)]. During the chronic phase of the

- 79 -

disease, mononuclear phagocytes or macrophages are believed to play a major role in destruction of the synovium. Furthermore, agents which suppress the recruitment of macrophages into the synovium effectively reduce the inflammation and pathology characteristic of arthritis.

5                   A central role for the macrophage in synovium destruction that leads to arthritis predicts that monoclonal antibodies to  $\alpha_d$  or inhibitors of  $\alpha_d$  function may have therapeutic potential in the treatment of this disease. As in other models previously described,  $\alpha_d$  monoclonal antibodies or small molecule inhibitors administered prophylactically are expected to block or moderate joint  
10 inflammation and prevent destruction of the synovium. Agents that interfere with  $\alpha_d$  function may also moderate ongoing inflammation by preventing the recruitment of additional macrophages to the joint or blocking macrophage activation. The net result would be to reverse ongoing destruction of the joint and facilitate tissue repair.

#### 15                   F. Multiple Sclerosis

                  Although pathogenesis of multiple sclerosis (MS) remains unclear, it is generally accepted that the disease is mediated by CD4<sup>+</sup> T cells which recognize autoantigens in the central nervous system and initiate an inflammatory cascade. The resulting immune response results in the recruitment of additional  
20 inflammatory cells, including activated macrophages which contribute to the disease. Experimental autoimmune encephalomyelitis (EAE) is an animal model which reproduces some aspects of MS. Recently, monoclonal antibodies reactive with CD11b/CD18 [Huitinga, *et al.*, *Eur.J.Immunol.* 23:709-715 (1993)] present on inflammatory macrophages have been shown to block both clinical and  
25 histologic disease. The results suggest that monoclonal antibodies or small molecule inhibitors to  $\alpha_d$  are likely to be effective in blocking the inflammatory response in EAE. Such agents also have important therapeutic applications in the treatment of MS.

- 80 -

Numerous modifications and variations in the invention as set forth in the above illustrative examples are expected to occur to those skilled in the art. Consequently only such limitations as appear in the appended claims should be placed on the invention.

**THIS PAGE WAS NOT FURNISHED UPON FILING  
THE INTERNATIONAL APPLICATION**

- 82 -

## (ix) FEATURE:

(A) NAME/KEY: CDS

(B) LOCATION: 3..3485

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

TG ACC TTC GGC ACT GTG CTT CTT CTG AGT GTC CTG GCT TCT TAT CAT	47
Thr Phe Gly Thr Val Leu Leu Leu Ser Val Leu Ala Ser Tyr His	
1 5 10 15	
GGA TTC AAC CTG GAT GTG GAG GAG CCT ACC ATC TTC CAG GAG GAT GCA	95
Gly Phe Asn Leu Asp Val Glu Glu Pro Thr Ile Phe Gln Glu Asp Ala	
20 25 30	
GGC GGC TTT GGG CAG AGC GTG GTG CAG TTC GGT GGA TCT CGA CTC GTG	143
Gly Gly Phe Gly Gln Ser Val Val Gln Phe Gly Gly Ser Arg Leu Val	
35 40 45	
GTG GGA GCA CCC CTG GAG GTG GTG GCG GCC AAC CAG ACG GGA CGG CTG	191
Val Gly Ala Pro Leu Glu Val Val Ala Ala Asn Gln Thr Gly Arg Leu	
50 55 60	
TAT GAC TGC GCA GCT GCC ACC GGC ATG TGC CAG CCC ATC CCG CTG CAC	239
Tyr Asp Cys Ala Ala Ala Thr Gly Met Cys Gln Pro Ile Pro Leu His	
65 70 75	
ATC CGC CCT GAG GCC GTG AAC ATG TCC TTG GGC CTG ACC CTG GCA GCC	287
Ile Arg Pro Glu Ala Val Asn Met Ser Leu Gly Leu Thr Leu Ala Ala	
80 85 90 95	
TCC ACC AAC GGC TCC CGG CTC CTG GCC TGT GGC CCG ACC CTG CAC AGA	335
Ser Thr Asn Gly Ser Arg Leu Leu Ala Cys Gly Pro Thr Leu His Arg	
100 105 110	
GTC TGT GGG GAG AAC TCA TAC TCA AAG GGT TCC TGC CTC CTG CTG GGC	383
Val Cys Gly Glu Asn Ser Tyr Ser Lys Gly Ser Cys Leu Leu Leu Gly	
115 120 125	
TCG CGC TGG GAG ATC ATC CAG ACA GTC CCC GAC GCC ACG CCA GAG TGT	431
Ser Arg Trp Glu Ile Ile Gln Thr Val Pro Asp Ala Thr Pro Glu Cys	
130 135 140	
CCA CAT CAA GAG ATG GAC ATC GTC TTC CTG ATT GAC GGC TCT GGA AGC	479
Pro His Gln Glu Met Asp Ile Val Phe Leu Ile Asp Gly Ser Gly Ser	
145 150 155	
ATT GAC CAA AAT GAC TTT AAC CAG ATG AAG GGC TTT GTC CAA GCT GTC	527
Ile Asp Gln Asn Asp Phe Asn Gln Met Lys Gly Phe Val Gln Ala Val	
160 165 170 175	
ATG GGC CAG TTT GAG GGC ACT GAC ACC CTG TTT GCA CTG ATG CAG TAC	575
Met Gly Gln Phe Glu Gly Thr Asp Thr Leu Phe Ala Leu Met Gln Tyr	
180 185 190	
TCA AAC CTC CTG AAG ATC CAC TTC ACC TTC ACC CAA TTC CGG ACC AGC	623
Ser Asn Leu Leu Lys Ile His Phe Thr Phe Thr Gln Phe Arg Thr Ser	
195 200 205	
CCG AGC CAG CAG AGC CTG GTG GAT CCC ATC GTC CAA CTG AAA GGC CTG	671
Pro Ser Gln Gln Ser Leu Val Asp Pro Ile Val Gln Leu Lys Gly Leu	
210 215 220	

- 83 -

ACG TTC ACG GCC ACG GGC ATC CTG ACA GTG GTG ACA CAG CTA TTT CAT	719
Thr Phe Thr Ala Thr Gly Ile Leu Thr Val Val Thr Gln Leu Phe His	
225 230 235	
CAT AAG AAT GGG GCC CGA AAA AGT GCC AAG AAG ATC CTC ATT GTC ATC	767
His Lys Asn Gly Ala Arg Lys Ser Ala Lys Lys Ile Leu Ile Val Ile	
240 245 250 255	
ACA GAT GGG CAG AAG TAC AAA GAC CCC CTG GAA TAC AGT GAT GTC ATC	815
Thr Asp Gly Gln Lys Tyr Lys Asp Pro Leu Glu Tyr Ser Asp Val Ile	
260 265 270	
CCC CAG GCA GAG AAG GCT GGC ATC ATC CGC TAC GCT ATC GGG GTG GGA	863
Pro Gln Ala Glu Lys Ala Gly Ile Ile Arg Tyr Ala Ile Gly Val Gly	
275 280 285	
CAC GCT TTC CAG GGA CCC ACT GCC AGG CAG GAG CTG AAT ACC ATC AGC	911
His Ala Phe Gln Gly Pro Thr Ala Arg Gln Glu Leu Asn Thr Ile Ser	
290 295 300	
TCA GCG CCT CCG CAG GAC CAC GTG TTC AAG GTG GAC AAC TTT GCA GCC	959
Ser Ala Pro Pro Gln Asp His Val Phe Lys Val Asp Asn Phe Ala Ala	
305 310 315	
CTT GGC AGC ATC CAG AAG CAG CTG CAG GAG AAG ATC TAT GCA GTT GAG	1007
Leu Gly Ser Ile Gln Lys Gln Leu Gln Glu Lys Ile Tyr Ala Val Glu	
320 325 330 335	
GGA ACC CAG TCC AGG GCA AGC AGC TCC TTC CAG CAC GAG ATG TCC CAA	1055
Gly Thr Gln Ser Arg Ala Ser Ser Ser Phe Gln His Glu Met Ser Gln	
340 345 350	
GAA GGC TTC AGC ACA GCC CTC ACA ATG GAT GGC CTC TTC CTG GGG GCT	1103
Glu Gly Phe Ser Thr Ala Leu Thr Met Asp Gly Leu Phe Leu Gly Ala	
355 360 365	
GTG GGG AGC TTT AGC TGG TCT GGA GGT GCC TTC CTG TAT CCC CCA AAT	1151
Val Gly Ser Phe Ser Trp Ser Gly Gly Ala Phe Leu Tyr Pro Pro Asn	
370 375 380	
ATG AGC CCC ACC TTC ATC AAC ATG TCT CAG GAG AAT GTG GAC ATG AGG	1199
Met Ser Pro Thr Phe Ile Asn Met Ser Gln Glu Asn Val Asp Met Arg	
385 390 395	
GAC TCT TAC CTG GGT TAC TCC ACC GAG CTA GCC CTG TGG AAG GGG GTA	1247
Asp Ser Tyr Leu Gly Tyr Ser Thr Glu Leu Ala Leu Trp Lys Gly Val	
400 405 410 415	
CAG AAC CTG GTC CTG GGG GCC CCC CGC TAC CAG CAT ACC GGG AAG GCT	1295
Gln Asn Leu Val Leu Gly Ala Pro Arg Tyr Gln His Thr Gly Lys Ala	
420 425 430	
GTC ATC TTC ACC CAG GTG TCC AGG CAA TGG AGG AAG AAG GCC GAA GTC	1343
Val Ile Phe Thr Gln Val Ser Arg Gln Trp Arg Lys Lys Ala Glu Val	
435 440 445	
ACA GGG ACG CAG ATC GGC TCC TAC TTC GGG GCC TCC CTC TGC TCC GTG	1391
Thr Gly Thr Gln Ile Gly Ser Tyr Phe Gly Ala Ser Leu Cys Ser Val	
450 455 460	

- 84 -

GAT Asp 465	GTG Val	GAC Asp	AGC Ser	GAT Asp	GGC Gly	AGC Ser 470	ACC Thr	GAC Asp	CTG Leu	ATC Ile	CTC Leu 475	ATT Ile	GGG Gly	GCC Ala	CCC Pro	1439
CAT His 480	TAC Tyr	TAT Tyr	GAG Glu	CAG Gln	ACC Thr 485	CGA Arg	GGG Gly	GGC Gly	CAG Gln	GTG Val 490	TCC Ser	GTG Val	TGT Cys	CCC Pro	TTG Leu 495	1487
CCT Pro	AGG Arg	GGG Gly	CAG Gln	AGG Arg 500	GTG Val	CAG Gln	TGG Trp	CAG Gln	TGT Cys 505	GAC Asp	GCT Ala	GTT Val	CTC Leu	CGT Arg 510	GGT Gly	1535
GAG Glu	CAG Gln	GGC Gly	CAC His 515	CCC Pro	TGG Trp	GGC Gly	CGC Arg	TTT Phe 520	GGG Gly	GCA Ala	GCC Ala	CTG Leu	ACA Thr 525	GTG Val	TTG Leu	1583
GGG Gly	GAT Asp	GTG Val 530	AAT Asn	GAG Glu	GAC Asp	AAG Lys 535	CTG Leu	ATA Ile	GAC Asp	GTG Val	GCC Ala 540	ATT Ile	GGG Gly	GCC Ala	CCG Pro	1631
GGA Gly	GAG Glu 545	CAG Gln	GAG Glu	AAC Asn	CGG Arg	GGT Gly 550	GCT Ala	GTC Val	TAC Tyr	CTG Leu	TTT Phe 555	CAC His	GGA Gly	GCC Ala	TCA Ser	1679
GAA Glu 560	TCC Ser	GGC Gly	ATC Ile	AGC Ser	CCC Pro 565	TCC Ser	CAC His	AGC Ser	CAG Gln	CGG Arg 570	ATT Ile	GCC Ala	AGC Ser	TCC Ser	CAG Gln 575	1727
CTC Leu	TCC Ser	CCC Pro	AGG Arg	CTG Leu 580	CAG Gln	TAT Tyr	TTT Phe	GGG Gly	CAG Gln 585	GCG Ala	CTG Leu	AGT Ser	GGG Gly	GGT Gly 590	CAG Gln	1775
GAC Asp	CTC Leu	ACC Thr	CAG Gln 595	GAT Asp	GGA Gly	CTG Leu	ATG Met	GAC Asp 600	CTG Leu	GCC Ala	GTG Val	GGG Gly	GCC Ala 605	CGG Arg	GGC Gly	1823
CAG Gln	GTG Val 610	CTC Leu	CTG Leu	CTC Leu	AGG Arg	AGT Ser	CTG Leu 615	CCG Pro	GTG Val	CTG Leu	AAA Lys 620	GTG Val	GGG Gly	GTG Val	GCC Ala	1871
ATG Met	AGA Arg 625	TTC Phe	AGC Ser	CCT Pro	GTG Val	GAG Glu 630	GTG Val	GCC Ala	AAG Lys	GCT Ala	GTG Val 635	TAC Tyr	CGG Arg	TGC Cys	TGG Trp	1919
GAA Glu 640	GAG Glu	AAG Lys	CCC Pro	AGT Ser	GCC Ala 645	CTG Leu	GAA Glu	GCT Ala	GGG Gly	GAC Asp 650	GCC Ala	ACC Thr	GTC Val	TGT Cys	CTC Leu 655	1967
ACC Thr	ATC Ile	CAG Gln	AAA Lys	AGC Ser 660	TCA Ser	CTG Leu	GAC Asp	CAG Gln	CTA Leu 665	GGT Gly	GAC Asp	ATC Ile	CAA Gln	AGC Ser 670	TCT Ser	2015
GTC Val	AGG Arg	TTT Phe	GAT Asp 675	CTG Leu	GCA Ala	CTG Leu	GAC Asp	CCA Pro 680	GGT Gly	CGT Arg	CTG Leu	ACT Thr	TCT Ser 685	CGT Arg	GCC Ala	2063
ATT Ile	TTC Phe	AAT Asn 690	GAA Glu	ACC Thr	AAG Lys	AAC Asn	CCC Pro 695	ACT Thr	TTG Leu	ACT Thr	CGA Arg 700	AGA Arg	AAA Lys	ACC Thr	CTG Leu	2111



- 85 -

GGA CTG GGG ATT CAC TGT GAA ACC CTG AAG CTG CTT TTG CCA GAT TGT Gly Leu Gly Ile His Cys Glu Thr Leu Lys Leu Leu Pro Asp Cys 705 710 715	2159
GTG GAG GAT GTG GTG AGC CCC ATC ATT CTG CAC CTC AAC TTC TCA CTG Val Glu Asp Val Val Ser Pro Ile Ile Leu His Leu Asn Phe Ser Leu 720 725 730 735	2207
GTG AGA GAG CCC ATC CCC TCC CCC CAG AAC CTG CGT CCT GTG CTG GCC Val Arg Glu Pro Ile Pro Ser Pro Gln Asn Leu Arg Pro Val Leu Ala 740 745 750	2255
GTG GGC TCA CAA GAC CTC TTC ACT GCT TCT CTC CCC TTC GAG AAG AAC Val Gly Ser Gln Asp Leu Phe Thr Ala Ser Leu Pro Phe Glu Lys Asn 755 760 765	2303
TGT GGG CAA GAT GGC CTC TGT GAA GGG GAC CTG GGT GTC ACC CTC AGC Cys Gly Gln Asp Gly Leu Cys Glu Gly Asp Leu Gly Val Thr Leu Ser 770 775 780	2351
TTC TCA GGC CTG CAG ACC CTG ACC GTG GGG AGC TCC CTG GAG CTC AAC Phe Ser Gly Leu Gln Thr Leu Thr Val Gly Ser Ser Leu Glu Leu Asn 785 790 795	2399
GTG ATT GTG ACT GTG TGG AAC GCA GGT GAG GAT TCC TAC GGA ACC GTG Val Ile Val Thr Val Trp Asn Ala Gly Glu Asp Ser Tyr Gly Thr Val 800 805 810 815	2447
GTC AGC CTC TAC TAT CCA GCA GGG CTG TCG CAC CGA CGG GTG TCA GGA Val Ser Leu Tyr Tyr Pro Ala Gly Leu Ser His Arg Arg Val Ser Gly 820 825 830	2495
GCC CAG AAG CAG CCC CAT CAG AGT GCC CTG CGC CTG GCA TGT GAG ACA Ala Gln Lys Gln Pro His Gln Ser Ala Leu Arg Leu Ala Cys Glu Thr 835 840 845	2543
GTG CCC ACT GAG GAT GAG GGC CTA AGA AGC AGC CGC TGC AGT GTC AAC Val Pro Thr Glu Asp Glu Gly Leu Arg Ser Ser Arg Cys Ser Val Asn 850 855 860	2591
CAC CCC ATC TTC CAT GAG GGC TCT AAC GGC ACC TTC ATA GTC ACA TTC His Pro Ile Phe His Glu Gly Ser Asn Gly Thr Phe Ile Val Thr Phe 865 870 875	2639
GAT GTC TCC TAC AAG GCC ACC CTG GGA GAC AGG ATG CTT ATG AGG GCC Asp Val Ser Tyr Lys Ala Thr Leu Gly Asp Arg Met Leu Met Arg Ala 880 885 890 895	2687
AGT GCA AGC AGT GAG AAC AAT AAG GCT TCA AGC AGC AAG GCC ACC TTC Ser Ala Ser Ser Glu Asn Asn Lys Ala Ser Ser Ser Lys Ala Thr Phe 900 905 910	2735
CAG CTG GAG CTC CCG GTG AAG TAT GCA GTC TAC ACC ATG ATC AGC AGG Gln Leu Glu Leu Pro Val Lys Tyr Ala Val Tyr Thr Met Ile Ser Arg 915 920 925	2783
CAG GAA GAA TCC ACC AAG TAC TTC AAC TTT GCA ACC TCC GAT GAG AAG Gln Glu Glu Ser Thr Lys Tyr Phe Asn Phe Ala Thr Ser Asp Glu Lys 930 935 940	2831

- 86 -

AAA ATG AAA GAG GCT GAG CAT CGA TAC CGT GTG AAT AAC CTC AGC CAG Lys Met Lys Glu Ala Glu His Arg Tyr Arg Val Asn Asn Leu Ser Gln 945 950 955	2879
CGA GAT CTG GCC ATC AGC ATT AAC TTC TGG GTT CCT GTC CTG CTG AAC Arg Asp Leu Ala Ile Ser Ile Asn Phe Trp Val Pro Val Leu Leu Asn 960 965 970 975	2927
GGG GTG GCT GTG TGG GAT GTG GTC ATG GAG GCC CCA TCT CAG AGT CTC Gly Val Ala Val Trp Asp Val Val Met Glu Ala Pro Ser Gln Ser Leu 980 985 990	2975
CCC TGT GTT TCA GAG AGA AAA CCT CCC CAG CAT TCT GAC TTC CTG ACC Pro Cys Val Ser Glu Arg Lys Pro Pro Gln His Ser Asp Phe Leu Thr 995 1000 1005	3023
CAG ATT TCA AGA AGT CCC ATG CTG GAC TGC TCC ATT GCT GAC TGC CTG Gln Ile Ser Arg Ser Pro Met Leu Asp Cys Ser Ile Ala Asp Cys Leu 1010 1015 1020	3071
CAG TTC CGC TGT GAC GTC CCC TCC TTC AGC GTC CAG GAG GAG CTG GAT Gln Phe Arg Cys Asp Val Pro Ser Phe Ser Val Gln Glu Glu Leu Asp 1025 1030 1035	3119
TTC ACC CTG AAG GGC AAT CTC AGT TTC GGC TGG GTC CGC GAG ACA TTG Phe Thr Leu Lys Gly Asn Leu Ser Phe Gly Trp Val Arg Glu Thr Leu 1040 1045 1050 1055	3167
CAG AAG AAG GTG TTG GTC GTG AGT GTG GCT GAA ATT ACG TTC GAC ACA Gln Lys Lys Val Leu Val Val Ser Val Ala Glu Ile Thr Phe Asp Thr 1060 1065 1070	3215
TCC GTG TAC TCC CAG CTT CCA GGA CAG GAG GCA TTT ATG AGA GCT CAG Ser Val Tyr Ser Gln Leu Pro Gly Gln Glu Ala Phe Met Arg Ala Gln 1075 1080 1085	3263
ATG GAG ATG GTG CTA GAA GAA GAC GAG GTC TAC AAT GCC ATT CCC ATC Met Glu Met Val Leu Glu Glu Asp Glu Val Tyr Asn Ala Ile Pro Ile 1090 1095 1100	3311
ATC ATG GGC AGC TCT GTG GGG GCT CTG CTA CTG CTG GCG CTC ATC ACA Ile Met Gly Ser Ser Val Gly Ala Leu Leu Leu Leu Ala Leu Ile Thr 1105 1110 1115	3359
GCC ACA CTG TAC AAG CTT GGC TTC TTC AAA CGC CAC TAC AAG GAA ATG Ala Thr Leu Tyr Lys Leu Gly Phe Phe Lys Arg His Tyr Lys Glu Met 1120 1125 1130 1135	3407
CTG GAG GAC AAG CCT GAA GAC ACT GCC ACA TTC AGT GGG GAC GAT TTC Leu Glu Asp Lys Pro Glu Asp Thr Ala Thr Phe Ser Gly Asp Asp Phe 1140 1145 1150	3455
AGC TGT GTG GCC CCA AAT GTG CCT TTG TCC TAATAATCCA CTTTCCTGTT Ser Cys Val Ala Pro Asn Val Pro Leu Ser 1155 1160	3505
TATCTCTACC ACTGTGGGCT GGACTTGCTT GCAACCATAA ATCAACTTAC ATGGAAACAA	3566
CTTCTGCATA GATCTGCACT GGCCTAAGCA ACCTACCAGG TGCTAAGCAC CTTCTCGGAG	3625
AGATAGAGAT TGTAATGTTT TTACATATCT GTCCATCTTT TTCAGCAATG ACCCACTTTT	3685

- 87 -

TACAGAAGCA GGCATGGTGC CAGCATAAAT TTTCATATGC T

3726

## (2) INFORMATION FOR SEQ ID NO:2:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1161 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

```

Thr Phe Gly Thr Val Leu Leu Leu Ser Val Leu Ala Ser Tyr His Gly
 1           5           10           15
Phe Asn Leu Asp Val Glu Glu Pro Thr Ile Phe Gln Glu Asp Ala Gly
 20           25           30
Gly Phe Gly Gln Ser Val Val Gln Phe Gly Gly Ser Arg Leu Val Val
 35           40           45
Gly Ala Pro Leu Glu Val Val Ala Ala Asn Gln Thr Gly Arg Leu Tyr
 50           55           60
Asp Cys Ala Ala Ala Thr Gly Met Cys Gln Pro Ile Pro Leu His Ile
 65           70           75           80
Arg Pro Glu Ala Val Asn Met Ser Leu Gly Leu Thr Leu Ala Ala Ser
 85           90           95
Thr Asn Gly Ser Arg Leu Leu Ala Cys Gly Pro Thr Leu His Arg Val
100          105          110
Cys Gly Glu Asn Ser Tyr Ser Lys Gly Ser Cys Leu Leu Leu Gly Ser
115          120          125
Arg Trp Glu Ile Ile Gln Thr Val Pro Asp Ala Thr Pro Glu Cys Pro
130          135          140
His Gln Glu Met Asp Ile Val Phe Leu Ile Asp Gly Ser Gly Ser Ile
145          150          155          160
Asp Gln Asn Asp Phe Asn Gln Met Lys Gly Phe Val Gln Ala Val Met
165          170          175
Gly Gln Phe Glu Gly Thr Asp Thr Leu Phe Ala Leu Met Gln Tyr Ser
180          185          190
Asn Leu Leu Lys Ile His Phe Thr Phe Thr Gln Phe Arg Thr Ser Pro
195          200          205
Ser Gln Gln Ser Leu Val Asp Pro Ile Val Gln Leu Lys Gly Leu Thr
210          215          220
Phe Thr Ala Thr Gly Ile Leu Thr Val Val Thr Gln Leu Phe His His
225          230          235          240
Lys Asn Gly Ala Arg Lys Ser Ala Lys Lys Ile Leu Ile Val Ile Thr
245          250          255

```

- 88 -

Asp Gly Gln Lys Tyr Lys Asp Pro Leu Glu Tyr Ser Asp Val Ile Pro  
 260 265 270  
 Gln Ala Glu Lys Ala Gly Ile Ile Arg Tyr Ala Ile Gly Val Gly His  
 275 280 285  
 Ala Phe Gln Gly Pro Thr Ala Arg Gln Glu Leu Asn Thr Ile Ser Ser  
 290 295 300  
 Ala Pro Pro Gln Asp His Val Phe Lys Val Asp Asn Phe Ala Ala Leu  
 305 310 315 320  
 Gly Ser Ile Gln Lys Gln Leu Gln Glu Lys Ile Tyr Ala Val Glu Gly  
 325 330 335  
 Thr Gln Ser Arg Ala Ser Ser Ser Phe Gln His Glu Met Ser Gln Glu  
 340 345 350  
 Gly Phe Ser Thr Ala Leu Thr Met Asp Gly Leu Phe Leu Gly Ala Val  
 355 360 365  
 Gly Ser Phe Ser Trp Ser Gly Gly Ala Phe Leu Tyr Pro Pro Asn Met  
 370 375 380  
 Ser Pro Thr Phe Ile Asn Met Ser Gln Glu Asn Val Asp Met Arg Asp  
 385 390 395 400  
 Ser Tyr Leu Gly Tyr Ser Thr Glu Leu Ala Leu Trp Lys Gly Val Gln  
 405 410 415  
 Asn Leu Val Leu Gly Ala Pro Arg Tyr Gln His Thr Gly Lys Ala Val  
 420 425 430  
 Ile Phe Thr Gln Val Ser Arg Gln Trp Arg Lys Lys Ala Glu Val Thr  
 435 440 445  
 Gly Thr Gln Ile Gly Ser Tyr Phe Gly Ala Ser Leu Cys Ser Val Asp  
 450 455 460  
 Val Asp Ser Asp Gly Ser Thr Asp Leu Ile Leu Ile Gly Ala Pro His  
 465 470 475 480  
 Tyr Tyr Glu Gln Thr Arg Gly Gly Gln Val Ser Val Cys Pro Leu Pro  
 485 490 495  
 Arg Gly Gln Arg Val Gln Trp Gln Cys Asp Ala Val Leu Arg Gly Glu  
 500 505 510  
 Gln Gly His Pro Trp Gly Arg Phe Gly Ala Ala Leu Thr Val Leu Gly  
 515 520 525  
 Asp Val Asn Glu Asp Lys Leu Ile Asp Val Ala Ile Gly Ala Pro Gly  
 530 535 540  
 Glu Gln Glu Asn Arg Gly Ala Val Tyr Leu Phe His Gly Ala Ser Glu  
 545 550 555 560  
 Ser Gly Ile Ser Pro Ser His Ser Gln Arg Ile Ala Ser Ser Gln Leu  
 565 570 575

- 89 -

Ser Pro Arg Leu Gln Tyr Phe Gly Gln Ala Leu Ser Gly Gly Gln Asp  
 580 585 590  
 Leu Thr Gln Asp Gly Leu Met Asp Leu Ala Val Gly Ala Arg Gly Gln  
 595 600 605  
 Val Leu Leu Leu Arg Ser Leu Pro Val Leu Lys Val Gly Val Ala Met  
 610 615 620  
 Arg Phe Ser Pro Val Glu Val Ala Lys Ala Val Tyr Arg Cys Trp Glu  
 625 630 635 640  
 Glu Lys Pro Ser Ala Leu Glu Ala Gly Asp Ala Thr Val Cys Leu Thr  
 645 650 655  
 Ile Gln Lys Ser Ser Leu Asp Gln Leu Gly Asp Ile Gln Ser Ser Val  
 660 665 670  
 Arg Phe Asp Leu Ala Leu Asp Pro Gly Arg Leu Thr Ser Arg Ala Ile  
 675 680 685  
 Phe Asn Glu Thr Lys Asn Pro Thr Leu Thr Arg Arg Lys Thr Leu Gly  
 690 695 700  
 Leu Gly Ile His Cys Glu Thr Leu Lys Leu Leu Leu Pro Asp Cys Val  
 705 710 715 720  
 Glu Asp Val Val Ser Pro Ile Ile Leu His Leu Asn Phe Ser Leu Val  
 725 730 735  
 Arg Glu Pro Ile Pro Ser Pro Gln Asn Leu Arg Pro Val Leu Ala Val  
 740 745 750  
 Gly Ser Gln Asp Leu Phe Thr Ala Ser Leu Pro Phe Glu Lys Asn Cys  
 755 760 765  
 Gly Gln Asp Gly Leu Cys Glu Gly Asp Leu Gly Val Thr Leu Ser Phe  
 770 775 780  
 Ser Gly Leu Gln Thr Leu Thr Val Gly Ser Ser Leu Glu Leu Asn Val  
 785 790 795 800  
 Ile Val Thr Val Trp Asn Ala Gly Glu Asp Ser Tyr Gly Thr Val Val  
 805 810 815  
 Ser Leu Tyr Tyr Pro Ala Gly Leu Ser His Arg Arg Val Ser Gly Ala  
 820 825 830  
 Gln Lys Gln Pro His Gln Ser Ala Leu Arg Leu Ala Cys Glu Thr Val  
 835 840 845  
 Pro Thr Glu Asp Glu Gly Leu Arg Ser Ser Arg Cys Ser Val Asn His  
 850 855 860  
 Pro Ile Phe His Glu Gly Ser Asn Gly Thr Phe Ile Val Thr Phe Asp  
 865 870 875 880  
 Val Ser Tyr Lys Ala Thr Leu Gly Asp Arg Met Leu Met Arg Ala Ser  
 885 890 895

- 90 -

Ala Ser Ser Glu Asn Asn Lys Ala Ser Ser Ser Lys Ala Thr Phe Gln  
                   900                                  905                                  910  
 Leu Glu Leu Pro Val Lys Tyr Ala Val Tyr Thr Met Ile Ser Arg Gln  
                   915                                  920                                  925  
 Glu Glu Ser Thr Lys Tyr Phe Asn Phe Ala Thr Ser Asp Glu Lys Lys  
                   930                                  935                                  940  
 Met Lys Glu Ala Glu His Arg Tyr Arg Val Asn Asn Leu Ser Gln Arg  
                   945                                  950                                  955                                  960  
 Asp Leu Ala Ile Ser Ile Asn Phe Trp Val Pro Val Leu Leu Asn Gly  
                                   965                                  970                                  975  
 Val Ala Val Trp Asp Val Val Met Glu Ala Pro Ser Gln Ser Leu Pro  
                                   980                                  985                                  990  
 Cys Val Ser Glu Arg Lys Pro Pro Gln His Ser Asp Phe Leu Thr Gln  
                   995                                  1000                                  1005  
 Ile Ser Arg Ser Pro Met Leu Asp Cys Ser Ile Ala Asp Cys Leu Gln  
                   1010                                  1015                                  1020  
 Phe Arg Cys Asp Val Pro Ser Phe Ser Val Gln Glu Glu Leu Asp Phe  
                   1025                                  1030                                  1035                                  1040  
 Thr Leu Lys Gly Asn Leu Ser Phe Gly Trp Val Arg Glu Thr Leu Gln  
                                   1045                                  1050                                  1055  
 Lys Lys Val Leu Val Val Ser Val Ala Glu Ile Thr Phe Asp Thr Ser  
                                   1060                                  1065                                  1070  
 Val Tyr Ser Gln Leu Pro Gly Gln Glu Ala Phe Met Arg Ala Gln Met  
                   1075                                  1080                                  1085  
 Glu Met Val Leu Glu Glu Asp Glu Val Tyr Asn Ala Ile Pro Ile Ile  
                   1090                                  1095                                  1100  
 Met Gly Ser Ser Val Gly Ala Leu Leu Leu Leu Ala Leu Ile Thr Ala  
                   1105                                  1110                                  1115                                  1120  
 Thr Leu Tyr Lys Leu Gly Phe Phe Lys Arg His Tyr Lys Glu Met Leu  
                                   1125                                  1130                                  1135  
 Glu Asp Lys Pro Glu Asp Thr Ala Thr Phe Ser Gly Asp Asp Phe Ser  
                   1140                                  1145                                  1150  
 Cys Val Ala Pro Asn Val Pro Lys Ser  
                   1155                                  1160

## (2) INFORMATION FOR SEQ ID NO:3:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1153 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: protein

- 91 -

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

Met	Ala	Leu	Arg	Val	Leu	Leu	Leu	Thr	Ala	Leu	Thr	Leu	Cys	His	Gly	1	5	10	15
Phe	Asn	Leu	Asp	Thr	Glu	Asn	Ala	Met	Thr	Phe	Gln	Glu	Asn	Ala	Arg	20	25	30	
Gly	Phe	Gly	Gln	Ser	Val	Val	Gln	Leu	Gln	Gly	Ser	Arg	Val	Val	Val	35	40	45	
Gly	Ala	Pro	Gln	Glu	Ile	Val	Ala	Ala	Asn	Gln	Arg	Gly	Ser	Leu	Tyr	50	55	60	
Gln	Cys	Asp	Tyr	Ser	Thr	Gly	Ser	Cys	Glu	Pro	Ile	Arg	Leu	Gln	Val	65	70	75	80
Pro	Val	Glu	Ala	Val	Asn	Met	Ser	Leu	Gly	Leu	Ser	Leu	Ala	Ala	Thr	85	90	95	
Thr	Ser	Pro	Pro	Gln	Leu	Leu	Ala	Cys	Gly	Pro	Thr	Val	His	Gln	Thr	100	105	110	
Cys	Ser	Glu	Asn	Thr	Tyr	Val	Lys	Gly	Leu	Cys	Phe	Leu	Phe	Gly	Ser	115	120	125	
Asn	Leu	Arg	Gln	Gln	Pro	Gln	Lys	Phe	Pro	Glu	Ala	Leu	Arg	Gly	Cys	130	135	140	
Pro	Gln	Glu	Asp	Ser	Asp	Ile	Ala	Phe	Leu	Ile	Asp	Gly	Ser	Gly	Ser	145	150	155	160
Ile	Ile	Pro	His	Asp	Phe	Arg	Arg	Met	Lys	Glu	Phe	Val	Ser	Thr	Val	165	170	175	
Met	Glu	Gln	Leu	Lys	Lys	Ser	Lys	Thr	Leu	Phe	Ser	Leu	Met	Gln	Tyr	180	185	190	
Ser	Glu	Glu	Phe	Arg	Ile	His	Phe	Thr	Phe	Lys	Glu	Phe	Gln	Asn	Asn	195	200	205	
Pro	Asn	Pro	Arg	Ser	Leu	Val	Lys	Pro	Ile	Thr	Gln	Leu	Leu	Gly	Arg	210	215	220	
Thr	His	Thr	Ala	Thr	Gly	Ile	Arg	Lys	Val	Val	Arg	Glu	Leu	Phe	Asn	225	230	235	240
Ile	Thr	Asn	Gly	Ala	Arg	Lys	Asn	Ala	Phe	Lys	Ile	Leu	Val	Val	Ile	245	250	255	
Thr	Asp	Gly	Glu	Lys	Phe	Gly	Asp	Pro	Leu	Gly	Tyr	Glu	Asp	Val	Ile	260	265	270	
Pro	Glu	Ala	Asp	Arg	Glu	Gly	Val	Ile	Arg	Tyr	Val	Ile	Gly	Val	Gly	275	280	285	
Asp	Ala	Phe	Arg	Ser	Glu	Lys	Ser	Arg	Gln	Glu	Leu	Asn	Thr	Ile	Ala	290	295	300	

- 92 -

Ser Lys Pro Pro Arg Asp His Val Phe Gln Val Asn Asn Phe Glu Ala  
 305 310 315 320  
 Leu Lys Thr Ile Gln Asn Gln Leu Arg Glu Lys Ile Phe Ala Ile Glu  
 325 330 335  
 Gly Thr Gln Thr Gly Ser Ser Ser Ser Phe Glu His Glu Met Ser Gln  
 340 345 350  
 Glu Gly Phe Ser Ala Ala Ile Thr Ser Asn Gly Pro Leu Leu Ser Thr  
 355 360 365  
 Val Gly Ser Tyr Asp Trp Ala Gly Gly Val Phe Leu Tyr Thr Ser Lys  
 370 375 380  
 Glu Lys Ser Thr Phe Ile Asn Met Thr Arg Val Asp Ser Asp Met Asn  
 385 390 395 400  
 Asp Ala Tyr Leu Gly Tyr Ala Ala Ala Ile Ile Leu Arg Asn Arg Val  
 405 410 415  
 Gln Ser Leu Val Leu Gly Ala Pro Arg Tyr Gln His Ile Gly Leu Val  
 420 425 430  
 Ala Met Phe Arg Gln Asn Thr Gly Met Trp Glu Ser Asn Ala Asn Val  
 435 440 445  
 Lys Gly Thr Gln Ile Gly Ala Tyr Phe Gly Ala Ser Leu Cys Ser Val  
 450 455 460  
 Asp Val Asp Ser Asn Gly Ser Thr Asp Leu Val Leu Ile Gly Ala Pro  
 465 470 475 480  
 His Tyr Tyr Glu Gln Thr Arg Gly Gly Gln Val Ser Val Cys Pro Leu  
 485 490 495  
 Pro Arg Gly Gln Arg Ala Arg Trp Gln Cys Asp Ala Val Leu Tyr Gly  
 500 505 510  
 Glu Gln Gly Gln Pro Trp Gly Arg Phe Gly Ala Ala Leu Thr Val Leu  
 515 520 525  
 Gly Asp Val Asn Gly Asp Lys Leu Thr Asp Val Ala Ile Gly Ala Pro  
 530 535 540  
 Gly Glu Glu Asp Asn Arg Gly Ala Val Tyr Leu Phe His Gly Thr Ser  
 545 550 555 560  
 Gly Ser Gly Ile Ser Pro Ser His Ser Gln Arg Ile Ala Gly Ser Lys  
 565 570 575  
 Leu Ser Pro Arg Leu Gln Tyr Phe Gly Gln Ser Leu Ser Gly Gly Gln  
 580 585 590  
 Asp Leu Thr Met Asp Gly Leu Val Asp Leu Thr Val Gly Ala Gln Gly  
 595 600 605  
 His Val Leu Leu Leu Arg Ser Gln Pro Val Leu Arg Val Lys Ala Ile  
 610 615 620



- 93 -

Met Glu Phe Asn Pro Arg Glu Val Ala Arg Asn Val Phe Glu Cys Asn  
 625 630 635 640  
 Asp Gln Val Val Lys Gly Lys Glu Ala Gly Glu Val Arg Val Cys Leu  
 645 650 655  
 His Val Gln Lys Ser Thr Arg Asp Arg Leu Arg Glu Gly Gln Ile Gln  
 660 665 670  
 Ser Val Val Thr Tyr Asp Leu Ala Leu Asp Ser Gly Arg Pro His Ser  
 675 680 685  
 Arg Ala Val Phe Asn Glu Thr Lys Asn Ser Thr Arg Arg Gln Thr Gln  
 690 695 700  
 Val Leu Gly Leu Thr Gln Thr Cys Glu Thr Leu Lys Leu Gln Leu Pro  
 705 710 715 720  
 Asn Cys Ile Glu Asp Pro Val Ser Pro Ile Val Leu Arg Leu Asn Phe  
 725 730 735  
 Ser Leu Val Gly Thr Pro Leu Ser Ala Phe Gly Asn Leu Arg Pro Val  
 740 745 750  
 Leu Ala Glu Asp Ala Gln Arg Leu Phe Thr Ala Leu Phe Pro Phe Glu  
 755 760 765  
 Lys Asn Cys Gly Asn Asp Asn Ile Cys Gln Asp Asp Leu Ser Ile Thr  
 770 775 780  
 Phe Ser Phe Met Ser Leu Asp Cys Leu Val Val Gly Gly Pro Arg Glu  
 785 790 795 800  
 Phe Asn Val Thr Val Thr Val Arg Asn Asp Gly Glu Asp Ser Tyr Arg  
 805 810 815  
 Thr Gln Val Thr Phe Phe Phe Pro Leu Asp Leu Ser Tyr Arg Lys Val  
 820 825 830  
 Ser Thr Leu Gln Asn Gln Arg Ser Gln Arg Ser Trp Arg Leu Ala Cys  
 835 840 845  
 Glu Ser Ala Ser Ser Thr Glu Val Ser Gly Ala Leu Lys Ser Thr Ser  
 850 855 860  
 Cys Ser Ile Asn His Pro Ile Phe Pro Glu Asn Ser Glu Val Thr Phe  
 865 870 875 880  
 Asn Ile Thr Phe Asp Val Asp Ser Lys Ala Ser Leu Gly Asn Lys Leu  
 885 890 895  
 Leu Leu Lys Ala Asn Val Thr Ser Glu Asn Asn Met Pro Arg Thr Asn  
 900 905 910  
 Lys Thr Glu Phe Gln Leu Glu Leu Pro Val Lys Tyr Ala Val Tyr Met  
 915 920 925  
 Val Val Thr Ser His Gly Val Ser Thr Lys Tyr Leu Asn Phe Thr Ala  
 930 935 940

- 94 -

Ser Glu Asn Thr Ser Arg Val Met Gln His Gln Tyr Gln Val Ser Asn  
 945 950 955 960  
 Leu Gly Gln Arg Ser Leu Pro Ile Ser Leu Val Phe Leu Val Pro Val  
 965 970 975  
 Arg Leu Asn Gln Thr Val Ile Trp Asp Arg Pro Gln Val Thr Phe Ser  
 980 985 990  
 Glu Asn Leu Ser Ser Thr Cys His Thr Lys Glu Arg Leu Pro Ser His  
 995 1000 1005  
 Ser Asp Phe Leu Ala Glu Leu Arg Lys Ala Pro Val Val Asn Cys Ser  
 1010 1015 1020  
 Ile Ala Val Cys Gln Arg Ile Gln Cys Asp Ile Pro Phe Phe Gly Ile  
 1025 1030 1035 1040  
 Gln Glu Glu Phe Asn Ala Thr Leu Lys Gly Asn Leu Ser Phe Asp Trp  
 1045 1050 1055  
 Tyr Ile Lys Thr Ser His Asn His Leu Leu Ile Val Ser Thr Ala Glu  
 1060 1065 1070  
 Ile Leu Phe Asn Asp Ser Val Phe Thr Leu Leu Pro Gly Gln Gly Ala  
 1075 1080 1085  
 Phe Val Arg Ser Gln Thr Glu Thr Lys Val Glu Pro Phe Glu Val Pro  
 1090 1095 1100  
 Asn Pro Leu Pro Leu Ile Val Gly Ser Ser Val Gly Gly Leu Leu Leu  
 1105 1110 1115 1120  
 Leu Ala Leu Ile Thr Ala Ala Leu Tyr Lys Leu Gly Phe Phe Lys Arg  
 1125 1130 1135  
 Gln Tyr Lys Asp Met Met Ser Glu Gly Gly Pro Pro Gly Ala Glu Pro  
 1140 1145 1150  
 Gln

## (2) INFORMATION FOR SEQ ID NO:4:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1163 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

Met Thr Arg Thr Arg Ala Ala Leu Leu Leu Phe Thr Ala Leu Ala Thr  
 1 5 10 15  
 Ser Leu Gly Phe Asn Leu Asp Thr Glu Leu Thr Ala Phe Arg Val  
 20 25 30

- 95 -

Asp Ser Ala Gly Phe Gly Asp Ser Val Val Gln Tyr Ala Asn Ser Trp  
 35 40 45  
 Val Val Val Gly Ala Pro Gln Lys Ile Ile Ala Ala Asn Gln Ile Gly  
 50 55 60  
 Gly Leu Tyr Gln Cys Gly Tyr Ser Thr Gly Ala Cys Glu Pro Ile Gly  
 65 70 75 80  
 Leu Gln Val Pro Pro Glu Ala Val Asn Met Ser Leu Gly Leu Ser Leu  
 85 90 95  
 Ala Ser Thr Thr Ser Pro Ser Gln Leu Leu Ala Cys Gly Pro Thr Val  
 100 105 110  
 His His Glu Cys Gly Arg Asn Met Tyr Leu Thr Gly Leu Cys Phe Leu  
 115 120 125  
 Leu Gly Pro Thr Gln Leu Thr Gln Arg Leu Pro Val Ser Arg Gln Glu  
 130 135 140  
 Cys Pro Arg Gln Glu Gln Asp Ile Val Phe Leu Ile Asp Gly Ser Gly  
 145 150 155 160  
 Ser Ile Ser Ser Arg Asn Phe Ala Thr Met Met Asn Phe Val Arg Ala  
 165 170 175  
 Val Ile Ser Gln Phe Gln Arg Pro Ser Thr Gln Phe Ser Leu Met Gln  
 180 185 190  
 Phe Ser Asn Lys Phe Gln Thr His Phe Thr Phe Glu Glu Phe Arg Arg  
 195 200 205  
 Thr Ser Asn Pro Leu Ser Leu Leu Ala Ser Val His Gln Leu Gln Gly  
 210 215 220  
 Phe Thr Tyr Thr Ala Thr Ala Ile Gln Asn Val Val His Arg Leu Phe  
 225 230 235 240  
 His Ala Ser Tyr Gly Ala Arg Arg Asp Ala Ile Lys Ile Leu Ile Val  
 245 250 255  
 Ile Thr Asp Gly Lys Lys Glu Gly Asp Ser Leu Asp Tyr Lys Asp Val  
 260 265 270  
 Ile Pro Met Ala Asp Ala Ala Gly Ile Ile Arg Tyr Ala Ile Gly Val  
 275 280 285  
 Gly Leu Ala Phe Gln Asn Arg Asn Ser Trp Lys Glu Leu Asn Asp Ile  
 290 295 300  
 Ala Ser Lys Pro Ser Gln Glu His Ile Phe Lys Val Glu Asp Phe Asp  
 305 310 315 320  
 Ala Leu Lys Asp Ile Gln Asn Gln Leu Lys Glu Lys Ile Phe Ala Ile  
 325 330 335  
 Glu Gly Thr Glu Thr Ile Ser Ser Ser Ser Phe Glu Leu Glu Met Ala  
 340 345 350

- 96 -

Gln Glu Gly Phe Ser Ala Val Phe Thr Pro Asp Gly Pro Val Leu Gly  
 355 360 365  
 Ala Val Gly Ser Phe Thr Trp Ser Gly Gly Ala Phe Leu Tyr Pro Pro  
 370 375 380  
 Asn Met Ser Pro Thr Phe Ile Asn Met Ser Gln Glu Asn Val Asp Met  
 385 390 395 400  
 Arg Asp Ser Tyr Leu Gly Tyr Ser Thr Glu Leu Ala Leu Trp Lys Gly  
 405 410 415  
 Val Gln Ser Leu Val Leu Gly Ala Pro Arg Tyr Gln His Ile Gly Lys  
 420 425 430  
 Ala Val Ile Phe Ile Gln Val Ser Arg Gln Trp Arg Met Lys Ala Glu  
 435 440 445  
 Val Ile Gly Thr Gln Ile Gly Ser Tyr Phe Gly Ala Ser Leu Cys Ser  
 450 455 460  
 Val Asp Val Asp Thr Asp Gly Ser Thr Asp Leu Val Leu Ile Gly Ala  
 465 470 475 480  
 Pro His Tyr Tyr Glu Gln Thr Arg Gly Gly Gln Val Ser Val Cys Pro  
 485 490 495  
 Leu Pro Arg Gly Trp Arg Arg Trp Trp Cys Asp Ala Val Leu Tyr Gly  
 500 505 510  
 Glu Gln Gly His Pro Trp Gly Arg Phe Gly Ala Ala Leu Thr Val Leu  
 515 520 525  
 Gly Asp Val Asn Gly Asp Lys Leu Thr Asp Val Val Ile Gly Ala Pro  
 530 535 540  
 Gly Glu Glu Glu Asn Arg Gly Ala Val Tyr Leu Phe His Gly Val Leu  
 545 550 555 560  
 Gly Pro Ser Ile Ser Pro Ser His Ser Gln Arg Ile Ala Gly Ser Gln  
 565 570 575  
 Leu Ser Ser Arg Leu Gln Tyr Phe Gly Gln Ala Leu Ser Gly Gly Gln  
 580 585 590  
 Asp Leu Thr Gln Asp Gly Leu Val Asp Leu Ala Val Gly Ala Arg Gly  
 595 600 605  
 Gln Val Leu Leu Leu Arg Thr Arg Pro Val Leu Trp Val Gly Val Ser  
 610 615 620  
 Met Gln Phe Ile Pro Ala Glu Ile Pro Arg Ser Ala Phe Glu Cys Arg  
 625 630 635 640  
 Glu Gln Val Val Ser Glu Gln Thr Leu Val Gln Ser Asn Ile Cys Leu  
 645 650 655  
 Tyr Ile Asp Lys Arg Ser Lys Asn Leu Leu Gly Ser Arg Asp Leu Gln  
 660 665 670

- 97 -

Ser Ser Val Thr Leu Asp Leu Ala Leu Ala Pro Gly Arg Leu Ser Pro  
 675 680 685  
 Arg Ala Ile Phe Gln Glu Thr Lys Asn Arg Ser Leu Ser Arg Val Arg  
 690 695 700  
 Val Leu Gly Leu Lys Ala His Cys Glu Asn Phe Asn Leu Leu Leu Pro  
 705 710 715 720  
 Ser Cys Val Glu Asp Ser Val Ile Pro Ile Ile Leu Arg Leu Asn Phe  
 725 730 735  
 Thr Leu Val Gly Lys Pro Leu Leu Ala Phe Arg Asn Leu Arg Pro Met  
 740 745 750  
 Leu Ala Ala Leu Ala Gln Arg Tyr Phe Thr Ala Ser Leu Pro Phe Glu  
 755 760 765  
 Lys Asn Cys Gly Ala Asp His Ile Cys Gln Asp Asn Leu Gly Ile Ser  
 770 775 780  
 Phe Ser Phe Pro Gly Leu Lys Ser Leu Leu Val Gly Ser Asn Leu Glu  
 785 790 795 800  
 Leu Asn Ala Glu Val Met Val Trp Asn Asp Gly Glu Asp Ser Tyr Gly  
 805 810 815  
 Thr Thr Ile Thr Phe Ser His Pro Ala Gly Leu Ser Tyr Arg Tyr Val  
 820 825 830  
 Ala Glu Gly Gln Lys Gln Gly Gln Leu Arg Ser Leu His Leu Thr Cys  
 835 840 845  
 Cys Ser Ala Pro Val Gly Ser Gln Gly Thr Trp Ser Thr Ser Cys Arg  
 850 855 860  
 Ile Asn His Leu Ile Phe Arg Gly Gly Ala Gln Ile Thr Phe Leu Ala  
 865 870 875 880  
 Thr Phe Asp Val Ser Pro Lys Ala Val Gly Leu Asp Arg Leu Leu Leu  
 885 890 895  
 Ile Ala Asn Val Ser Ser Glu Asn Asn Ile Pro Arg Thr Ser Lys Thr  
 900 905 910  
 Ile Phe Gln Leu Glu Leu Pro Val Lys Tyr Ala Val Tyr Ile Val Val  
 915 920 925  
 Ser Ser His Glu Gln Phe Thr Lys Tyr Leu Asn Phe Ser Glu Ser Glu  
 930 935 940  
 Glu Lys Glu Ser His Val Ala Met His Arg Tyr Gln Val Asn Asn Leu  
 945 950 955 960  
 Gly Gln Arg Asp Leu Pro Val Ser Ile Asn Phe Trp Val Pro Val Glu  
 965 970 975  
 Leu Asn Gln Glu Ala Val Trp Met Asp Val Glu Val Ser His Pro Gln  
 980 985 990

- 98 -

Asn Pro Ser Leu Arg Cys Ser Ser Glu Lys Ile Ala Pro Pro Ala Ser  
 995 1000 1005  
 Asp Phe Leu Ala His Ile Gln Lys Asn Pro Val Leu Asp Cys Ser Ile  
 1010 1015 1020  
 Ala Gly Cys Leu Arg Phe Arg Cys Asp Val Pro Ser Phe Ser Val Gln  
 1025 1030 1035 1040  
 Glu Glu Leu Asp Phe Thr Leu Lys Gly Asn Leu Ser Phe Gly Trp Val  
 1045 1050 1055  
 Arg Gln Ile Leu Gln Lys Lys Val Ser Val Val Ser Val Ala Glu Ile  
 1060 1065 1070  
 Ile Phe Asp Thr Ser Val Tyr Ser Gln Leu Pro Gly Gln Glu Ala Phe  
 1075 1080 1085  
 Met Arg Ala Gln Thr Ile Thr Val Leu Glu Lys Tyr Lys Val His Asn  
 1090 1095 1100  
 Pro Ile Pro Leu Ile Val Gly Ser Ser Ile Gly Gly Leu Leu Leu Leu  
 1105 1110 1115 1120  
 Ala Leu Ile Thr Ala Val Leu Tyr Lys Val Gly Phe Phe Lys Arg Gln  
 1125 1130 1135  
 Tyr Lys Glu Met Met Glu Glu Ala Asn Gly Gln Ile Ala Pro Glu Asn  
 1140 1145 1150  
 Gly Thr Gln Thr Pro Ser Pro Pro Ser Glu Lys  
 1155 1160

## (2) INFORMATION FOR SEQ ID NO:5:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 12 amino acids
  - (B) TYPE: amino acid
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

Phe Asn Leu Asp Val Glu Glu Pro Met Val Phe Gln  
 1 5 10

## (2) INFORMATION FOR SEQ ID NO:6:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 35 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

- 99 -

- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:  
TTYAAYYTGG AYGTINGARGA RCCNATGGTN TTYCA 35
- (2) INFORMATION FOR SEQ ID NO:7:
- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 36 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: DNA
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:  
TTCAACCTGG ACGTGGAGGA GCCCATGGTG TTCAA 36
- (2) INFORMATION FOR SEQ ID NO:8:
- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 36 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: DNA
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:  
TTCAACCTGG ACGTNGAASA NCCCATGGTC TTCAA 36
- (2) INFORMATION FOR SEQ ID NO:9:
- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 23 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: DNA
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:  
TTYAAYYTNG AYGTINGARGA RCC 23
- (2) INFORMATION FOR SEQ ID NO:10:
- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 20 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single

- 100 -

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:

TTYAAYYTGG ACGTNGAAGA

20

(2) INFORMATION FOR SEQ ID NO:11:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 17 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:11:

TGRAANACCA TNGGYTC

17

(2) INFORMATION FOR SEQ ID NO:12:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 18 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:12:

TTGGAAGACC ATNGGYTC

18

(2) INFORMATION FOR SEQ ID NO:13:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 17 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:13:

ATTAACCCTC ACTAAG

17



- 101 -

## (2) INFORMATION FOR SEQ ID NO:14:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 17 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:14:

AATACGACTC ACTATAG

17

## (2) INFORMATION FOR SEQ ID NO:15:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 11 amino acids
  - (B) TYPE: amino acid
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:15:

Val	Phe	Gln	Glu	Xaa	Gly	Ala	Gly	Phe	Gly	Gln
1				5					10	

## (2) INFORMATION FOR SEQ ID NO:16:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 14 amino acids
  - (B) TYPE: amino acid
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:16:

Leu	Tyr	Asp	Xaa	Val	Ala	Ala	Thr	Gly	Leu	Xaa	Gln	Pro	Ile
1				5					10				

## (2) INFORMATION FOR SEQ ID NO:17:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 12 amino acids
  - (B) TYPE: amino acid
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

- 102 -

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:17:

Pro Leu Glu Tyr Xaa Asp Val Ile Pro Gln Ala Glu  
1 5 10

(2) INFORMATION FOR SEQ ID NO:18:

(i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 10 amino acids  
(B) TYPE: amino acid  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:18:

Phe Gln Glu Gly Phe Ser Xaa Val Leu Xaa  
1 5 10

(2) INFORMATION FOR SEQ ID NO:19:

(i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 14 amino acids  
(B) TYPE: amino acid  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:19:

Thr Ser Pro Thr Phe Ile Xaa Met Ser Gln Glu Asn Val Asp  
1 5 10

(2) INFORMATION FOR SEQ ID NO:20:

(i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 17 amino acids  
(B) TYPE: amino acid  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:20:

Leu Val Val Gly Ala Pro Leu Glu Val Val Ala Val Xaa Gln Thr Gly  
1 5 10 15

Arg

(2) INFORMATION FOR SEQ ID NO:21:

(i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 9 amino acids  
(B) TYPE: amino acid

- 103 -

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:21:

Leu Asp Xaa Lys Pro Xaa Asp Thr Ala  
1 5

(2) INFORMATION FOR SEQ ID NO:22:

(i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 7 amino acids  
(B) TYPE: amino acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:22:

Phe Gly Glu Gln Phe Ser Glu  
1 5

(2) INFORMATION FOR SEQ ID NO:23:

(i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 21 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:23:

RAANCCYTCY TGRAAACTYT C

21

(2) INFORMATION FOR SEQ ID NO:24:

(i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 1006 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:24:

- 104 -

TTCAACCTGG	ACGTGGAGGA	GCCCATGGTG	TTCAAGAGGA	TGGAGCTGGC	TTTGGACAGA	60
GCGTGGCCCA	GCTTGGCGGA	TCTAGACTCG	TGGTGGGAGC	CCCCCTGGAG	GTGGTGGCGG	120
TCAACCAAAC	AGGAAGGTTG	TATGACTGTG	TGGCTGCCAC	TGGCCTTGTC	AACCCATACC	180
CCTGCACACA	CCCCCAGATG	CTGTGAACAT	GTCCCTGGGT	CTGTCCCTGT	CAGCCGCCGC	240
CAGTCGCCCC	TGGCTGCTGG	CCTGTGGCCC	AACCATGCAC	AGAGCCTGTG	GGGAGAATAT	300
GTATGCAGAA	GGCTTTTGCC	TCCTGTTGGA	CTCCCATCTG	CAGACCATT	GGACAGTACC	360
TGCTGCCCTA	CCAGAGTGTC	CAAGTCAAGA	GATGGACATT	GTCTTCCTGA	TTGATGGTTC	420
TGGCAGTATG	AGCAAAGTGA	CTTTAAACAA	ATGAAGGATT	TGTGAGAGCT	GTGATGGGAC	480
AGTTTGAGGG	CACCCAAACC	CTGTTCTCAC	TGATACAGTA	TCCCACCTCC	CTGAAGATCC	540
ACTTCACCTT	CACGCAATTC	CAGAGCAGCT	GGAACCCTCT	GAGCCTGGTG	GATCCCATTG	600
TCCAACCTGA	CGGCCTGACA	TATACAGCCA	CGGGCATCCG	GAAAGTGGTG	GAGGAACGTG	660
TTCATAGTAA	GAATGGGGCC	CGTAAAAGTG	CCAAGAAGAT	CCTCATTGTC	ATCACAGATG	720
GCAAAAATAC	AAAGACCCCC	TGGAGTACGA	GGACGTATCC	CCAGGCAGAG	AGAGCGGATC	780
ATCCGCTATG	CCATTGGGGT	GGGAGATGCT	TTCTGGAAAC	CCAGTGCCAA	GCAGGAGCTG	840
GACAACATTG	GCTCAGAGCC	GGCTCAGGAC	CATGTGTTCA	GGGTGGACAA	CTTTCAGCA	900
CTCAGCAGCA	TCCAGGAGCA	GCTGCAGGAG	AAGATCTTTG	CACTCGAAGG	AACCCAGTCG	960
ACGACAAGTA	GCTCTTTCCA	ACATGAGATG	TTCCAAGAAG	GGTTCA		1006

## (2) INFORMATION FOR SEQ ID NO:25:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 17 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:25:

GTNTTYCARG ARGAYGG

17

## (2) INFORMATION FOR SEQ ID NO:26:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 20 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

- 105 -

- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:26:  
CCACTGTCAG GATGCCCGTG 20
- (2) INFORMATION FOR SEQ ID NO:27:
- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 42 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: DNA
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:27:  
AGTTACGAAT TCGCCACCAT GGCTCTACGG GTGCTTCTTC TG 42
- (2) INFORMATION FOR SEQ ID NO:28:
- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 42 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: DNA
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:28:  
AGTTACGAAT TCGCCACCAT GACTCGGACT GTGCTTCTTC TG 42
- (2) INFORMATION FOR SEQ ID NO:29:
- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 36 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: DNA
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:29:  
AGTTACGAAT TCGCCACCAT GACCTTCGGC ACTGTG 36
- (2) INFORMATION FOR SEQ ID NO:30:
- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 20 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: DNA

- 106 -

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:30:

TTGCTGACTG CCTGCAGTTC

20

(2) INFORMATION FOR SEQ ID NO:31:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 36 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:31:

GTTCTGACGC GTAATGGCAT TGTAGACCTC GTCTTC

36

(2) INFORMATION FOR SEQ ID NO:32:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 36 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:32:

ACGTATGCAG GATCCCATCA AGAGATGGAC ATCGCT

36

(2) INFORMATION FOR SEQ ID NO:33:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 37 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:33:

ACTGCATGTC TCGAGGCTGA AGCCTTCTTG GGACATC

37

(2) INFORMATION FOR SEQ ID NO:34:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 24 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

- 107 -

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:34:

TATAGACTGC TGGGTAGTCC CCAC

24

(2) INFORMATION FOR SEQ ID NO:35:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 24 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:35:

TGAAGATTGG GGGTAAATAA CAGA

24

(2) INFORMATION FOR SEQ ID NO:36:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 3528 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(ix) FEATURE:

- (A) NAME/KEY: CDS
- (B) LOCATION: 1..3456

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:36:

GGC	TGG	GCC	CTG	GCT	TCC	TGT	CAT	GGG	TCT	AAC	CTG	GAT	GTG	GAG	GAA	48
Gly	Trp	Ala	Leu	Ala	Ser	Cys	His	Gly	Ser	Asn	Leu	Asp	Val	Glu	Glu	
1				5				10						15		
CCC	ATC	GTG	TTC	AGA	GAG	GAT	GCA	GCC	AGC	TTT	GGA	CAG	ACT	GTG	GTG	96
Pro	Ile	Val	Phe	Arg	Glu	Asp	Ala	Ala	Ser	Phe	Gly	Gln	Thr	Val	Val	
			20				25						30			
CAG	TTT	GGT	GGA	TCT	CGA	CTC	GTG	GTG	GGA	GCC	CCT	CTG	GAG	GCG	GTG	144
Gln	Phe	Gly	Gly	Ser	Arg	Leu	Val	Val	Gly	Ala	Pro	Leu	Glu	Ala	Val	
		35				40						45				
GCA	GTC	AAC	CAA	ACA	GGA	CGG	TTG	TAT	GAC	TGT	GCA	CCT	GCC	ACT	GGC	192
Ala	Val	Asn	Gln	Thr	Gly	Arg	Leu	Tyr	Asp	Cys	Ala	Pro	Ala	Thr	Gly	
50					55						60					

- 108 -

ATG TGC CAG CCC ATC GTA CTG CGC AGT CCC CTA GAG GCA GTG AAC ATG Met Cys Gln Pro Ile Val Leu Arg Ser Pro Leu Glu Ala Val Asn Met 65 70 75 80	240
TCC CTG GGC CTG TCT CTG GTG ACT GCC ACC AAT AAC GCC CAG TTG CTG Ser Leu Gly Leu Ser Leu Val Thr Ala Thr Asn Asn Ala Gln Leu Leu 85 90 95	288
GCT TGT GGT CCA ACT GCA CAG AGA GCT TGT GTG AAG AAC ATG TAT GCG Ala Cys Gly Pro Thr Ala Gln Arg Ala Cys Val Lys Asn Met Tyr Ala 100 105 110	336
AAA GGT TCC TGC CTC CTT CTC GGC TCC AGC TTG CAG TTC ATC CAG GCA Lys Gly Ser Cys Leu Leu Leu Gly Ser Ser Leu Gln Phe Ile Gln Ala 115 120 125	384
GTC CCT GCC TCC ATG CCA GAG TGT CCA AGA CAA GAG ATG GAC ATT GCT Val Pro Ala Ser Met Pro Glu Cys Pro Arg Gln Glu Met Asp Ile Ala 130 135 140	432
TTC CTG ATT GAT GGT TCT GGC AGC ATT AAC CAA AGG GAC TTT GCC CAG Phe Leu Ile Asp Gly Ser Gly Ser Ile Asn Gln Arg Asp Phe Ala Gln 145 150 155 160	480
ATG AAG GAC TTT GTC AAA GCT TTG ATG GGA GAG TTT GCG AGC ACC AGC Met Lys Asp Phe Val Lys Ala Leu Met Gly Glu Phe Ala Ser Thr Ser 165 170 175	528
ACC TTG TTC TCC CTG ATG CAA TAC TCG AAC ATC CTG AAG ACC CAT TTT Thr Leu Phe Ser Leu Met Gln Tyr Ser Asn Ile Leu Lys Thr His Phe 180 185 190	576
ACC TTC ACT GAA TTC AAG AAC ATC CTG GAC CCT CAG AGC CTG GTG GAT Thr Phe Thr Glu Phe Lys Asn Ile Leu Asp Pro Gln Ser Leu Val Asp 195 200 205	624
CCC ATT GTC CAG CTG CAA GGC CTG ACC TAC ACA GCC ACA GGC ATC CGG Pro Ile Val Gln Leu Gln Gly Leu Thr Tyr Thr Ala Thr Gly Ile Arg 210 215 220	672
ACA GTG ATG GAA GAG CTA TTT CAT AGC AAG AAT GGG TCC CGT AAA AGT Thr Val Met Glu Glu Leu Phe His Ser Lys Asn Gly Ser Arg Lys Ser 225 230 235 240	720
GCC AAG AAG ATC CTC CTT GTC ATC ACA GAT GGG CAG AAA TAC AGA GAC Ala Lys Lys Ile Leu Leu Val Ile Thr Asp Gly Gln Lys Tyr Arg Asp 245 250 255	768
CCC CTG GAG TAT AGT GAT GTC ATT CCC GCC GCA GAC AAA GCT GGC ATC Pro Leu Glu Tyr Ser Asp Val Ile Pro Ala Ala Asp Lys Ala Gly Ile 260 265 270	816
ATT CGT TAT GCT ATT GGG GTG GGA GAT GCC TTC CAG GAG CCC ACT GCC Ile Arg Tyr Ala Ile Gly Val Gly Asp Ala Phe Gln Glu Pro Thr Ala 275 280 285	864
CTG AAG GAG CTG AAC ACC ATT GGC TCA GCT CCC CCA CAG GAC CAC GTG Leu Lys Glu Leu Asn Thr Ile Gly Ser Ala Pro Pro Gln Asp His Val 290 295 300	912



- 109 -

TTC Phe 305	AAG Lys	GTA Val	GGC Gly	AAC Asn	TTT Phe 310	GCA Ala	GCA Ala	CTT Leu	CGC Arg	AGC Ser 315	ATC Ile	CAG Gln	AGG Arg	CAA Gln 320	CTT Leu	960
CAG Gln	GAG Glu	AAA Lys	ATC Ile	TTT Phe 325	GCC Ala	ATT Ile	GAG Glu	GGA Gly	ACT Thr 330	CAA Gln	TCA Ser	AGG Arg	TCA Ser	AGT Ser 335	AGT Ser	1008
TCC Ser	TTT Phe	CAG Gln	CAC His 340	GAG Glu	ATG Met	TCA Ser	CAA Gln	GAA Glu 345	GGT Gly	TTT Phe	AGT Ser	TCA Ser	GCT Ala 350	CTC Leu	ACA Thr	1056
TCG Ser	GAT Asp	GGA Gly 355	CCC Pro	GTT Val	CTG Leu	GGG Gly	GCC Ala 360	GYG Xaa	GGA Gly	AGC Ser	TTT Phe	AGC Ser 365	TGG Trp	TCC Ser	GGA Gly	1104
GGT Gly 370	GCC Ala	TTT Phe	TTA Leu	TAT Tyr	CCC Pro	CCA Pro 375	AAT Asn	ACG Thr	AGA Arg	CCC Pro	ACC Thr 380	TTT Phe	ATC Ile	AAC Asn	ATG Met	1152
TCT Ser 385	CAG Gln	GAG Glu	AAT Asn	GTG Val	GAC Asp 390	ATG Met	AGA Arg	GAC Asp	TCC Ser	TAC Tyr 395	CTG Leu	GGT Gly	TAC Tyr	TCC Ser	ACC Thr 400	1200
GCA Ala	GTG Val	GCC Ala	TTT Phe	TGG Trp 405	AAG Lys	GGG Gly	GTT Val	CAC His	AGC Ser 410	CTG Leu	ATC Ile	CTG Leu	GGG Gly	GCC Ala 415	CCG Pro	1248
CGT Arg	CAC His	CAG Gln	CAC His 420	ACG Thr	GGG Gly	AAG Lys	GTT Val	GTC Val 425	ATC Ile	TTT Phe	ACC Thr	CAG Gln	GAA Glu 430	GCC Ala	AGG Arg	1296
CAT His	TGG Trp	AGG Arg 435	CCC Pro	AAG Lys	TCT Ser	GAA Glu	GTC Val 440	AGA Arg	GGG Gly	ACA Thr	CAG Gln	ATC Ile 445	GGC Gly	TCC Ser	TAC Tyr	1344
TTC Phe 450	GGG Gly	GCC Ala	TCT Ser	CTC Leu	TGT Cys	TCT Ser 455	GTG Val	GAC Asp	GTG Val	GAT Asp 460	AGA Arg	GAT Asp	GGC Gly	AGC Ser	ACY Xaa	1392
GAC Asp 465	CTG Leu	GTC Val	CTG Leu	ATC Ile	GGA Gly 470	GCC Ala	CCC Pro	CAT His	TAC Tyr	TAT Tyr 475	GAG Glu	CAG Gln	ACC Thr	CGA Arg	GGG Gly 480	1440
GGG Gly	CAG Gln	GTC Val	TCA Ser	GTG Val 485	TKC Xaa	CCC Pro	GTG Val	CCC Pro	GGT Gly 490	GTG Val	AGG Arg	GGC Gly	AGG Arg	TGG Trp 495	CAG Gln	1488
TGT Cys	GAG Glu	GCC Ala	ACC Thr 500	CTC Leu	CAC His	GGG Gly	GAG Glu	CAG Gln 505	GRC Xaa	CAT His	CCT Pro	TGG Trp	GGC Gly 510	CGC Arg	TTT Phe	1536
GGG Gly	GTG Val	GCT Ala 515	CTG Leu	ACA Thr	GTG Val	CTG Leu	GGG Gly 520	GAC Asp	GTA Val	AAC Asn	GGG Gly	GAC Asp 525	AAT Asn	CTG Leu	GCA Ala	1584
GAC Asp 530	GTG Val	GCT Ala	ATT Ile	GGT Gly	GCC Ala	CCT Pro 535	GGA Gly	GAG Glu	GAG Glu	GAG Glu	AGC Ser 540	AGA Arg	GGT Gly	GCT Ala	GTC Val	1632

- 110 -

TAC ATA TTT CAT GGA GCC TCG AGA CTG GAG ATC ATG CCC TCA CCC AGC Tyr Ile Phe His Gly Ala Ser Arg Leu Glu Ile Met Pro Ser Pro Ser 545 550 555 560	1680
CAG CGG GTC ACT GGC TCC CAG CTC TCC CTG AGA CTG CAG TAT TTT GGG Gln Arg Val Thr Gly Ser Gln Leu Ser Leu Arg Leu Gln Tyr Phe Gly 565 570 575	1728
CAG TCA TTG AGT GGG GGT CAG GAC CTT ACA CAG GAT GGC CTG GTG GAC Gln Ser Leu Ser Gly Gly Gln Asp Leu Thr Gln Asp Gly Leu Val Asp 580 585 590	1776
CTG GCC GTG GGA GCC CAG GGG CAC GTA CTG CTG CTC AGG AGT CTG CCT Leu Ala Val Gly Ala Gln Gly His Val Leu Leu Leu Arg Ser Leu Pro 595 600 605	1824
CTG CTG AAA GTC GAG CTC TCC ATA AGA TTC GCC CCC ATG GAG GTG GCA Leu Leu Lys Val Glu Leu Ser Ile Arg Phe Ala Pro Met Glu Val Ala 610 615 620	1872
AAG GCT GTG TAC CAG TGC TGG GAA AGG ACT CCC ACT GTC CTC GAA GCT Lys Ala Val Tyr Gln Cys Trp Glu Arg Thr Pro Thr Val Leu Glu Ala 625 630 635 640	1920
GGA GAG GCC ACT GTC TGT CTC ACT GTC CAC AAA GGC TCA CCT GAC CTG Gly Glu Ala Thr Val Cys Leu Thr Val His Lys Gly Ser Pro Asp Leu 645 650 655	1968
TTA GGT AAT GTC CAA GGC TCT GTC AGG TAT GAT CTG GCG TTA GAT CCG Leu Gly Asn Val Gln Gly Ser Val Arg Tyr Asp Leu Ala Leu Asp Pro 660 665 670	2016
GGC CGC CTG ATT TCT CGT GCC ATT TTT GAT GAG ACT AAG AAC TGC ACT Gly Arg Leu Ile Ser Arg Ala Ile Phe Asp Glu Thr Lys Asn Cys Thr 675 680 685	2064
TTG ACG GGA AGC AAG ACT CTG GGG CTT GGT GAT CAC TGC GAA ACA GTG Leu Thr Gly Arg Lys Thr Leu Gly Leu Gly Asp His Cys Glu Thr Val 690 695 700	2112
AAG CTG CTT TTG CCG GAC TGT GTG GAA GAT GCA GTG AGC CCT ATC ATC Lys Leu Leu Leu Pro Asp Cys Val Glu Asp Ala Val Ser Pro Ile Ile 705 710 715 720	2160
CTG CGC CTC AAC TTT TCC CTG GTG AGA GAC TCT GCT TCA CCC AGG AAC Leu Arg Leu Asn Phe Ser Leu Val Arg Asp Ser Ala Ser Pro Arg Asn 725 730 735	2208
CTG CAT CCT GTG CTG GCT GTG GGC TCA CAA GAC CAC ATA ACT GCT TCT Leu His Pro Val Leu Ala Val Gly Ser Gln Asp His Ile Thr Ala Ser 740 745 750	2256
CTG CCG TTT GAG AAG AAC TGT AAG CAA GAA CTC CTG TGT GAG GGG GAC Leu Pro Phe Glu Lys Asn Cys Lys Gln Glu Leu Leu Cys Glu Gly Asp 755 760 765	2304
CTG GGC ATC AGC TTT AAC TTC TCA GGC CTG CAG GTC TTG GTG GTG GGA Leu Gly Ile Ser Phe Asn Phe Ser Gly Leu Gln Val Leu Val Val Gly 770 775 780	2352

- 111 -

GGC	TCC	CCA	GAG	CTC	ACT	GTG	ACA	GTC	ACT	GTG	TGG	AAT	GAG	GGT	GAG	2400
Gly	Ser	Pro	Glu	Leu	Thr	Val	Thr	Val	Thr	Val	Trp	Asn	Glu	Gly	Glu	
785					790					795					800	
GAC	AGC	TAT	GGA	ACT	TTA	GTC	AAG	TTC	TAC	TAC	CCA	GCA	GGG	CTA	TCT	2448
Asp	Ser	Tyr	Gly	Thr	Leu	Val	Lys	Phe	Tyr	Tyr	Pro	Ala	Gly	Leu	Ser	
				805					810					815		
TAC	CGA	CGG	GTA	ACA	GGG	ACT	CAG	CAA	CCT	CAT	CAG	TAC	CCA	CTA	CGC	2496
Tyr	Arg	Arg	Val	Thr	Gly	Thr	Gln	Gln	Pro	His	Gln	Tyr	Pro	Leu	Arg	
			820					825					830			
TTG	GCC	TGT	GAG	GCT	GAG	CCC	GCT	GCC	CAG	GAG	GAC	CTG	AGG	AGC	AGC	2544
Leu	Ala	Cys	Glu	Ala	Glu	Pro	Ala	Ala	Gln	Glu	Asp	Leu	Arg	Ser	Ser	
		835					840					845				
AGC	TGT	AGC	ATT	AAT	CAC	CCC	ATC	TTC	CGA	GAA	GGT	GCA	AAG	ACC	ACC	2592
Ser	Cys	Ser	Ile	Asn	His	Pro	Ile	Phe	Arg	Glu	Gly	Ala	Lys	Thr	Thr	
	850					855					860					
TTC	ATG	ATC	ACA	TTC	GAT	GTC	TCC	TAC	AAG	GCC	TTC	CTA	GGA	GAC	AGG	2640
Phe	Met	Ile	Thr	Phe	Asp	Val	Ser	Tyr	Lys	Ala	Phe	Leu	Gly	Asp	Arg	
865					870					875					880	
TTG	CTT	CTG	AGG	GCC	AAA	GCC	AGC	AGT	GAG	AAT	AAT	AAG	CCT	GAT	ACC	2688
Leu	Leu	Leu	Arg	Ala	Lys	Ala	Ser	Ser	Glu	Asn	Asn	Lys	Pro	Asp	Thr	
				885					890					895		
AAC	AAG	ACT	GCC	TTC	CAG	CTG	GAG	CTC	CCA	GTG	AAG	TAC	ACC	GTC	TAT	2736
Asn	Lys	Thr	Ala	Phe	Gln	Leu	Glu	Leu	Pro	Val	Lys	Tyr	Thr	Val	Tyr	
			900					905					910			
ACC	CTG	ATC	AGT	AGG	CAA	GAA	GAT	TCC	ACC	AAC	CAT	GTC	AAC	TTT	TCA	2784
Thr	Leu	Ile	Ser	Arg	Gln	Glu	Asp	Ser	Thr	Asn	His	Val	Asn	Phe	Ser	
		915					920					925				
TCT	TCC	CAC	GGG	GGG	AGA	AGG	CAA	GAA	GCC	GCA	CAT	CGC	TAT	CGT	GTG	2832
Ser	Ser	His	Gly	Gly	Arg	Arg	Gln	Glu	Ala	Ala	His	Arg	Tyr	Arg	Val	
	930					935						940				
AAT	AAC	CTG	AGT	CCA	CTG	AAG	CTG	GCC	GTC	AGA	GTT	AAC	TTC	TGG	GTC	2880
Asn	Asn	Leu	Ser	Pro	Leu	Lys	Leu	Ala	Val	Arg	Val	Asn	Phe	Trp	Val	
945					950				955					960		
CCT	GTC	CTT	CTG	AAC	GGT	GTG	GCT	GTG	TGG	GAC	GTG	ACT	CTG	AGC	AGC	2928
Pro	Val	Leu	Leu	Asn	Gly	Val	Ala	Val	Trp	Asp	Val	Thr	Leu	Ser	Ser	
				965					970					975		
CCA	GCA	CAG	GGT	GTC	TCC	TGC	GTG	TCC	CAG	ATG	AAA	CCT	CCT	CAG	AAT	2976
Pro	Ala	Gln	Gly	Val	Ser	Cys	Val	Ser	Gln	Met	Lys	Pro	Pro	Gln	Asn	
			980					985					990			
CCC	GAC	TTT	CTG	ACC	CAG	ATT	CAG	AGA	CGT	TCT	GTG	CTG	GAC	TGC	TCC	3024
Pro	Asp	Phe	Leu	Thr	Gln	Ile	Gln	Arg	Arg	Ser	Val	Leu	Asp	Cys	Ser	
		995				1000						1005				
ATT	GCT	GAC	TGC	CTG	CAC	TCC	CGC	TGT	GAC	ATC	CCC	TCC	TTG	GAC	ATC	3072
Ile	Ala	Asp	Cys	Leu	His	Ser	Arg	Cys	Asp	Ile	Pro	Ser	Leu	Asp	Ile	
	1010					1015					1020					

- 112 -

CAG GAT GAA CTT GAC TTC ATT CTG AGG GGC AAC CTC AGC TTC GGC TGG Gln Asp Glu Leu Asp Phe Ile Leu Arg Gly Asn Leu Ser Phe Gly Trp 1025 1030 1035 1040	3120
GTC AGT CAG ACA TTG CAG GAA AAG GTG TTG CTT GTG AGT GAG GCT GAA Val Ser Gln Thr Leu Gln Glu Lys Val Leu Leu Val Ser Glu Ala Glu 1045 1050 1055	3168
ATC ACT TTC GAC ACA TCT GTG TAC TCC CAG CTG CCA GGA CAG GAG GCA Ile Thr Phe Asp Thr Ser Val Tyr Ser Gln Leu Pro Gly Gln Glu Ala 1060 1065 1070	3216
TTT CTG AGA GCC CAG GTG GAG ACA ACG TTA GAA GAA TAC GTG GTC TAT Phe Leu Arg Ala Gln Val Glu Thr Thr Leu Glu Glu Tyr Val Val Tyr 1075 1080 1085	3264
GAG CCC ATC TTC CTC GTG GCG GGC AGC TCG GTG GGA GGT CTG CTG TTA Glu Pro Ile Phe Leu Val Ala Gly Ser Ser Val Gly Gly Leu Leu Leu 1090 1095 1100	3312
CTG GCT CTC ATC ACA GTG GTA CTG TAC AAG CTT GGC TYC TYC AAA CGT Leu Ala Leu Ile Thr Val Val Leu Tyr Lys Leu Gly Xaa Xaa Lys Arg 1105 1110 1115 1120	3360
CAG TAC AAA GAA ATG CTG GAC GGC AAG GCT GCA GAT CCT GTC ACA GCC Gln Tyr Lys Glu Met Leu Asp Gly Lys Ala Ala Asp Pro Val Thr Ala 1125 1130 1135	3408
GGC CAG GCA GAT TTC GGC TGT GAG ACT CCT CCA TAT CTC GTG AGC TAGGAATCCA Gly Gln Ala Asp Phe Gly Cys Glu Thr Pro Pro Tyr Leu Val Ser 1140 1145 1150	3463
CTCTCCTGCC TATCTCTGNA ATGAAGATTG GTCCTGCCTA TGAGTCTACT GGCATGGGAA	3523
CGAGT	3528

## (2) INFORMATION FOR SEQ ID NO:37:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1151 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:37:

Gly Trp Ala Leu Ala Ser Cys His Gly Ser Asn Leu Asp Val Glu Glu 1 5 10 15
Pro Ile Val Phe Arg Glu Asp Ala Ala Ser Phe Gly Gln Thr Val Val 20 25 30
Gln Phe Gly Gly Ser Arg Leu Val Val Gly Ala Pro Leu Glu Ala Val 35 40 45
Ala Val Asn Gln Thr Gly Arg Leu Tyr Asp Cys Ala Pro Ala Thr Gly 50 55 60

- 113 -

Met Cys Gln Pro Ile Val Leu Arg Ser Pro Leu Glu Ala Val Asn Met  
 65 70 75 80  
 Ser Leu Gly Leu Ser Leu Val Thr Ala Thr Asn Asn Ala Gln Leu Leu  
 85 90 95  
 Ala Cys Gly Pro Thr Ala Gln Arg Ala Cys Val Lys Asn Met Tyr Ala  
 100 105 110  
 Lys Gly Ser Cys Leu Leu Leu Gly Ser Ser Leu Gln Phe Ile Gln Ala  
 115 120 125  
 Val Pro Ala Ser Met Pro Glu Cys Pro Arg Gln Glu Met Asp Ile Ala  
 130 135 140  
 Phe Leu Ile Asp Gly Ser Gly Ser Ile Asn Gln Arg Asp Phe Ala Gln  
 145 150 155 160  
 Met Lys Asp Phe Val Lys Ala Leu Met Gly Glu Phe Ala Ser Thr Ser  
 165 170 175  
 Thr Leu Phe Ser Leu Met Gln Tyr Ser Asn Ile Leu Lys Thr His Phe  
 180 185 190  
 Thr Phe Thr Glu Phe Lys Asn Ile Leu Asp Pro Gln Ser Leu Val Asp  
 195 200 205  
 Pro Ile Val Gln Leu Gln Gly Leu Thr Tyr Thr Ala Thr Gly Ile Arg  
 210 215 220  
 Thr Val Met Glu Glu Leu Phe His Ser Lys Asn Gly Ser Arg Lys Ser  
 225 230 235 240  
 Ala Lys Lys Ile Leu Leu Val Ile Thr Asp Gly Gln Lys Tyr Arg Asp  
 245 250 255  
 Pro Leu Glu Tyr Ser Asp Val Ile Pro Ala Ala Asp Lys Ala Gly Ile  
 260 265 270  
 Ile Arg Tyr Ala Ile Gly Val Gly Asp Ala Phe Gln Glu Pro Thr Ala  
 275 280 285  
 Leu Lys Glu Leu Asn Thr Ile Gly Ser Ala Pro Pro Gln Asp His Val  
 290 295 300  
 Phe Lys Val Gly Asn Phe Ala Ala Leu Arg Ser Ile Gln Arg Gln Leu  
 305 310 315 320  
 Gln Glu Lys Ile Phe Ala Ile Glu Gly Thr Gln Ser Arg Ser Ser Ser  
 325 330 335  
 Ser Phe Gln His Glu Met Ser Gln Glu Gly Phe Ser Ser Ala Leu Thr  
 340 345 350  
 Ser Asp Gly Pro Val Leu Gly Ala Xaa Gly Ser Phe Ser Trp Ser Gly  
 355 360 365  
 Gly Ala Phe Leu Tyr Pro Pro Asn Thr Arg Pro Thr Phe Ile Asn Met  
 370 375 380

- 114 -

Ser Gln Glu Asn Val Asp Met Arg Asp Ser Tyr Leu Gly Tyr Ser Thr  
 385 390 395 400  
 Ala Val Ala Phe Trp Lys Gly Val His Ser Leu Ile Leu Gly Ala Pro  
 405 410 415  
 Arg His Gln His Thr Gly Lys Val Val Ile Phe Thr Gln Glu Ala Arg  
 420 425 430  
 His Trp Arg Pro Lys Ser Glu Val Arg Gly Thr Gln Ile Gly Ser Tyr  
 435 440 445  
 Phe Gly Ala Ser Leu Cys Ser Val Asp Val Asp Arg Asp Gly Ser Xaa  
 450 455 460  
 Asp Leu Val Leu Ile Gly Ala Pro His Tyr Tyr Glu Gln Thr Arg Gly  
 465 470 475 480  
 Gly Gln Val Ser Val Xaa Pro Val Pro Gly Val Arg Gly Arg Trp Gln  
 485 490 495  
 Cys Glu Ala Thr Leu His Gly Glu Gln Xaa His Pro Trp Gly Arg Phe  
 500 505 510  
 Gly Val Ala Leu Thr Val Leu Gly Asp Val Asn Gly Asp Asn Leu Ala  
 515 520 525  
 Asp Val Ala Ile Gly Ala Pro Gly Glu Glu Glu Ser Arg Gly Ala Val  
 530 535 540  
 Tyr Ile Phe His Gly Ala Ser Arg Leu Glu Ile Met Pro Ser Pro Ser  
 545 550 555 560  
 Gln Arg Val Thr Gly Ser Gln Leu Ser Leu Arg Leu Gln Tyr Phe Gly  
 565 570 575  
 Gln Ser Leu Ser Gly Gly Gln Asp Leu Thr Gln Asp Gly Leu Val Asp  
 580 585 590  
 Leu Ala Val Gly Ala Gln Gly His Val Leu Leu Leu Arg Ser Leu Pro  
 595 600 605  
 Leu Leu Lys Val Glu Leu Ser Ile Arg Phe Ala Pro Met Glu Val Ala  
 610 615 620  
 Lys Ala Val Tyr Gln Cys Trp Glu Arg Thr Pro Thr Val Leu Glu Ala  
 625 630 635 640  
 Gly Glu Ala Thr Val Cys Leu Thr Val His Lys Gly Ser Pro Asp Leu  
 645 650 655  
 Leu Gly Asn Val Gln Gly Ser Val Arg Tyr Asp Leu Ala Leu Asp Pro  
 660 665 670  
 Gly Arg Leu Ile Ser Arg Ala Ile Phe Asp Glu Thr Lys Asn Cys Thr  
 675 680 685  
 Leu Thr Gly Arg Lys Thr Leu Gly Leu Gly Asp His Cys Glu Thr Val  
 690 695 700

- 115 -

Lys Leu Leu Leu Pro Asp Cys Val Glu Asp Ala Val Ser Pro Ile Ile  
 705 710 715 720  
 Leu Arg Leu Asn Phe Ser Leu Val Arg Asp Ser Ala Ser Pro Arg Asn  
 725 730 735  
 Leu His Pro Val Leu Ala Val Gly Ser Gln Asp His Ile Thr Ala Ser  
 740 745 750  
 Leu Pro Phe Glu Lys Asn Cys Lys Gln Glu Leu Leu Cys Glu Gly Asp  
 755 760 765  
 Leu Gly Ile Ser Phe Asn Phe Ser Gly Leu Gln Val Leu Val Val Gly  
 770 775 780  
 Gly Ser Pro Glu Leu Thr Val Thr Val Thr Val Trp Asn Glu Gly Glu  
 785 790 795 800  
 Asp Ser Tyr Gly Thr Leu Val Lys Phe Tyr Tyr Pro Ala Gly Leu Ser  
 805 810 815  
 Tyr Arg Arg Val Thr Gly Thr Gln Gln Pro His Gln Tyr Pro Leu Arg  
 820 825 830  
 Leu Ala Cys Glu Ala Glu Pro Ala Ala Gln Glu Asp Leu Arg Ser Ser  
 835 840 845  
 Ser Cys Ser Ile Asn His Pro Ile Phe Arg Glu Gly Ala Lys Thr Thr  
 850 855 860  
 Phe Met Ile Thr Phe Asp Val Ser Tyr Lys Ala Phe Leu Gly Asp Arg  
 865 870 875 880  
 Leu Leu Leu Arg Ala Lys Ala Ser Ser Glu Asn Asn Lys Pro Asp Thr  
 885 890 895  
 Asn Lys Thr Ala Phe Gln Leu Glu Leu Pro Val Lys Tyr Thr Val Tyr  
 900 905 910  
 Thr Leu Ile Ser Arg Gln Glu Asp Ser Thr Asn His Val Asn Phe Ser  
 915 920 925  
 Ser Ser His Gly Gly Arg Arg Gln Glu Ala Ala His Arg Tyr Arg Val  
 930 935 940  
 Asn Asn Leu Ser Pro Leu Lys Leu Ala Val Arg Val Asn Phe Trp Val  
 945 950 955 960  
 Pro Val Leu Leu Asn Gly Val Ala Val Trp Asp Val Thr Leu Ser Ser  
 965 970 975  
 Pro Ala Gln Gly Val Ser Cys Val Ser Gln Met Lys Pro Pro Gln Asn  
 980 985 990  
 Pro Asp Phe Leu Thr Gln Ile Gln Arg Arg Ser Val Leu Asp Cys Ser  
 995 1000 1005  
 Ile Ala Asp Cys Leu His Ser Arg Cys Asp Ile Pro Ser Leu Asp Ile  
 1010 1015 1020

- 116 -

Gln Asp Glu Leu Asp Phe Ile Leu Arg Gly Asn Leu Ser Phe Gly Trp  
 1025 1030 1035 1040  
 Val Ser Gln Thr Leu Gln Glu Lys Val Leu Leu Val Ser Glu Ala Glu  
 1045 1050 1055  
 Ile Thr Phe Asp Thr Ser Val Tyr Ser Gln Leu Pro Gly Gln Glu Ala  
 1060 1065 1070  
 Phe Leu Arg Ala Gln Val Glu Thr Thr Leu Glu Glu Tyr Val Val Tyr  
 1075 1080 1085  
 Glu Pro Ile Phe Leu Val Ala Gly Ser Ser Val Gly Gly Leu Leu Leu  
 1090 1095 1100  
 Leu Ala Leu Ile Thr Val Val Leu Tyr Lys Leu Gly Xaa Xaa Lys Arg  
 1105 1110 1115 1120  
 Gln Tyr Lys Glu Met Leu Asp Gly Lys Ala Ala Asp Pro Val Thr Ala  
 1125 1130 1135  
 Gly Gln Ala Asp Phe Gly Cys Glu Thr Pro Pro Tyr Leu Val Ser  
 1140 1145 1150

## (2) INFORMATION FOR SEQ ID NO:38:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 21 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:38:

GTCCAAGCTG TCATGGGCCA G

21

## (2) INFORMATION FOR SEQ ID NO:39:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 23 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:39:

GTCCAGCAGA CTGAAGAGCA CGG

23

## (2) INFORMATION FOR SEQ ID NO:40:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 18 base pairs  
 (B) TYPE: nucleic acid



- 117 -

- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:40:

TGTAAAACGA CGGCCAGT

18

(2) INFORMATION FOR SEQ ID NO:41:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 19 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:41:

GGAAACAGCT ATGACCATG

19

(2) INFORMATION FOR SEQ ID NO:42:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 22 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:42:

GGACATGTTC ACTGCCTCTA GG

22

(2) INFORMATION FOR SEQ ID NO:43:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 25 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:43:

GGCGGACAGT CAGACGACTG TCCTG

25

(2) INFORMATION FOR SEQ ID NO:44:

- 118 -

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 38 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:44:

CTGGTTCGGC CCACCTCTGA AGGTTCCAGA ATCGATAG

38

(2) INFORMATION FOR SEQ ID NO:45:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 3519 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(ix) FEATURE:

- (A) NAME/KEY: CDS  
 (B) LOCATION: 52..3519

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:45:

GCTTTCTGAA GGTTCAGAA TCGATAGTGA ATTCGTGGGC ACTGCTCAGA T ATG GTC	57
Met Val 1	
CGT GGA GTT GTG ATC CTC CTG TGT GGC TGG GCC CTG GCT TCC TGT CAT	105
Arg Gly Val Val Ile Leu Leu Cys Gly Trp Ala Leu Ala Ser Cys His 5 10 15	
GGG TCT AAC CTG GAT GTG GAG AAG CCC GTC GTG TTC AAA GAG GAT GCA	153
Gly Ser Asn Leu Asp Val Glu Lys Pro Val Val Phe Lys Glu Asp Ala 20 25 30	
GCC AGC TTC GGA CAG ACT GTG GTG CAG TTT GGT GGA TCT CGA CTC GTG	201
Ala Ser Phe Gly Gln Thr Val Val Gln Phe Gly Gly Ser Arg Leu Val 35 40 45 50	
GTG GGA GCC CCT CTG GAG GCG GTG GCA GTC AAC CAA ACA GGA CAG TCG	249
Val Gly Ala Pro Leu Glu Ala Val Ala Val Asn Gln Thr Gly Gln Ser 55 60 65	
TCT GAC TGT CCG CCT GCC ACT GGC GTG TGC CAG CCC ATC TTA CTG CAC	297
Ser Asp Cys Pro Pro Ala Thr Gly Val Cys Gln Pro Ile Leu Leu His 70 75 80	
ATT CCC CTA GAG GCA GTG AAC ATG TCC CTG GGC CTG TCT CTG GTG GCT	345
Ile Pro Leu Glu Ala Val Asn Met Ser Leu Gly Leu Ser Leu Val Ala 85 90 95	

- 119 -

GAC ACC AAT AAC TCC CAG TTG CTG GCT TGT GGT CCA ACT GCA CAG AGA Asp Thr Asn Asn Ser Gln Leu Ala Cys Gly Pro Thr Ala Gln Arg 100 105 110	393
GCT TGT GCA AAG AAC ATG TAT GCA AAA GGT TCC TGC CTC CTT CTG GGC Ala Cys Ala Lys Asn Met Tyr Ala Lys Gly Ser Cys Leu Leu Leu Gly 115 120 125 130	441
TCC AGC TTG CAG TTC ATC CAG GCA ATC CCT GCT ACC ATG CCA GAG TGT Ser Ser Leu Gln Phe Ile Gln Ala Ile Pro Ala Thr Met Pro Glu Cys 135 140 145	489
CCA GGA CAA GAG ATG GAC ATT GCT TTC CTG ATT GAT GGC TCC GGC AGC Pro Gly Gln Glu Met Asp Ile Ala Phe Leu Ile Asp Gly Ser Gly Ser 150 155 160	537
ATT GAT CAA AGT GAC TTT ACC CAG ATG AAG GAC TTC GTC AAA GCT TTG Ile Asp Gln Ser Asp Phe Thr Gln Met Lys Asp Phe Val Lys Ala Leu 165 170 175	585
ATG GGC CAG TTG GCG AGC ACC AGC ACC TCG TTC TCC CTG ATG CAA TAC Met Gly Gln Leu Ala Ser Thr Ser Thr Ser Phe Ser Leu Met Gln Tyr 180 185 190	633
TCA AAC ATC CTG AAG ACT CAT TTT ACC TTC ACG GAA TTC AAG AGC AGC Ser Asn Ile Leu Lys Thr His Phe Thr Phe Thr Glu Phe Lys Ser Ser 195 200 205 210	681
CTG AGC CCT CAG AGC CTG GTG GAT GCC ATC GTC CAG CTC CAA GGC CTG Leu Ser Pro Gln Ser Leu Val Asp Ala Ile Val Gln Leu Gln Gly Leu 215 220 225	729
ACG TAC ACA GCC TCG GGC ATC CAG AAA GTG GTG AAA GAG CTA TTT CAT Thr Tyr Thr Ala Ser Gly Ile Gln Lys Val Val Lys Glu Leu Phe His 230 235 240	777
AGC AAG AAT GGG GCC CGA AAA AGT GCC AAG AAG ATA CTA ATT GTC ATC Ser Lys Asn Gly Ala Arg Lys Ser Ala Lys Lys Ile Leu Ile Val Ile 245 250 255	825
ACA GAT GGG CAG AAA TTC AGA GAC CCC CTG GAG TAT AGA CAT GTC ATC Thr Asp Gly Gln Lys Phe Arg Asp Pro Leu Glu Tyr Arg His Val Ile 260 265 270	873
CCT GAA GCA GAG AAA GCT GGG ATC ATT CGC TAT GCT ATA GGG GTG GGA Pro Glu Ala Glu Lys Ala Gly Ile Ile Arg Tyr Ala Ile Gly Val Gly 275 280 285 290	921
GAT GCC TTC CGG GAA CCC ACT GCC CTA CAG GAG CTG AAC ACC ATT GGC Asp Ala Phe Arg Glu Pro Thr Ala Leu Gln Glu Leu Asn Thr Ile Gly 295 300 305	969
TCA GCT CCC TCG CAG GAC CAC GTG TTC AAG GTG GGC AAT TTT GTA GCA Ser Ala Pro Ser Gln Asp His Val Phe Lys Val Gly Asn Phe Val Ala 310 315 320	1017
CTT CGC AGC ATC CAG CGG CAA ATT CAG GAG AAA ATC TTT GCC ATT GAA Leu Arg Ser Ile Gln Arg Gln Ile Gln Glu Lys Ile Phe Ala Ile Glu 325 330 335	1065

- 120 -

GGA ACC GAA TCA AGG TCA AGT AGT TCC TTT CAG CAC GAG ATG TCA CAA Gly Thr Glu Ser Arg Ser Ser Ser Ser Phe Gln His Glu Met Ser Gln 340 345 350	1113
GAA GGT TTC AGC TCA GCT CTC TCA ATG GAT GGA CCA GTT CTG GGG GCT Glu Gly Phe Ser Ser Ala Leu Ser Met Asp Gly Pro Val Leu Gly Ala 355 360 365 370	1161
GTG GGA GGC TTC AGC TGG TCT GGA GGT GCC TTC TTG TAC CCC TCA AAT Val Gly Gly Phe Ser Trp Ser Gly Gly Ala Phe Leu Tyr Pro Ser Asn 375 380 385	1209
ATG AGA TCC ACC TTC ATC AAC ATG TCT CAG GAG AAC GAG GAT ATG AGG Met Arg Ser Thr Phe Ile Asn Met Ser Gln Glu Asn Glu Asp Met Arg 390 395 400	1257
GAC GCT TAC CTG GGT TAC TCC ACC GCA CTG GCC TTT TGG AAG GGG GTC Asp Ala Tyr Leu Gly Tyr Ser Thr Ala Leu Ala Phe Trp Lys Gly Val 405 410 415	1305
CAC AGC CTG ATC CTG GGG GCC CCT CGC CAC CAG CAC ACG GGG AAG GTT His Ser Leu Ile Leu Gly Ala Pro Arg His Gln His Thr Gly Lys Val 420 425 430	1353
GTC ATC TTT ACC CAG GAA TCC AGG CAC TGG AGG CCC AAG TCT GAA GTC Val Ile Phe Thr Gln Glu Ser Arg His Trp Arg Pro Lys Ser Glu Val 435 440 445 450	1401
AGA GGG ACA CAG ATC GGC TCC TAC TTT GGG GCA TCT CTC TGT TCT GTG Arg Gly Thr Gln Ile Gly Ser Tyr Phe Gly Ala Ser Leu Cys Ser Val 455 460 465	1449
GAC ATG GAT AGA GAT GGC AGC ACT GAC CTG GTC CTG ATT GGA GTC CCC Asp Met Asp Arg Asp Gly Ser Thr Asp Leu Val Leu Ile Gly Val Pro 470 475 480	1497
CAT TAC TAT GAG CAC ACC CGA GGG GGG CAG GTG TCG GTG TGC CCC ATG His Tyr Tyr Glu His Thr Arg Gly Gly Gln Val Ser Val Cys Pro Met 485 490 495	1545
CCT GGT GTG AGG AGC AGG TGG CAT TGT GGG ACC ACC CTC CAT GGG GAG Pro Gly Val Arg Ser Arg Trp His Cys Gly Thr Thr Leu His Gly Glu 500 505 510	1593
CAG GGC CAT CCT TGG GGC CGC TTT GGG GCG GCT CTG ACA GTG CTA GGG Gln Gly His Pro Trp Gly Arg Phe Gly Ala Ala Leu Thr Val Leu Gly 515 520 525 530	1641
GAC GTG AAT GGG GAC AGT CTG GCG GAT GTG GCT ATT GGT GCA CCC GGA Asp Val Asn Gly Asp Ser Leu Ala Asp Val Ala Ile Gly Ala Pro Gly 535 540 545	1689
GAG GAG GAG AAC AGA GGT GCT GTC TAC ATA TTT CAT GGA GCC TCG AGA Glu Glu Glu Asn Arg Gly Ala Val Tyr Ile Phe His Gly Ala Ser Arg 550 555 560	1737
CAG GAC ATC GCT CCC TCG CCT AGC CAG CGG GTC ACT GGC TCC CAG CTC Gln Asp Ile Ala Pro Ser Pro Ser Gln Arg Val Thr Gly Ser Gln Leu 565 570 575	1785

- 121 -

TTC	CTG	AGG	CTC	CAA	TAT	TTT	GGG	CAG	TCA	TTA	AGT	GGG	GGT	CAG	GAC	1833
Phe	Leu	Arg	Leu	Gln	Tyr	Phe	Gly	Gln	Ser	Leu	Ser	Gly	Gly	Gln	Asp	
580						585					590					
CTT	ACA	CAG	GAT	GGC	CTG	GTG	GAC	CTG	GCC	GTG	GGA	GCC	CAG	GGG	CAC	1881
Leu	Thr	Gln	Asp	Gly	Leu	Val	Asp	Leu	Ala	Val	Gly	Ala	Gln	Gly	His	
595					600					605					610	
GTG	CTG	CTG	CTT	AGG	AGT	CTG	CCT	TTG	CTG	AAA	GTG	GGG	ATC	TCC	ATT	1929
Val	Leu	Leu	Leu	Arg	Ser	Leu	Pro	Leu	Leu	Lys	Val	Gly	Ile	Ser	Ile	
				615					620					625		
AGA	TTT	GCC	CCC	TCA	GAG	GTG	GCA	AAG	ACT	GTG	TAC	CAG	TGC	TGG	GGA	1977
Arg	Phe	Ala	Pro	Ser	Glu	Val	Ala	Lys	Thr	Val	Tyr	Gln	Cys	Trp	Gly	
			630					635					640			
AGG	ACT	CCC	ACT	GTC	CTC	GAA	GCT	GGA	GAG	GCC	ACC	GTC	TGT	CTC	ACT	2025
Arg	Thr	Pro	Thr	Val	Leu	Glu	Ala	Gly	Glu	Ala	Thr	Val	Cys	Leu	Thr	
		645					650					655				
GTC	CGC	AAA	GGT	TCA	CCT	GAC	CTG	TTA	GGT	GAT	GTC	CAA	AGC	TCT	GTC	2073
Val	Arg	Lys	Gly	Ser	Pro	Asp	Leu	Leu	Gly	Asp	Val	Gln	Ser	Ser	Val	
660						665					670					
AGG	TAT	GAT	CTG	GGC	TTG	GAT	CCG	GGC	CGT	CTG	ATT	TCT	CGT	GCC	ATT	2121
Arg	Tyr	Asp	Leu	Ala	Leu	Asp	Pro	Gly	Arg	Leu	Ile	Ser	Arg	Ala	Ile	
675					680					685					690	
TTT	GAT	GAG	ACG	AAG	AAC	TGC	ACT	TTG	ACC	CGA	AGG	AAG	ACT	CTG	GGG	2169
Phe	Asp	Glu	Thr	Lys	Asn	Cys	Thr	Leu	Thr	Arg	Arg	Lys	Thr	Leu	Gly	
				695					700					705		
CTT	GGT	GAT	CAC	TGC	GAA	ACA	ATG	AAG	CTG	CTT	TTG	CCA	GAC	TGT	GTG	2217
Leu	Gly	Asp	His	Cys	Glu	Thr	Met	Lys	Leu	Leu	Leu	Pro	Asp	Cys	Val	
			710					715					720			
GAG	GAT	GCA	GTG	ACC	CCT	ATC	ATC	CTG	CGC	CTT	AAC	TTA	TCC	CTG	GCA	2265
Glu	Asp	Ala	Val	Thr	Pro	Ile	Ile	Leu	Arg	Leu	Asn	Leu	Ser	Leu	Ala	
		725					730					735				
GGG	GAC	TCT	GCT	CCA	TCC	AGG	AAC	CTT	CGT	CCT	GTG	CTG	GCT	GTG	GGC	2313
Gly	Asp	Ser	Ala	Pro	Ser	Arg	Asn	Leu	Arg	Pro	Val	Leu	Ala	Val	Gly	
740						745					750					
TCA	CAA	GAC	CAT	GTA	ACA	GCT	TCT	TTC	CCG	TTT	GAG	AAG	AAC	TGT	GAG	2361
Ser	Gln	Asp	His	Val	Thr	Ala	Ser	Phe	Pro	Phe	Glu	Lys	Asn	Cys	Glu	
755					760					765					770	
GGG	AAC	CTG	GGC	GTC	AGC	TTC	AAC	TTC	TCA	GGC	CTG	CAG	GTC	TTG	GAG	2409
Gly	Asn	Leu	Gly	Val	Ser	Phe	Asn	Phe	Ser	Gly	Leu	Gln	Val	Leu	Glu	
			775						780					785		
GTA	GGA	AGC	TCC	CCA	GAG	CTC	ACT	GTG	ACA	GTA	ACA	GTT	TGG	AAT	GAG	2457
Val	Gly	Ser	Ser	Pro	Glu	Leu	Thr	Val	Thr	Val	Thr	Val	Trp	Asn	Glu	
			790					795					800			
GGT	GAG	GAC	AGC	TAT	GGA	ACC	TTA	ATC	AAG	TTC	TAC	TAC	CCA	GCA	GAG	2505
Gly	Glu	Asp	Ser	Tyr	Gly	Thr	Leu	Ile	Lys	Phe	Tyr	Tyr	Pro	Ala	Glu	
		805					810					815				

- 122 -

CTA TCT TAC CGA CGG GTG ACA AGA GCC CAG CAA CCT CAT CCG TAC CCA Leu Ser Tyr Arg Arg Val Thr Arg Ala Gln Gln Pro His Pro Tyr Pro 820 825 830	2553
CTA CGC CTG GCA TGT GAG GCT GAG CCC ACG GGC CAG GAG AGC CTG AGG Leu Arg Leu Ala Cys Glu Ala Glu Pro Thr Gly Gln Glu Ser Leu Arg 835 840 845 850	2601
AGC AGC AGC TGT AGC ATC AAT CAC CCC ATC TTC CGA GAA GGT GCC AAG Ser Ser Ser Cys Ser Ile Asn His Pro Ile Phe Arg Glu Gly Ala Lys 855 860 865	2649
GCC ACC TTC ATG ATC ACA TTT GAT GTC TCC TAC AAG GCC TTC CTG GGA Ala Thr Phe Met Ile Thr Phe Asp Val Ser Tyr Lys Ala Phe Leu Gly 870 875 880	2697
GAC AGG TTG CTT CTG AGG GCC AGC GCA AGC AGT GAG AAT AAT AAG CCT Asp Arg Leu Leu Leu Arg Ala Ser Ala Ser Ser Glu Asn Asn Lys Pro 885 890 895	2745
GAA ACC AGC AAG ACT GCC TTC CAG CTG GAG CTT CCG GTG AAG TAC ACG Glu Thr Ser Lys Thr Ala Phe Gln Leu Glu Leu Pro Val Lys Tyr Thr 900 905 910	2793
GTC TAT ACC GTG ATC AGT AGG CAG GAA GAT TCT ACC AAG CAT TTC AAC Val Tyr Thr Val Ile Ser Arg Gln Glu Asp Ser Thr Lys His Phe Asn 915 920 925 930	2841
TTC TCA TCT TCC CAC GGG GAG AGA CAG AAA GAG GCC GAA CAT CGA TAT Phe Ser Ser Ser His Gly Glu Arg Gln Lys Glu Ala Glu His Arg Tyr 935 940 945	2889
CGT GTG AAT AAC CTG AGT CCA TTG ACG CTG GCC ATC AGC GTT AAC TTC Arg Val Asn Asn Leu Ser Pro Leu Thr Leu Ala Ile Ser Val Asn Phe 950 955 960	2937
TGG GTC CCC ATC CTT CTG AAT GGT GTG GCC GTG TGG GAT GTG ACT CTG Trp Val Pro Ile Leu Leu Asn Gly Val Ala Val Trp Asp Val Thr Leu 965 970 975	2985
AGG AGC CCA GCA CAG GGT GTC TCC TGT GTG TCA CAG AGG GAA CCT CCT Arg Ser Pro Ala Gln Gly Val Ser Cys Val Ser Gln Arg Glu Pro Pro 980 985 990	3033
CAA CAT TCC GAC CTT CTG ACC CAG ATC CAA GGA CGC TCT GTG CTG GAC Gln His Ser Asp Leu Leu Thr Gln Ile Gln Gly Arg Ser Val Leu Asp 995 1000 1005 1010	3081
TGC GCC ATC GCC GAC TGC CTG CAC CTC CGC TGT GAC ATC CCC TCC TTG Cys Ala Ile Ala Asp Cys Leu His Leu Arg Cys Asp Ile Pro Ser Leu 1015 1020 1025	3129
GGC ACC CTG GAT GAG CTT GAC TTC ATT CTG AAG GGC AAC CTC AGC TTC Gly Thr Leu Asp Glu Leu Asp Phe Ile Leu Lys Gly Asn Leu Ser Phe 1030 1035 1040	3177
GGC TGG ATC AGT CAG ACA TTG CAG AAA AAG GTG TTG CTC CTG AGT GAG Gly Trp Ile Ser Gln Thr Leu Gln Lys Lys Val Leu Leu Leu Ser Glu 1045 1050 1055	3225

- 123 -

GCT GAA ATC ACA TTC AAC ACA TCT GTG TAT TCC CAG CTG CCG GGA CAG Ala Glu Ile Thr Phe Asn Thr Ser Val Tyr Ser Gln Leu Pro Gly Gln 1060 1065 1070	3273
GAG GCA TTT CTG AGA GCC CAG GTG TCA ACG ATG CTA GAA GAA TAC GTG Glu Ala Phe Leu Arg Ala Gln Val Ser Thr Met Leu Glu Glu Tyr Val 1075 1080 1085 1090	3321
GTC TAT GAG CCC GTC TTC CTC ATG GTG TTC AGC TCA GTG GGA GGT CTG Val Tyr Glu Pro Val Phe Leu Met Val Phe Ser Ser Val Gly Gly Leu 1095 1100 1105	3369
CTG TTA CTG GCT CTC ATC ACT GTG GCG CTG TAC AAG CTT GGC TTC TTC Leu Leu Leu Ala Leu Ile Thr Val Ala Leu Tyr Lys Leu Gly Phe Phe 1110 1115 1120	3417
AAA CGT CAG TAT AAA GAG ATG CTG GAT CTA CCA TCT GCA GAT CCT GAC Lys Arg Gln Tyr Lys Glu Met Leu Asp Leu Pro Ser Ala Asp Pro Asp 1125 1130 1135	3465
CCA GCC GGC CAG GCA GAT TCC AAC CAT GAG ACT CCT CCA CAT CTC ACG Pro Ala Gly Gln Ala Asp Ser Asn His Glu Thr Pro Pro His Leu Thr 1140 1145 1150	3513
TCC TAG Ser 1155	3519

## (2) INFORMATION FOR SEQ ID NO:46:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 1155 amino acids
  - (B) TYPE: amino acid
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:46:

Met Val Arg Gly Val Val Ile Leu Leu Cys Gly Trp Ala Leu Ala Ser 1 5 10 15
Cys His Gly Ser Asn Leu Asp Val Glu Lys Pro Val Val Phe Lys Glu 20 25 30
Asp Ala Ala Ser Phe Gly Gln Thr Val Val Gln Phe Gly Gly Ser Arg 35 40 45
Leu Val Val Gly Ala Pro Leu Glu Ala Val Ala Val Asn Gln Thr Gly 50 55 60
Gln Ser Ser Asp Cys Pro Pro Ala Thr Gly Val Cys Gln Pro Ile Leu 65 70 75 80
Leu His Ile Pro Leu Glu Ala Val Asn Met Ser Leu Gly Leu Ser Leu 85 90 95
Val Ala Asp Thr Asn Asn Ser Gln Leu Leu Ala Cys Gly Pro Thr Ala 100 105 110

- 124 -

Gln Arg Ala Cys Ala Lys Asn Met Tyr Ala Lys Gly Ser Cys Leu Leu  
 115 120 125  
 Leu Gly Ser Ser Leu Gln Phe Ile Gln Ala Ile Pro Ala Thr Met Pro  
 130 135 140  
 Glu Cys Pro Gly Gln Glu Met Asp Ile Ala Phe Leu Ile Asp Gly Ser  
 145 150 155 160  
 Gly Ser Ile Asp Gln Ser Asp Phe Thr Gln Met Lys Asp Phe Val Lys  
 165 170 175  
 Ala Leu Met Gly Gln Leu Ala Ser Thr Ser Thr Ser Phe Ser Leu Met  
 180 185 190  
 Gln Tyr Ser Asn Ile Leu Lys Thr His Phe Thr Phe Thr Glu Phe Lys  
 195 200 205  
 Ser Ser Leu Ser Pro Gln Ser Leu Val Asp Ala Ile Val Gln Leu Gln  
 210 215 220  
 Gly Leu Thr Tyr Thr Ala Ser Gly Ile Gln Lys Val Val Lys Glu Leu  
 225 230 235 240  
 Phe His Ser Lys Asn Gly Ala Arg Lys Ser Ala Lys Lys Ile Leu Ile  
 245 250 255  
 Val Ile Thr Asp Gly Gln Lys Phe Arg Asp Pro Leu Glu Tyr Arg His  
 260 265 270  
 Val Ile Pro Glu Ala Glu Lys Ala Gly Ile Ile Arg Tyr Ala Ile Gly  
 275 280 285  
 Val Gly Asp Ala Phe Arg Glu Pro Thr Ala Leu Gln Glu Leu Asn Thr  
 290 295 300  
 Ile Gly Ser Ala Pro Ser Gln Asp His Val Phe Lys Val Gly Asn Phe  
 305 310 315 320  
 Val Ala Leu Arg Ser Ile Gln Arg Gln Ile Gln Glu Lys Ile Phe Ala  
 325 330 335  
 Ile Glu Gly Thr Glu Ser Arg Ser Ser Ser Phe Gln His Glu Met  
 340 345 350  
 Ser Gln Glu Gly Phe Ser Ser Ala Leu Ser Met Asp Gly Pro Val Leu  
 355 360 365  
 Gly Ala Val Gly Gly Phe Ser Trp Ser Gly Gly Ala Phe Leu Tyr Pro  
 370 375 380  
 Ser Asn Met Arg Ser Thr Phe Ile Asn Met Ser Gln Glu Asn Glu Asp  
 385 390 395 400  
 Met Arg Asp Ala Tyr Leu Gly Tyr Ser Thr Ala Leu Ala Phe Trp Lys  
 405 410 415  
 Gly Val His Ser Leu Ile Leu Gly Ala Pro Arg His Gln His Thr Gly  
 420 425 430



- 125 -

Lys Val Val Ile Phe Thr Gln Glu Ser Arg His Trp Arg Pro Lys Ser  
                   435                                  440                                  445  
 Glu Val Arg Gly Thr Gln Ile Gly Ser Tyr Phe Gly Ala Ser Leu Cys  
                   450                                  455                                  460  
 Ser Val Asp Met Asp Arg Asp Gly Ser Thr Asp Leu Val Leu Ile Gly  
                   465                                  470                                  475                                  480  
 Val Pro His Tyr Tyr Glu His Thr Arg Gly Gly Gln Val Ser Val Cys  
                                   485                                  490                                  495  
 Pro Met Pro Gly Val Arg Ser Arg Trp His Cys Gly Thr Thr Leu His  
                                   500                                  505                                  510  
 Gly Glu Gln Gly His Pro Trp Gly Arg Phe Gly Ala Ala Leu Thr Val  
                                   515                                  520                                  525  
 Leu Gly Asp Val Asn Gly Asp Ser Leu Ala Asp Val Ala Ile Gly Ala  
                   530                                  535                                  540  
 Pro Gly Glu Glu Glu Asn Arg Gly Ala Val Tyr Ile Phe His Gly Ala  
                   545                                  550                                  555                                  560  
 Ser Arg Gln Asp Ile Ala Pro Ser Pro Ser Gln Arg Val Thr Gly Ser  
                                   565                                  570                                  575  
 Gln Leu Phe Leu Arg Leu Gln Tyr Phe Gly Gln Ser Leu Ser Gly Gly  
                                   580                                  585                                  590  
 Gln Asp Leu Thr Gln Asp Gly Leu Val Asp Leu Ala Val Gly Ala Gln  
                   595                                  600                                  605  
 Gly His Val Leu Leu Leu Arg Ser Leu Pro Leu Leu Lys Val Gly Ile  
                   610                                  615                                  620  
 Ser Ile Arg Phe Ala Pro Ser Glu Val Ala Lys Thr Val Tyr Gln Cys  
                   625                                  630                                  635                                  640  
 Trp Gly Arg Thr Pro Thr Val Leu Glu Ala Gly Glu Ala Thr Val Cys  
                                   645                                  650                                  655  
 Leu Thr Val Arg Lys Gly Ser Pro Asp Leu Leu Gly Asp Val Gln Ser  
                                   660                                  665                                  670  
 Ser Val Arg Tyr Asp Leu Ala Leu Asp Pro Gly Arg Leu Ile Ser Arg  
                   675                                  680                                  685  
 Ala Ile Phe Asp Glu Thr Lys Asn Cys Thr Leu Thr Arg Arg Lys Thr  
                   690                                  695                                  700  
 Leu Gly Leu Gly Asp His Cys Glu Thr Met Lys Leu Leu Leu Pro Asp  
                   705                                  710                                  715                                  720  
 Cys Val Glu Asp Ala Val Thr Pro Ile Ile Leu Arg Leu Asn Leu Ser  
                                   725                                  730                                  735  
 Leu Ala Gly Asp Ser Ala Pro Ser Arg Asn Leu Arg Pro Val Leu Ala  
                   740                                  745                                  750

- 126 -

Val Gly Ser Gln Asp His Val Thr Ala Ser Phe Pro Phe Glu Lys Asn  
 755 760 765  
 Cys Glu Gly Asn Leu Gly Val Ser Phe Asn Phe Ser Gly Leu Gln Val  
 770 775 780  
 Leu Glu Val Gly Ser Ser Pro Glu Leu Thr Val Thr Val Thr Val Trp  
 785 790 795 800  
 Asn Glu Gly Glu Asp Ser Tyr Gly Thr Leu Ile Lys Phe Tyr Tyr Pro  
 805 810 815  
 Ala Glu Leu Ser Tyr Arg Arg Val Thr Arg Ala Gln Gln Pro His Pro  
 820 825 830  
 Tyr Pro Leu Arg Leu Ala Cys Glu Ala Glu Pro Thr Gly Gln Glu Ser  
 835 840 845  
 Leu Arg Ser Ser Ser Cys Ser Ile Asn His Pro Ile Phe Arg Glu Gly  
 850 855 860  
 Ala Lys Ala Thr Phe Met Ile Thr Phe Asp Val Ser Tyr Lys Ala Phe  
 865 870 875 880  
 Leu Gly Asp Arg Leu Leu Leu Arg Ala Ser Ala Ser Ser Glu Asn Asn  
 885 890 895  
 Lys Pro Glu Thr Ser Lys Thr Ala Phe Gln Leu Glu Leu Pro Val Lys  
 900 905 910  
 Tyr Thr Val Tyr Thr Val Ile Ser Arg Gln Glu Asp Ser Thr Lys His  
 915 920 925  
 Phe Asn Phe Ser Ser Ser His Gly Glu Arg Gln Lys Glu Ala Glu His  
 930 935 940  
 Arg Tyr Arg Val Asn Asn Leu Ser Pro Leu Thr Leu Ala Ile Ser Val  
 945 950 955 960  
 Asn Phe Trp Val Pro Ile Leu Leu Asn Gly Val Ala Val Trp Asp Val  
 965 970 975  
 Thr Leu Arg Ser Pro Ala Gln Gly Val Ser Cys Val Ser Gln Arg Glu  
 980 985 990  
 Pro Pro Gln His Ser Asp Leu Leu Thr Gln Ile Gln Gly Arg Ser Val  
 995 1000 1005  
 Leu Asp Cys Ala Ile Ala Asp Cys Leu His Leu Arg Cys Asp Ile Pro  
 1010 1015 1020  
 Ser Leu Gly Thr Leu Asp Glu Leu Asp Phe Ile Leu Lys Gly Asn Leu  
 1025 1030 1035 1040  
 Ser Phe Gly Trp Ile Ser Gln Thr Leu Gln Lys Lys Val Leu Leu Leu  
 1045 1050 1055  
 Ser Glu Ala Glu Ile Thr Phe Asn Thr Ser Val Tyr Ser Gln Leu Pro  
 1060 1065 1070

- 127 -

Gly Gln Glu Ala Phe Leu Arg Ala Gln Val Ser Thr Met Leu Glu Glu  
 1075 1080 1085  
 Tyr Val Val Tyr Glu Pro Val Phe Leu Met Val Phe Ser Ser Val Gly  
 1090 1095 1100  
 Gly Leu Leu Leu Leu Ala Leu Ile Thr Val Ala Leu Tyr Lys Leu Gly  
 1105 1110 1115 1120  
 Phe Phe Lys Arg Gln Tyr Lys Glu Met Leu Asp Leu Pro Ser Ala Asp  
 1125 1130 1135  
 Pro Asp Pro Ala Gly Gln Ala Asp Ser Asn His Glu Thr Pro Pro His  
 1140 1145 1150  
 Leu Thr Ser  
 1155

## (2) INFORMATION FOR SEQ ID NO:47:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 49 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:47:

AGTTACGGAT CCGGCACCAT GACCTTCGGC ACTGTGATCC TCCTGTGTG

49

## (2) INFORMATION FOR SEQ ID NO:48:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 19 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:48:

GCTGGACGAT GGCATCCAC

19

## (2) INFORMATION FOR SEQ ID NO:49:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 24 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

- 128 -

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:49:

GTAGAGTTAC GGATCCGGCA CCAT

24

(2) INFORMATION FOR SEQ ID NO:50:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 20 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:50:

GCAGCCAGCT TCGGACAGAC

20

(2) INFORMATION FOR SEQ ID NO:51:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 21 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:51:

CCATGTCCAC AGAACAGAGA G

21

(2) INFORMATION FOR SEQ ID NO:52:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 3803 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(ix) FEATURE:

- (A) NAME/KEY: CDS
- (B) LOCATION: 1..3486

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:52:

ATG GTC CGT GGA GTT GTG ATC CTC CTG TGT GGC TGG GCC CTG GCT TCC  
 Met Val Arg Gly Val Val Ile Leu Leu Cys Gly Trp Ala Leu Ala Ser  
 1 5 10 15

48

TGT CAT GGG TCT AAC CTG GAT GTG GAG AAG CCC GTC GTG TTC AAA GAG  
 Cys His Gly Ser Asn Leu Asp Val Glu Lys Pro Val Val Phe Lys Glu  
 20 25 30

96

- 129 -

GAT	GCA	GCC	AGC	TTC	GGA	CAG	ACT	GTG	GTG	CAG	TTT	GGT	GGA	TCT	CGA	144
Asp	Ala	Ala	Ser	Phe	Gly	Gln	Thr	Val	Val	Gln	Phe	Gly	Gly	Ser	Arg	
		35					40					45				
CTC	GTG	GTG	GGA	GCC	CCT	CTG	GAG	GCG	GTG	GCA	GTC	AAC	CAA	ACA	GGA	192
Leu	Val	Val	Gly	Ala	Pro	Leu	Glu	Ala	Val	Ala	Val	Asn	Gln	Thr	Gly	
	50					55					60					
CAG	TCG	TCT	GAC	TGT	CCG	CCT	GCC	ACT	GGC	GTG	TGC	CAG	CCC	ATC	TTA	240
Gln	Ser	Ser	Asp	Cys	Pro	Pro	Ala	Thr	Gly	Val	Cys	Gln	Pro	Ile	Leu	
65					70					75					80	
CTG	CAC	ATT	CCC	CTA	GAG	GCA	GTG	AAC	ATG	TCC	CTG	GGC	CTG	TCT	CTG	288
Leu	His	Ile	Pro	Leu	Glu	Ala	Val	Asn	Met	Ser	Leu	Gly	Leu	Ser	Leu	
				85					90					95		
GTG	GCT	GAC	ACC	AAT	AAC	TCC	CAG	TTG	CTG	GCT	TGT	GGT	CCA	ACT	GCA	336
Val	Ala	Asp	Thr	Asn	Asn	Ser	Gln	Leu	Leu	Ala	Cys	Gly	Pro	Thr	Ala	
			100					105					110			
CAG	AGA	GCT	TGT	GCA	AAG	AAC	ATG	TAT	GCA	AAA	GGT	TCC	TGC	CTC	CTT	384
Gln	Arg	Ala	Cys	Ala	Lys	Asn	Met	Tyr	Ala	Lys	Gly	Ser	Cys	Leu	Leu	
		115					120					125				
CTG	GGC	TCC	AGC	TTG	CAG	TTC	ATC	CAG	GCA	ATC	CCT	GCT	ACC	ATG	CCA	432
Leu	Gly	Ser	Ser	Leu	Gln	Phe	Ile	Gln	Ala	Ile	Pro	Ala	Thr	Met	Pro	
	130					135					140					
GAG	TGT	CCA	GGA	CAA	GAG	ATG	GAC	ATT	GCT	TTC	CTG	ATT	GAT	GGC	TCC	480
Glu	Cys	Pro	Gly	Gln	Met	Asp	Ile	Ala	Phe	Leu	Ile	Asp	Gly	Ser		
145					150				155					160		
GGC	AGC	ATT	GAT	CAA	AGT	GAC	TTT	ACC	CAG	ATG	AAG	GAC	TTC	GTC	AAA	528
Gly	Ser	Ile	Asp	Gln	Ser	Asp	Phe	Thr	Gln	Met	Lys	Asp	Phe	Val	Lys	
				165					170					175		
GCT	TTG	ATG	GGC	CAG	TTG	GCG	AGC	ACC	AGC	ACC	TCG	TTC	TCC	CTG	ATG	576
Ala	Leu	Met	Gly	Gln	Leu	Ala	Ser	Thr	Ser	Thr	Ser	Phe	Ser	Leu	Met	
			180					185					190			
CAA	TAC	TCA	AAC	ATC	CTG	AAG	ACT	CAT	TTT	ACC	TTC	ACG	GAA	TTC	AAG	624
Gln	Tyr	Ser	Asn	Ile	Leu	Lys	Thr	His	Phe	Thr	Phe	Thr	Glu	Phe	Lys	
		195					200					205				
AGC	AGC	CTG	AGC	CCT	CAG	AGC	CTG	GTG	GAT	GCC	ATC	GTC	CAG	CTC	CAA	672
Ser	Ser	Leu	Ser	Pro	Gln	Ser	Leu	Val	Asp	Ala	Ile	Val	Gln	Leu	Gln	
		210				215					220					
GGC	CTG	ACG	TAC	ACA	GCC	TCG	GGC	ATC	CAG	AAA	GTG	GTG	AAA	GAG	CTA	720
Gly	Leu	Thr	Tyr	Thr	Ala	Ser	Gly	Ile	Gln	Lys	Val	Val	Lys	Glu	Leu	
225					230					235				240		
TTT	CAT	AGC	AAG	AAT	GGG	GCC	CGA	AAA	AGT	GCC	AAG	AAG	ATA	CTA	ATT	768
Phe	His	Ser	Lys	Asn	Gly	Ala	Arg	Lys	Ser	Ala	Lys	Lys	Ile	Leu	Ile	
				245					250					255		
GTC	ATC	ACA	GAT	GGG	CAG	AAA	TTC	AGA	GAC	CCC	CTG	GAG	TAT	AGA	CAT	816
Val	Ile	Thr	Asp	Gly	Gln	Lys	Phe	Arg	Asp	Pro	Leu	Glu	Tyr	Arg	His	
			260					265					270			

- 130 -

GTC ATC CCT GAA GCA GAG AAA GCT GGG ATC ATT CGC TAT GCT ATA GGG Val Ile Pro Glu Ala Glu Lys Ala Gly Ile Ile Arg Tyr Ala Ile Gly 275 280 285	864
GTG GGA GAT GCC TTC CGG GAA CCC ACT GCC CTA CAG GAG CTG AAC ACC Val Gly Asp Ala Phe Arg Glu Pro Thr Ala Leu Gln Glu Leu Asn Thr 290 295 300	912
ATT GGC TCA GCT CCC TCG CAG GAC CAC GTG TTC AAG GTG GGC AAT TTT Ile Gly Ser Ala Pro Ser Gln Asp His Val Phe Lys Val Gly Asn Phe 305 310 315 320	960
GTA GCA CTT CGC AGC ATC CAG CGG CAA ATT CAG GAG AAA ATC TTT GCC Val Ala Leu Arg Ser Ile Gln Arg Gln Ile Gln Glu Lys Ile Phe Ala 325 330 335	1008
ATT GAA GGA ACC GAA TCA AGG TCA AGT AGT TCC TTT CAG CAC GAG ATG Ile Glu Gly Thr Glu Ser Arg Ser Ser Ser Phe Gln His Glu Met 340 345 350	1056
TCA CAA GAA GGT TTC AGC TCA GCT CTC TCA ATG GAT GGA CCA GTT CTG Ser Gln Glu Gly Phe Ser Ser Ala Leu Ser Met Asp Gly Pro Val Leu 355 360 365	1104
GGG GCT GTG GGA GGC TTC AGC TGG TCT GGA GGT GCC TTC TTG TAC CCC Gly Ala Val Gly Gly Phe Ser Trp Ser Gly Gly Ala Phe Leu Tyr Pro 370 375 380	1152
TCA AAT ATG AGA TCC ACC TTC ATC AAC ATG TCT CAG GAG AAC GAG GAT Ser Asn Met Arg Ser Thr Phe Ile Asn Met Ser Gln Glu Asn Glu Asp 385 390 395 400	1200
ATG AGG GAC GCT TAC CTG GGT TAC TCC ACC GCA CTG GCC TTT TGG AAG Met Arg Asp Ala Tyr Leu Gly Tyr Ser Thr Ala Leu Ala Phe Trp Lys 405 410 415	1248
GGG GTC CAC AGC CTG ATC CTG GGG GCC CCT CGC CAC CAG CAC ACG GGG Gly Val His Ser Leu Ile Leu Gly Ala Pro Arg His Gln His Thr Gly 420 425 430	1296
AAG GTT GTC ATC TTT ACC CAG GAA TCC AGG CAC TGG AGG CCC AAG TCT Lys Val Val Ile Phe Thr Gln Glu Ser Arg His Trp Arg Pro Lys Ser 435 440 445	1344
GAA GTC AGA GGG ACA CAG ATC GGC TCC TAC TTT GGG GCA TCT CTC TGT Glu Val Arg Gly Thr Gln Ile Gly Ser Tyr Phe Gly Ala Ser Leu Cys 450 455 460	1392
TCT GTG GAC ATG GAT AGA GAT GGC AGC ACT GAC CTG GTC CTG ATT GGA Ser Val Asp Met Asp Arg Asp Gly Ser Thr Asp Leu Val Leu Ile Gly 465 470 475 480	1440
GTC CCC CAT TAC TAT GAG CAC ACC CGA GGG GGG CAG GTG TCG GTG TGC Val Pro His Tyr Tyr Glu His Thr Arg Gly Gly Gln Val Ser Val Cys 485 490 495	1488
CCC ATG CCT GGT GTG AGG AGC AGG TGG CAT TGT GGG ACC ACC CTC CAT Pro Met Pro Gly Val Arg Ser Arg Trp His Cys Gly Thr Thr Leu His 500 505 510	1536

- 131 -

GGG GAG CAG GGC CAT CCT TGG GGC CGC TTT GGG GCG GCT CTG ACA GTG Gly Glu Gln Gly His Pro Trp Gly Arg Phe Gly Ala Ala Leu Thr Val 515 520 525	1584
CTA GGG GAC GTG AAT GGG GAC AGT CTG GCG GAT GTG GCT ATT GGT GCA Leu Gly Asp Val Asn Gly Asp Ser Leu Ala Asp Val Ala Ile Gly Ala 530 535 540	1632
CCC GGA GAG GAG GAG AAC AGA GGT GCT GTC TAC ATA TTT CAT GGA GCC Pro Gly Glu Glu Glu Asn Arg Gly Ala Val Tyr Ile Phe His Gly Ala 545 550 555 560	1680
TCG AGA CAG GAC ATC GCT CCC TCG CCT AGC CAG CGG GTC ACT GGC TCC Ser Arg Gln Asp Ile Ala Pro Ser Pro Ser Gln Arg Val Thr Gly Ser 565 570 575	1728
CAG CTC TTC CTG AGG CTC CAA TAT TTT GGG CAG TCA TTA AGT GGG GGT Gln Leu Phe Leu Arg Leu Gln Tyr Phe Gly Gln Ser Leu Ser Gly Gly 580 585 590	1776
CAG GAC CTT ACA CAG GAT GGC CTG GTG GAC CTG GCC GTG GGA GCC CAG Gln Asp Leu Thr Gln Asp Gly Leu Val Asp Leu Ala Val Gly Ala Gln 595 600 605	1824
GGG CAC GTG CTG CTG CTT AGG AGT CTG CCT TTG CTG AAA GTG GGG ATC Gly His Val Leu Leu Leu Arg Ser Leu Pro Leu Leu Lys Val Gly Ile 610 615 620	1872
TCC ATT AGA TTT GCC CCC TCA GAG GTG GCA AAG ACT GTG TAC CAG TGC Ser Ile Arg Phe Ala Pro Ser Glu Val Ala Lys Thr Val Tyr Gln Cys 625 630 635 640	1920
TGG GGA AGG ACT CCC ACT GTC CTC GAA GCT GGA GAG GCC ACC GTC TGT Trp Gly Arg Thr Pro Thr Val Leu Glu Ala Gly Glu Ala Thr Val Cys 645 650 655	1968
CTC ACT GTC CGC AAA GGT TCA CCT GAC CTG TTA GGT GAT GTC CAA AGC Leu Thr Val Arg Lys Gly Ser Pro Asp Leu Leu Gly Asp Val Gln Ser 660 665 670	2016
TCT GTC AGG TAT GAT CTG GCG TTG GAT CCG GGC CGT CTG ATT TCT CGT Ser Val Arg Tyr Asp Leu Ala Leu Asp Pro Gly Arg Leu Ile Ser Arg 675 680 685	2064
GCC ATT TTT GAT GAG ACG AAG AAC TGC ACT TTG ACC CGA AGG AAG ACT Ala Ile Phe Asp Glu Thr Lys Asn Cys Thr Leu Thr Arg Arg Lys Thr 690 695 700	2112
CTG GGG CTT GGT GAT CAC TGC GAA ACA ATG AAG CTG CTT TTG CCA GAC Leu Gly Leu Gly Asp His Cys Glu Thr Met Lys Leu Leu Leu Pro Asp 705 710 715 720	2160
TGT GTG GAG GAT GCA GTG ACC CCT ATC ATC CTG CGC CTT AAC TTA TCC Cys Val Glu Asp Ala Val Thr Pro Ile Ile Leu Arg Leu Asn Leu Ser 725 730 735	2208
CTG GCA GGG GAC TCT GCT CCA TCC AGG AAC CTT CGT CCT GTG CTG GCT Leu Ala Gly Asp Ser Ala Pro Ser Arg Asn Leu Arg Pro Val Leu Ala 740 745 750	2256

- 132 -

GTG GGC TCA CAA GAC CAT GTA ACA GCT TCT TTC CCG TTT GAG AAG AAC Val Gly Ser Gln Asp His Val Thr Ala Ser Phe Pro Phe Glu Lys Asn 755 760 765	2304
TGT AAG CAG GAG CTC CTG TGT GAG GGG AAC CTG GGC GTC AGC TTC AAC Cys Lys Gln Glu Leu Leu Cys Glu Gly Asn Leu Gly Val Ser Phe Asn 770 775 780	2352
TTC TCA GGC CTG CAG GTC TTG GAG GTA GGA AGC TCC CCA GAG CTC ACT Phe Ser Gly Leu Gln Val Leu Glu Val Gly Ser Ser Pro Glu Leu Thr 785 790 795 800	2400
GTG ACA GTA ACA GTT TGG AAT GAG GGT GAG GAC AGC TAT GGA ACC TTA Val Thr Val Thr Val Trp Asn Glu Gly Glu Asp Ser Tyr Gly Thr Leu 805 810 815	2448
ATC AAG TTC TAC TAC CCA GCA GAG CTA TCT TAC CGA CGG GTG ACA AGA Ile Lys Phe Tyr Tyr Pro Ala Glu Leu Ser Tyr Arg Arg Val Thr Arg 820 825 830	2496
GCC CAG CAA CCT CAT CCG TAC CCA CTA CGC CTG GCA TGT GAG GCT GAG Ala Gln Gln Pro His Pro Tyr Pro Leu Arg Leu Ala Cys Glu Ala Glu 835 840 845	2544
CCC ACG GGC CAG GAG AGC CTG AGG AGC AGC AGC TGT AGC ATC AAT CAC Pro Thr Gly Gln Glu Ser Leu Arg Ser Ser Ser Cys Ser Ile Asn His 850 855 860	2592
CCC ATC TTC CGA GAA GGT GCC AAG GCC ACC TTC ATG ATC ACA TTT GAT Pro Ile Phe Arg Glu Gly Ala Lys Ala Thr Phe Met Ile Thr Phe Asp 865 870 875 880	2640
CTC TCC TAC AAG GCC TTC CTG GGA GAC AGG TTG CTT CTG AGG GCC AGC Val Ser Tyr Lys Ala Phe Leu Gly Asp Arg Leu Leu Leu Arg Ala Ser 885 890 895	2688
GCA AGC AGT GAG AAT AAT AAG CCT GAA ACC AGC AAG ACT GCC TTC CAG Ala Ser Ser Glu Asn Asn Lys Pro Glu Thr Ser Lys Thr Ala Phe Gln 900 905 910	2736
CTG GAG CTT CCG GTG AAG TAC ACG GTC TAT ACC GTG ATC AGT AGG CAG Leu Glu Leu Pro Val Lys Tyr Thr Val Tyr Thr Val Ile Ser Arg Gln 915 920 925	2784
GAA GAT TCT ACC AAG CAT TTC AAC TTC TCA TCT TCC CAC GGG GAG AGA Glu Asp Ser Thr Lys His Phe Asn Phe Ser Ser Ser His Gly Glu Arg 930 935 940	2832
CAG AAA GAG GCC GAA CAT CGA TAT CGT GTG AAT AAC CTG AGT CCA TTG Gln Lys Glu Ala Glu His Arg Tyr Arg Val Asn Asn Leu Ser Pro Leu 945 950 955 960	2880
ACG CTG GCC ATC AGC GTT AAC TTC TGG GTC CCC ATC CTT CTG AAT GGT Thr Leu Ala Ile Ser Val Asn Phe Trp Val Pro Ile Leu Leu Asn Gly 965 970 975	2928
TGT GCC GTG TGG GAT GTG ACT CTG AGG AGC CCA GCA CAG GGT GTC TCC Ala Val Trp Asp Val Thr Leu Arg Ser Pro Ala Gln Gly Val Ser 980 985 990	2976



- 133 -

TGT GTG TCA CAG AGG GAA CCT CCT CAA CAT TCC GAC CTT CTG ACC CAG Cys Val Ser Gln Arg Glu Pro Pro Gln His Ser Asp Leu Leu Thr Gln 995 1000 1005	3024
ATC CAA GGA CGC TCT GTG CTG GAC TGC GCC ATC GCC GAC TGC CTG CAC Ile Gln Gly Arg Ser Val Leu Asp Cys Ala Ile Ala Asp Cys Leu His 1010 1015 1020	3072
CTC CGC TGT GAC ATC CCC TCC TTG GGC ACC CTG GAT GAG CTT GAC TTC Leu Arg Cys Asp Ile Pro Ser Leu Gly Thr Leu Asp Glu Leu Asp Phe 1025 1030 1035 1040	3120
ATT CTG AAG GGC AAC CTC AGC TTC GGC TGG ATC AGT CAG ACA TTG CAG Ile Leu Lys Gly Asn Leu Ser Phe Gly Trp Ile Ser Gln Thr Leu Gln 1045 1050 1055	3168
AAA AAG GTG TTG CTC CTG AGT GAG GCT GAA ATC ACA TTC AAC ACA TCT Lys Lys Val Leu Leu Leu Ser Glu Ala Glu Ile Thr Phe Asn Thr Ser 1060 1065 1070	3216
GTG TAT TCC CAG CTG CCG GGA CAG GAG GCA TTT CTG AGA GCC CAG GTG Val Tyr Ser Gln Leu Pro Gly Gln Glu Ala Phe Leu Arg Ala Gln Val 1075 1080 1085	3264
TCA ACG ATG CTA GAA GAA TAC GTG GTC TAT GAG CCC GTC TTC CTC ATG Ser Thr Met Leu Glu Glu Tyr Val Val Tyr Glu Pro Val Phe Leu Met 1090 1095 1100	3312
GTG TTC AGC TCA GTG GGA GGT CTG CTG TTA CTG GCT CTC ATC ACT GTG Val Phe Ser Ser Val Gly Gly Leu Leu Leu Leu Ala Leu Ile Thr Val 1105 1110 1115 1120	3360
GCG CTG TAC AAG CTT GGC TTC TTC AAA CGT CAG TAT AAA GAG ATG CTG Ala Leu Tyr Lys Leu Gly Phe Phe Lys Arg Gln Tyr Lys Glu Met Leu 1125 1130 1135	3408
GAT CTA CCA TCT GCA GAT CCT GAC CCA GCC GGC CAG GCA GAT TCC AAC Asp Leu Pro Ser Ala Asp Pro Asp Pro Ala Gly Gln Ala Asp Ser Asn 1140 1145 1150	3456
CAT GAG ACT CCT CCA CAT CTC ACG TCC TAGGAATCTA CTTTCCTGTA His Glu Thr Pro Pro His Leu Thr Ser 1155 1160	3503
TATCTCCACA ATTACGAGAT TGGTTTTGCT TTTGCCTATG AATCTACTGG CATGGGAACA	3563
AGTTCTCTTC AGCTCTGGGC TAGCCTGGGA AACTTCCCAG AAATGATGCC CTACCTCCTG	3623
AGCTGGGAGA TTTTATGGT TTGCCCATGT GTCAGATTTT AGTGCTGATC CACTTTTTTT	3683
GCAAGAGCAG GAATGGGGTC AGCATAAATT TACATATGGA TAAGAACTAA CACAAGACTG	3743
AGTAATATGC TCAATATTCA ATGTATTGCT TGTATAAATT TTTAAAAAAT AAAATGAAAN	3803

## (2) INFORMATION FOR SEQ ID NO:53:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1161 amino acids  
(B) TYPE: amino acid

- 134 -

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:53:

```

Met Val Arg Gly Val Val Ile Leu Leu Cys Gly Trp Ala Leu Ala Ser
 1           5           10           15
Cys His Gly Ser Asn Leu Asp Val Glu Lys Pro Val Val Phe Lys Glu
          20           25           30
Asp Ala Ala Ser Phe Gly Gln Thr Val Val Gln Phe Gly Gly Ser Arg
          35           40           45
Leu Val Val Gly Ala Pro Leu Glu Ala Val Ala Val Asn Gln Thr Gly
          50           55           60
Gln Ser Ser Asp Cys Pro Pro Ala Thr Gly Val Cys Gln Pro Ile Leu
          65           70           75           80
Leu His Ile Pro Leu Glu Ala Val Asn Met Ser Leu Gly Leu Ser Leu
          85           90           95
Val Ala Asp Thr Asn Asn Ser Gln Leu Leu Ala Cys Gly Pro Thr Ala
          100          105          110
Gln Arg Ala Cys Ala Lys Asn Met Tyr Ala Lys Gly Ser Cys Leu Leu
          115          120          125
Leu Gly Ser Ser Leu Gln Phe Ile Gln Ala Ile Pro Ala Thr Met Pro
          130          135          140
Glu Cys Pro Gly Gln Glu Met Asp Ile Ala Phe Leu Ile Asp Gly Ser
          145          150          155          160
Gly Ser Ile Asp Gln Ser Asp Phe Thr Gln Met Lys Asp Phe Val Lys
          165          170          175
Ala Leu Met Gly Gln Leu Ala Ser Thr Ser Thr Ser Phe Ser Leu Met
          180          185          190
Gln Tyr Ser Asn Ile Leu Lys Thr His Phe Thr Phe Thr Glu Phe Lys
          195          200          205
Ser Ser Leu Ser Pro Gln Ser Leu Val Asp Ala Ile Val Gln Leu Gln
          210          215          220
Gly Leu Thr Tyr Thr Ala Ser Gly Ile Gln Lys Val Val Lys Glu Leu
          225          230          235          240
Phe His Ser Lys Asn Gly Ala Arg Lys Ser Ala Lys Lys Ile Leu Ile
          245          250          255
Val Ile Thr Asp Gly Gln Lys Phe Arg Asp Pro Leu Glu Tyr Arg His
          260          265          270
Val Ile Pro Glu Ala Glu Lys Ala Gly Ile Ile Arg Tyr Ala Ile Gly
          275          280          285

```

- 135 -

Val Gly Asp Ala Phe Arg Glu Pro Thr Ala Leu Gln Glu Leu Asn Thr  
 290 295 300  
 Ile Gly Ser Ala Pro Ser Gln Asp His Val Phe Lys Val Gly Asn Phe  
 305 310 315 320  
 Val Ala Leu Arg Ser Ile Gln Arg Gln Ile Gln Glu Lys Ile Phe Ala  
 325 330 335  
 Ile Glu Gly Thr Glu Ser Arg Ser Ser Ser Phe Gln His Glu Met  
 340 345 350  
 Ser Gln Glu Gly Phe Ser Ser Ala Leu Ser Met Asp Gly Pro Val Leu  
 355 360 365  
 Gly Ala Val Gly Gly Phe Ser Trp Ser Gly Gly Ala Phe Leu Tyr Pro  
 370 375 380  
 Ser Asn Met Arg Ser Thr Phe Ile Asn Met Ser Gln Glu Asn Glu Asp  
 385 390 395 400  
 Met Arg Asp Ala Tyr Leu Gly Tyr Ser Thr Ala Leu Ala Phe Trp Lys  
 405 410 415  
 Gly Val His Ser Leu Ile Leu Gly Ala Pro Arg His Gln His Thr Gly  
 420 425 430  
 Lys Val Val Ile Phe Thr Gln Glu Ser Arg His Trp Arg Pro Lys Ser  
 435 440 445  
 Glu Val Arg Gly Thr Gln Ile Gly Ser Tyr Phe Gly Ala Ser Leu Cys  
 450 455 460  
 Ser Val Asp Met Asp Arg Asp Gly Ser Thr Asp Leu Val Leu Ile Gly  
 465 470 475 480  
 Val Pro His Tyr Tyr Glu His Thr Arg Gly Gly Gln Val Ser Val Cys  
 485 490 495  
 Pro Met Pro Gly Val Arg Ser Arg Trp His Cys Gly Thr Thr Leu His  
 500 505 510  
 Gly Glu Gln Gly His Pro Trp Gly Arg Phe Gly Ala Ala Leu Thr Val  
 515 520 525  
 Leu Gly Asp Val Asn Gly Asp Ser Leu Ala Asp Val Ala Ile Gly Ala  
 530 535 540  
 Pro Gly Glu Glu Glu Asn Arg Gly Ala Val Tyr Ile Phe His Gly Ala  
 545 550 555 560  
 Ser Arg Gln Asp Ile Ala Pro Ser Pro Ser Gln Arg Val Thr Gly Ser  
 565 570 575  
 Gln Leu Phe Leu Arg Leu Gln Tyr Phe Gly Gln Ser Leu Ser Gly Gly  
 580 585 590  
 Gln Asp Leu Thr Gln Asp Gly Leu Val Asp Leu Ala Val Gly Ala Gln  
 595 600 605

- 136 -

Gly His Val Leu Leu Leu Arg Ser Leu Pro Leu Leu Lys Val Gly Ile  
 610 615 620  
 Ser Ile Arg Phe Ala Pro Ser Glu Val Ala Lys Thr Val Tyr Gln Cys  
 625 630 635 640  
 Trp Gly Arg Thr Pro Thr Val Leu Glu Ala Gly Glu Ala Thr Val Cys  
 645 650 655  
 Leu Thr Val Arg Lys Gly Ser Pro Asp Leu Leu Gly Asp Val Gln Ser  
 660 665 670  
 Ser Val Arg Tyr Asp Leu Ala Leu Asp Pro Gly Arg Leu Ile Ser Arg  
 675 680 685  
 Ala Ile Phe Asp Glu Thr Lys Asn Cys Thr Leu Thr Arg Arg Lys Thr  
 690 695 700  
 Leu Gly Leu Gly Asp His Cys Glu Thr Met Lys Leu Leu Leu Pro Asp  
 705 710 715 720  
 Cys Val Glu Asp Ala Val Thr Pro Ile Ile Leu Arg Leu Asn Leu Ser  
 725 730 735  
 Leu Ala Gly Asp Ser Ala Pro Ser Arg Asn Leu Arg Pro Val Leu Ala  
 740 745 750  
 Val Gly Ser Gln Asp His Val Thr Ala Ser Phe Pro Phe Glu Lys Asn  
 755 760 765  
 Cys Lys Gln Glu Leu Leu Cys Glu Gly Asn Leu Gly Val Ser Phe Asn  
 770 775 780  
 Phe Ser Gly Leu Gln Val Leu Glu Val Gly Ser Ser Pro Glu Leu Thr  
 785 790 795 800  
 Val Thr Val Thr Val Trp Asn Glu Gly Glu Asp Ser Tyr Gly Thr Leu  
 805 810 815  
 Ile Lys Phe Tyr Tyr Pro Ala Glu Leu Ser Tyr Arg Arg Val Thr Arg  
 820 825 830  
 Ala Gln Gln Pro His Pro Tyr Pro Leu Arg Leu Ala Cys Glu Ala Glu  
 835 840 845  
 Pro Thr Gly Gln Glu Ser Leu Arg Ser Ser Ser Cys Ser Ile Asn His  
 850 855 860  
 Pro Ile Phe Arg Glu Gly Ala Lys Ala Thr Phe Met Ile Thr Phe Asp  
 865 870 875 880  
 Val Ser Tyr Lys Ala Phe Leu Gly Asp Arg Leu Leu Leu Arg Ala Ser  
 885 890 895  
 Ala Ser Ser Glu Asn Asn Lys Pro Glu Thr Ser Lys Thr Ala Phe Gln  
 900 905 910  
 Leu Glu Leu Pro Val Lys Tyr Thr Val Tyr Thr Val Ile Ser Arg Gln  
 915 920 925

- 137 -

Glu Asp Ser Thr Lys His Phe Asn Phe Ser Ser Ser His Gly Glu Arg  
 930 935 940  
 Gln Lys Glu Ala Glu His Arg Tyr Arg Val Asn Asn Leu Ser Pro Leu  
 945 950 955 960  
 Thr Leu Ala Ile Ser Val Asn Phe Trp Val Pro Ile Leu Leu Asn Gly  
 965 970 975  
 Val Ala Val Trp Asp Val Thr Leu Arg Ser Pro Ala Gln Gly Val Ser  
 980 985 990  
 Cys Val Ser Gln Arg Glu Pro Pro Gln His Ser Asp Leu Leu Thr Gln  
 995 1000 1005  
 Ile Gln Gly Arg Ser Val Leu Asp Cys Ala Ile Ala Asp Cys Leu His  
 1010 1015 1020  
 Leu Arg Cys Asp Ile Pro Ser Leu Gly Thr Leu Asp Glu Leu Asp Phe  
 1025 1030 1035 1040  
 Ile Leu Lys Gly Asn Leu Ser Phe Gly Trp Ile Ser Gln Thr Leu Gln  
 1045 1050 1055  
 Lys Lys Val Leu Leu Leu Ser Glu Ala Glu Ile Thr Phe Asn Thr Ser  
 1060 1065 1070  
 Val Tyr Ser Gln Leu Pro Gly Gln Glu Ala Phe Leu Arg Ala Gln Val  
 1075 1080 1085  
 Ser Thr Met Leu Glu Glu Tyr Val Val Tyr Glu Pro Val Phe Leu Met  
 1090 1095 1100  
 Val Phe Ser Ser Val Gly Gly Leu Leu Leu Leu Ala Leu Ile Thr Val  
 1105 1110 1115 1120  
 Ala Leu Tyr Lys Leu Gly Phe Phe Lys Arg Gln Tyr Lys Glu Met Leu  
 1125 1130 1135  
 Asp Leu Pro Ser Ala Asp Pro Asp Pro Ala Gly Gln Ala Asp Ser Asn  
 1140 1145 1150  
 His Glu Thr Pro Pro His Leu Thr Ser  
 1155 1160

## (2) INFORMATION FOR SEQ ID NO:54:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 3597 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

- (ix) FEATURE:
- (A) NAME/KEY: CDS
  - (B) LOCATION: 40..3525

- 138 -

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:54:

AGCTTTACAG CTCTCTACTT CTCAGTGCAC TGCTCAGTG ATG GCC GGT GGA GTT	54
Met Ala Gly Gly Val	
1 5	
GTG ATC CTC CTG TGT GGC TGG GTC CTG GCT TCC TGT CAT GGG TCT AAC	102
Val Ile Leu Leu Cys Gly Trp Val Leu Ala Ser Cys His Gly Ser Asn	
10 15 20	
CTG GAT GTG GAG GAA CCC ATC GTG TTC AGA GAG GAT GCA GCC AGC TTT	150
Leu Asp Val Glu Pro Ile Val Phe Arg Glu Asp Ala Ala Ser Phe	
25 30 35	
GGA CAG ACT GTG GTG CAG TTT GGT GGA TCT CGA CTC GTG GTG GGA GCC	198
Gly Gln Thr Val Val Gln Phe Gly Ser Arg Leu Val Val Gly Ala	
40 45 50	
CCT CTG GAG GCG GTG GCA GTC AAC CAA ACA GGA CGG TTG TAT GAC TGT	246
Pro Leu Glu Ala Val Ala Val Asn Gln Thr Gly Arg Leu Tyr Asp Cys	
55 60 65	
GCA CCT GCC ACT GGC ATG TGC CAG CCC ATC GTA CTG CGC AGT CCC CTA	294
Ala Pro Ala Thr Gly Met Cys Gln Pro Ile Val Leu Arg Ser Pro Leu	
70 75 80 85	
GAG GCA GTG AAC ATG TCC CTG GGC CTG TCT CTG GTG ACT GCC ACC AAT	342
Glu Ala Val Asn Met Ser Leu Gly Leu Ser Leu Val Thr Ala Thr Asn	
90 95 100	
AAC GCC CAG TTG CTG GCT TGT GGT CCA ACT GCA CAG AGA GCT TGT GTG	390
Asn Ala Gln Leu Leu Ala Cys Gly Pro Thr Ala Gln Arg Ala Cys Val	
105 110 115	
AAG AAC ATG TAT GCG AAA GGT TCC TGC CTC CTT CTC GGC TCC AGC TTG	438
Lys Asn Met Tyr Ala Lys Gly Ser Cys Leu Leu Leu Ser Ser Leu	
120 125 130	
CAG TTC ATC CAG GCA GTC CCT GCC TCC ATG CCA GAG TGT CCA AGA CAA	486
Gln Phe Ile Gln Ala Val Pro Ala Ser Met Pro Glu Cys Pro Arg Gln	
135 140 145	
GAG ATG GAC ATT GCT TTC CTG ATT GAT GGT TCT GGC AGC ATT AAC CAA	534
Glu Met Asp Ile Ala Phe Leu Ile Asp Gly Ser Gly Ser Ile Asn Gln	
150 155 160 165	
AGG GAC TTT GCC CAG ATG AAG GAC TTT GTC AAA GCT TTG ATG GGA GAG	582
Arg Asp Phe Ala Gln Met Lys Asp Phe Val Lys Ala Leu Met Gly Glu	
170 175 180	
TTT GCG AGC ACC AGC ACC TTG TTC TCC CTG ATG CAA TAC TCG AAC ATC	630
Phe Ala Ser Thr Ser Thr Leu Phe Ser Leu Met Gln Tyr Ser Asn Ile	
185 190 195	
CTG AAG ACC CAT TTT ACC TTC ACT GAA TTC AAG AAC ATC CTG GAC CCT	678
Leu Lys Thr His Phe Thr Phe Thr Glu Phe Lys Asn Ile Leu Asp Pro	
200 205 210	
CAG AGC CTG GTG GAT CCC ATT GTC CAG CTG CAA GGC CTG ACC TAC ACA	726
Gln Ser Leu Val Asp Pro Ile Val Gln Leu Gln Gly Leu Thr Tyr Thr	
215 220 225	

- 139 -

GCC Ala 230	ACA Thr 230	GGC Gly 230	ATC Ile 230	CGG Arg 235	ACA Thr 235	GTG Val 235	ATG Met 235	GAA Glu 240	GAG Glu 240	CTA Leu 240	TTT Phe 240	CAT His 240	AGC Ser 245	AAG Lys 245	AAT Asn 245	774
GGG Gly 250	TCC Ser 250	CGT Arg 250	AAA Lys 250	AGT Ser 250	GCC Ala 255	AAG Lys 255	AAG Lys 255	ATC Ile 255	CTC Leu 255	CTT Leu 260	GTC Val 260	ATC Ile 260	ACA Thr 260	GAT Asp 260	GGG Gly 260	822
CAG Gln 265	AAA Lys 265	TAC Tyr 265	AGA Arg 265	GAC Asp 265	CCC Pro 270	CTG Leu 270	GAG Glu 270	TAT Tyr 270	AGT Ser 270	GAT Asp 275	GTC Val 275	ATT Ile 275	CCC Pro 275	GCC Ala 275	GCA Ala 275	870
GAC Asp 280	AAA Lys 280	GCT Ala 280	GGC Gly 280	ATC Ile 285	ATT Ile 285	CGT Arg 285	TAT Tyr 285	GCT Ala 285	ATT Ile 285	GGG Gly 290	GTG Val 290	GGA Gly 290	GAT Asp 290	GCC Ala 290	TTC Phe 290	918
CAG Gln 295	GAG Glu 295	CCC Pro 295	ACT Thr 295	GCC Ala 300	CTG Leu 300	AAG Lys 300	GAG Glu 305	CTG Leu 305	AAC Asn 305	ACC Thr 305	ATT Ile 305	GGC Gly 305	TCA Ser 305	GCT Ala 305	CCC Pro 305	966
CCA Pro 310	CAG Gln 310	GAC Asp 315	CAC His 315	GTG Val 315	TTC Phe 315	AAG Lys 315	GTA Val 315	GGC Gly 320	AAC Asn 320	TTT Phe 320	GCA Ala 320	GCA Ala 320	CTT Leu 325	CGC Arg 325	AGC Ser 325	1014
ATC Ile 330	CAG Gln 330	AGG Arg 330	CAA Gln 330	CTT Leu 330	CAG Gln 330	GAG Glu 335	AAA Lys 335	ATC Ile 335	TTC Phe 335	GCC Ala 335	ATT Ile 340	GAG Glu 340	GGA Gly 340	ACT Thr 340	CAA Gln 340	1062
TCA Ser 345	AGG Arg 345	TCA Ser 345	AGT Ser 345	AGT Ser 345	TCC Ser 350	TTT Phe 350	CAG Gln 350	CAC His 350	GAG Glu 350	ATG Met 355	TCA Ser 355	CAA Gln 355	GAA Glu 355	GGT Gly 355	TTC Phe 355	1110
AGT Ser 360	TCA Ser 360	GCT Ala 360	CTC Leu 360	ACA Thr 365	TCG Ser 365	GAT Asp 365	GGA Gly 365	CCC Pro 370	GTT Val 370	CTG Leu 370	GGG Gly 370	GCC Ala 370	GTG Val 370	GGA Gly 370	AGC Ser 370	1158
TTC Phe 375	AGC Ser 375	TGG Trp 375	TCC Ser 380	GGA Gly 380	GGT Gly 380	GCC Ala 380	TTC Phe 380	TTA Leu 385	TAT Tyr 385	CCC Pro 385	CCA Pro 385	AAT Asn 385	ACG Thr 385	AGA Arg 385	CCC Pro 385	1206
ACC Thr 390	TTT Phe 390	ATC Ile 395	AAC Asn 395	ATG Met 395	TCT Ser 395	CAG Gln 400	GAG Glu 400	AAT Asn 400	GTG Val 400	GAC Met 400	ATG Met 405	AGA Arg 405	GAC Asp 405	TCC Ser 405	TAC Tyr 405	1254
CTG Leu 410	GGT Gly 410	TAC Tyr 410	TCC Ser 410	ACC Thr 410	GCA Ala 415	GTG Val 415	GCC Ala 415	TTT Phe 415	TGG Trp 415	AAG Lys 420	GGG Gly 420	GTT Val 420	CAC His 420	AGC Ser 420	CTG Leu 420	1302
ATC Ile 425	CTG Leu 425	GGG Gly 425	GCC Ala 430	CCG Pro 430	CGT Arg 430	CAC His 430	CAG Gln 430	CAC His 430	ACG Thr 435	GGG Gly 435	AAG Lys 435	GTT Val 435	GTC Val 435	ATC Ile 435	TTT Phe 435	1350
ACC Thr 440	CAG Gln 440	GAA Glu 440	GCC Ala 445	AGG Arg 445	CAT His 445	TGG Trp 445	AGG Arg 445	CCC Pro 450	AAG Lys 450	TCT Ser 450	GAA Glu 450	GTC Val 450	AGA Arg 450	GGG Gly 450	ACA Thr 450	1398
CAG Gln 455	ATC Ile 455	GGC Gly 455	TCC Ser 460	TAC Tyr 460	TTC Phe 460	GGG Gly 460	GCC Ala 460	TCT Ser 465	CTC Leu 465	TGT Cys 465	TCT Ser 465	GTG Val 465	GAC Asp 465	GTG Val 465	GAT Asp 465	1446

- 140 -

AGA Arg 470	GAT Asp 470	GGC Gly 470	AGC Ser 470	ACY Xaa 475	GAC Asp 475	CTG Leu 475	GTC Val 475	CTG Leu 475	ATC Ile 480	GGA Gly 480	GCC Ala 480	CCC Pro 480	CAT His 485	TAC Tyr 485	TAT Tyr 485	1494
GAG Glu 490	CAG Gln 490	ACC Thr 490	CGA Arg 490	GGG Gly 490	GGG Gly 490	CAG Gln 495	GTC Val 495	TCA Ser 495	GTG Val 495	TTC Phe 495	CCC Pro 500	GTG Val 500	CCC Pro 500	GGT Gly 500	GTG Val 500	1542
AGG Arg 505	GGC Gly 505	AGG Arg 505	TGG Trp 505	CAG Gln 505	TGT Cys 510	GAG Glu 510	GCC Ala 510	ACC Thr 510	CTC Leu 510	CAC His 515	GGG Gly 515	GAG Glu 515	CAG Gln 515	GGC Gly 515	CAT His 515	1590
CCT Pro 520	TGG Trp 520	GGC Gly 520	CGC Arg 520	TTT Phe 520	GGG Gly 525	GTG Val 525	GCT Ala 525	CTG Leu 525	ACA Thr 530	GTG Val 530	CTG Leu 530	GGG Gly 530	GAC Asp 530	GTA Val 530	AAC Asn 530	1638
GGG Gly 535	GAC Asp 535	AAT Asn 535	CTG Leu 535	GCA Ala 535	GAC Asp 540	GTG Val 540	GCT Ala 540	ATT Ile 540	GGT Gly 545	GCC Ala 545	CCT Pro 545	GGA Gly 545	GAG Glu 545	GAG Glu 545	GAG Glu 545	1686
AGC Ser 550	AGA Arg 550	GGT Gly 550	GCT Ala 550	GTC Val 555	TAC Tyr 555	ATA Ile 555	TTT Phe 555	CAT His 560	GGA Gly 560	GCC Ala 560	TCG Ser 560	AGA Arg 560	CTG Leu 565	GAG Glu 565	ATC Ile 565	1734
ATG Met 570	CCC Pro 570	TCA Ser 570	CCC Pro 570	AGC Ser 570	CAG Gln 570	CGG Arg 570	GTC Val 575	ACT Thr 575	GGC Gly 575	TCC Ser 575	CAG Gln 580	CTC Leu 580	TCC Ser 580	CTG Leu 580	AGA Arg 580	1782
CTG Leu 585	CAG Gln 585	TAT Tyr 585	TTT Phe 585	GGG Gly 585	CAG Gln 590	TCA Ser 590	TTG Leu 590	AGT Ser 590	GGG Gly 590	GGT Gly 590	CAG Gln 595	GAC Asp 595	CTT Leu 595	ACA Thr 595	CAG Gln 595	1830
GAT Asp 600	GGC Gly 600	CTG Leu 600	GTG Val 600	GAC Asp 605	CTG Leu 605	GCC Ala 605	GTG Val 605	GGA Gly 605	GCC Ala 610	CAG Gln 610	GGG Gly 610	CAC His 610	GTA Val 610	CTG Leu 610	CTG Leu 610	1878
CTC Leu 615	AGG Arg 615	ACT Ser 615	CTG Leu 615	CCT Pro 620	CTG Leu 620	CTG Leu 620	AAA Lys 620	GTG Val 625	GAG Glu 625	CTC Leu 625	TCC Ser 625	ATA Ile 625	AGA Arg 625	TTC Phe 625	GCC Ala 625	1926
CCC Pro 630	ATG Met 630	GAG Glu 630	GTG Val 630	GCA Ala 635	AAG Lys 635	GCT Ala 635	GTG Val 635	TAC Tyr 640	CAG Gln 640	TGC Cys 640	TGG Trp 640	GAA Glu 645	AGG Arg 645	ACT Thr 645	CCC Pro 645	1974
ACT Thr 650	GTC Val 650	CTC Leu 650	GAA Glu 650	GCT Ala 650	GGA Gly 650	GAG Glu 655	GCC Ala 655	ACT Thr 655	GTC Val 655	TGT Cys 655	CTC Leu 660	ACT Thr 660	GTC Val 660	CAC His 660	AAA Lys 660	2022
GGC Gly 665	TCA Ser 665	CCT Pro 665	GAC Asp 665	CTG Leu 665	TTA Leu 670	GGT Gly 670	AAT Asn 670	GTC Val 670	CAA Gln 670	GGC Gly 675	TCT Ser 675	GTC Val 675	AGG Arg 675	TAT Tyr 675	GAT Asp 675	2070
CTG Leu 680	GCG Ala 680	TTA Leu 680	GAT Asp 680	CCG Pro 685	GGC Gly 685	CGC Arg 685	CTG Leu 685	ATT Ile 685	TCT Ser 690	CGT Arg 690	GCC Ala 690	ATT Ile 690	TTT Phe 690	GAT Asp 690	GAG Glu 690	2118
ACT Thr 695	AAG Lys 695	AAC Asn 695	TGC Cys 695	ACT Thr 700	TTG Leu 700	ACG Thr 700	GGA Gly 700	AGG Arg 705	AAG Lys 705	ACT Thr 705	CTG Leu 705	GGG Gly 705	CTT Leu 705	GGT Gly 705	GAT Asp 705	2166



- 141 -

CAC His 710	TGC Cys 710	GAA Glu 710	ACA Thr 710	GTG Val 710	AAG Lys 715	CTG Leu 715	CTT Leu 715	TTG Leu 715	CCG Pro 715	GAC Asp 720	TGT Cys 720	GTG Val 720	GAA Glu 720	GAT Asp 725	GCA Ala 725	2214
GTG Val 730	AGC Ser 730	CCT Pro 730	ATC Ile 730	ATC Ile 730	CTG Leu 730	CGC Arg 730	CTC Leu 730	AAC Asn 735	TTT Phe 735	TCC Ser 735	CTG Leu 735	GTG Val 735	AGA Arg 740	GAC Asp 740	TCT Ser 740	2262
GCT Ala 745	TCA Ser 745	CCC Pro 745	AGG Arg 745	AAC Asn 745	CTG Leu 745	CAT His 745	CCT Pro 745	GTG Val 750	CTG Leu 750	GCT Ala 750	GTG Val 750	GGC Gly 755	TCA Ser 755	CAA Gln 755	GAC Asp 755	2310
CAC His 760	ATA Ile 760	ACT Thr 760	GCT Ala 760	TCT Ser 760	CTG Leu 760	CCG Pro 765	TTT Phe 765	GAG Glu 765	AAG Lys 765	AAC Asn 765	TGT Cys 770	AAG Lys 770	CAA Gln 770	GAA Glu 770	CTC Leu 770	2358
CTG Leu 775	TGT Cys 775	GAG Glu 775	GGG Gly 775	GAC Asp 775	CTG Leu 780	GGC Gly 780	ATC Ile 780	AGC Ser 780	TTT Phe 785	AAC Asn 785	TTC Ser 785	TCA Ser 785	GGC Gly 785	CTG Leu 785	CAG Gln 785	2406
GTC Val 790	TTG Leu 790	GTG Val 790	GTG Val 790	GGA Gly 795	GGC Gly 795	TCC Ser 795	CCA Pro 795	GAG Glu 795	CTC Leu 800	ACT Thr 800	GTG Val 800	ACA Thr 800	GTC Val 805	ACT Thr 805	GTG Val 805	2454
TGG Trp 810	AAT Asn 810	GAG Glu 810	GGT Gly 810	GAG Asp 810	GAC Asp 810	AGC Ser 810	TAT Tyr 810	GGA Gly 815	ACT Thr 815	TTA Leu 815	GTC Val 815	AAG Lys 815	TTC Phe 820	TAC Tyr 820	TAC Tyr 820	2502
CCA Pro 825	GCA Ala 825	GGG Gly 825	CTA Leu 825	TCT Ser 825	TAC Tyr 825	CGA Arg 830	CGG Arg 830	GTA Val 830	ACA Thr 830	GGG Gly 835	ACT Thr 835	CAG Gln 835	CAA Gln 835	CCT Pro 835	CAT His 835	2550
CAG Gln 840	TAC Tyr 840	CCA Pro 840	CTA Leu 840	CGC Arg 840	TTG Leu 840	GCC Ala 845	TGT Cys 845	GAG Glu 845	GCT Ala 845	GAG Glu 845	CCC Pro 850	GCT Ala 850	GCC Ala 850	CAG Gln 850	GAG Glu 850	2598
GAC Asp 855	CTG Leu 855	AGG Arg 855	AGC Ser 855	AGC Ser 855	AGC Ser 855	TGT Cys 860	AGC Ser 860	ATT Ile 860	AAT Asn 860	CAC His 865	CCC Pro 865	ATC Ile 865	TTC Phe 865	CGA Arg 865	GAA Glu 865	2646
GGT Gly 870	GCA Ala 870	AAG Lys 870	ACC Thr 870	ACC Thr 870	Phe Met 875	ATG Met 875	ATC Ile 875	ACA Thr 875	TTC Phe 880	GAT Asp 880	GTC Val 880	TCC Ser 880	TAC Tyr 880	AAG Lys 885	GCC Ala 885	2694
TTC Phe 890	CTA Leu 890	GGA Gly 890	GAC Asp 890	AGG Arg 890	TTG Leu 890	CTT Leu 890	CTG Leu 890	AGG Arg 895	GCC Ala 895	AAA Lys 895	GCC Ala 895	AGC Ser 895	AGT Ser 895	GAG Glu 900	AAT Asn 900	2742
AAT Asn 905	AAG Lys 905	CCT Pro 905	GAT Asp 905	ACC Thr 905	AAC Asn 905	AAG Lys 905	ACT Thr 910	GCC Ala 910	TTC Phe 910	CAG Gln 915	CTG Leu 915	GAG Glu 915	CTC Leu 915	CCA Pro 915	GTG Val 915	2790
AAG Lys 920	TAC Tyr 920	ACC Thr 920	GTC Val 920	TAT Tyr 920	ACC Thr 920	CTG Leu 925	ATC Ile 925	AGT Ser 925	AGG Arg 925	CAA Gln 930	GAA Glu 930	GAT Asp 930	TCC Ser 930	ACC Thr 930	AAC Asn 930	2838
CAT His 935	GTC Val 935	AAC Asn 935	TTT Phe 935	TCA Ser 935	TCT Ser 935	TCC Ser 940	CAC His 940	GGG Gly 940	GGG Gly 940	AGA Arg 945	AGG Arg 945	CAA Gln 945	GAA Glu 945	GCC Ala 945	GCA Ala 945	2886

- 142 -

CAT CGC TAT CGT GTG AAT AAC CTG AGT CCA CTG AAG CTG GCC GTC AGA His Arg Tyr Arg Val Asn Asn Leu Ser Pro Leu Lys Leu Ala Val Arg 950 955 960 965	2934
GTT AAC TTC TGG GTC CCT GTC CTT CTG AAC GGT GTG GCT GTG TGG GAC Val Asn Phe Trp Val Pro Val Leu Leu Asn Gly Val Ala Val Trp Asp 970 975 980	2982
GTG ACT CTG AGC AGC CCA GCA CAG GGT GTC TCC TGC GTG TCC CAG ATG Val Thr Leu Ser Ser Pro Ala Gln Gly Val Ser Cys Val Ser Gln Met 985 990 995	3030
AAA CCT CCT CAG AAT CCC GAC TTT CTG ACC CAG ATT CAG AGA CGT TCT Lys Pro Pro Gln Asn Pro Asp Phe Leu Thr Gln Ile Gln Arg Arg Ser 1000 1005 1010	3078
GTG CTG GAC TGC TCC ATT GCT GAC TGC CTG CAC TTC CGC TGT GAC ATC Val Leu Asp Cys Ser Ile Ala Asp Cys Leu His Phe Arg Cys Asp Ile 1015 1020 1025	3126
CCC TCC TTG GAC ATC CAG GAT GAA CTT GAC TTC ATT CTG AGG GGC AAC Pro Ser Leu Asp Ile Gln Asp Glu Leu Asp Phe Ile Leu Arg Gly Asn 1030 1035 1040 1045	3174
CTC AGC TTC GGC TGG GTC AGT CAG ACA TTG CAG GAA AAG GTG TTG CTT Leu Ser Phe Gly Trp Val Ser Gln Thr Leu Gln Glu Lys Val Leu Leu 1050 1055 1060	3222
GTG AGT GAG GCT GAA ATC ACT TTC GAC ACA TCT GTG TAC TCC CAG CTG Val Ser Glu Ala Glu Ile Thr Phe Asp Thr Ser Val Tyr Ser Gln Leu 1065 1070 1075	3270
CCA GGA CAG GAG GCA TTT CTG AGA GCC CAG GTG GAG ACA ACG TTA GAA Pro Gly Gln Glu Ala Phe Leu Arg Ala Gln Val Glu Thr Thr Leu Glu 1080 1085 1090	3318
GAA TAC GTG GTC TAT GAG CCC ATC TTC CTC GTG GCG GGC AGC TCG GTG Glu Tyr Val Val Tyr Glu Pro Ile Phe Leu Val Ala Gly Ser Ser Val 1095 1100 1105	3366
GGA GGT CTG CTG TTA CTG GCT CTC ATC ACA GTG GTA CTG TAC AAG CTT Gly Gly Leu Leu Leu Leu Ala Leu Ile Thr Val Val Leu Tyr Lys Leu 1110 1115 1120 1125	3414
GGC TTC TYC AAA CGT CAG TAC AAA GAA ATG CTG GAC GGC AAG GCT GCA Gly Phe Xaa Lys Arg Gln Tyr Lys Glu Met Leu Asp Gly Lys Ala Ala 1130 1135 1140	3462
GAT CCT GTC ACA GCC GGC CAG GCA GAT TTC GGC TGT GAG ACT CCT CCA Asp Pro Val Thr Ala Gly Gln Ala Asp Phe Gly Cys Glu Thr Pro Pro 1145 1150 1155	3510
TAT CTC GTG AGC TAGGAATCCA CTCTCTGCC TATCTCTGCA ATGAAGATTG Tyr Leu Val Ser 1160	3562
GTCCTGCCTA TGAGTCTACT GGCATGGGAA CGAGT	3597

- 143 -

## (2) INFORMATION FOR SEQ ID NO:55:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1161 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:55:

```

Met Ala Gly Gly Val Val Ile Leu Leu Cys Gly Trp Val Leu Ala Ser
 1             5             10             15
Cys His Gly Ser Asn Leu Asp Val Glu Glu Pro Ile Val Phe Arg Glu
 20             25             30
Asp Ala Ala Ser Phe Gly Gln Thr Val Val Gln Phe Gly Gly Ser Arg
 35             40             45
Leu Val Val Gly Ala Pro Leu Glu Ala Val Ala Val Asn Gln Thr Gly
 50             55             60
Arg Leu Tyr Asp Cys Ala Pro Ala Thr Gly Met Cys Gln Pro Ile Val
 65             70             75             80
Leu Arg Ser Pro Leu Glu Ala Val Asn Met Ser Leu Gly Leu Ser Leu
 85             90             95
Val Thr Ala Thr Asn Asn Ala Gln Leu Leu Ala Cys Gly Pro Thr Ala
100            105            110
Gln Arg Ala Cys Val Lys Asn Met Tyr Ala Lys Gly Ser Cys Leu Leu
115            120            125
Leu Gly Ser Ser Leu Gln Phe Ile Gln Ala Val Pro Ala Ser Met Pro
130            135            140
Glu Cys Pro Arg Gln Glu Met Asp Ile Ala Phe Leu Ile Asp Gly Ser
145            150            155            160
Gly Ser Ile Asn Gln Arg Asp Phe Ala Gln Met Lys Asp Phe Val Lys
165            170            175
Ala Leu Met Gly Glu Phe Ala Ser Thr Ser Thr Leu Phe Ser Leu Met
180            185            190
Gln Tyr Ser Asn Ile Leu Lys Thr His Phe Thr Phe Thr Glu Phe Lys
195            200            205
Asn Ile Leu Asp Pro Gln Ser Leu Val Asp Pro Ile Val Gln Leu Gln
210            215            220
Gly Leu Thr Tyr Thr Ala Thr Gly Ile Arg Thr Val Met Glu Glu Leu
225            230            235            240
Phe His Ser Lys Asn Gly Ser Arg Lys Ser Ala Lys Lys Ile Leu Leu
245            250            255
Val Ile Thr Asp Gly Gln Lys Tyr Arg Asp Pro Leu Glu Tyr Ser Asp
260            265            270

```

- 144 -

Val Ile Pro Ala Ala Asp Lys Ala Gly Ile Ile Arg Tyr Ala Ile Gly  
 275 280 285  
 Val Gly Asp Ala Phe Gln Glu Pro Thr Ala Leu Lys Glu Leu Asn Thr  
 290 295 300  
 Ile Gly Ser Ala Pro Pro Gln Asp His Val Phe Lys Val Gly Asn Phe  
 305 310 315 320  
 Ala Ala Leu Arg Ser Ile Gln Arg Gln Leu Gln Glu Lys Ile Phe Ala  
 325 330 335  
 Ile Glu Gly Thr Gln Ser Arg Ser Ser Ser Ser Phe Gln His Glu Met  
 340 345 350  
 Ser Gln Glu Gly Phe Ser Ser Ala Leu Thr Ser Asp Gly Pro Val Leu  
 355 360 365  
 Gly Ala Val Gly Ser Phe Ser Trp Ser Gly Gly Ala Phe Leu Tyr Pro  
 370 375 380  
 Pro Asn Thr Arg Pro Thr Phe Ile Asn Met Ser Gln Glu Asn Val Asp  
 385 390 395 400  
 Met Arg Asp Ser Tyr Leu Gly Tyr Ser Thr Ala Val Ala Phe Trp Lys  
 405 410 415  
 Gly Val His Ser Leu Ile Leu Gly Ala Pro Arg His Gln His Thr Gly  
 420 425 430  
 Lys Val Val Ile Phe Thr Gln Glu Ala Arg His Trp Arg Pro Lys Ser  
 435 440 445  
 Glu Val Arg Gly Thr Gln Ile Gly Ser Tyr Phe Gly Ala Ser Leu Cys  
 450 455 460  
 Ser Val Asp Val Asp Arg Asp Gly Ser Xaa Asp Leu Val Leu Ile Gly  
 465 470 475 480  
 Ala Pro His Tyr Tyr Glu Gln Thr Arg Gly Gly Gln Val Ser Val Phe  
 485 490 495  
 Pro Val Pro Gly Val Arg Gly Arg Trp Gln Cys Glu Ala Thr Leu His  
 500 505 510  
 Gly Glu Gln Gly His Pro Trp Gly Arg Phe Gly Val Ala Leu Thr Val  
 515 520 525  
 Leu Gly Asp Val Asn Gly Asp Asn Leu Ala Asp Val Ala Ile Gly Ala  
 530 535 540  
 Pro Gly Glu Glu Glu Ser Arg Gly Ala Val Tyr Ile Phe His Gly Ala  
 545 550 555 560  
 Ser Arg Leu Glu Ile Met Pro Ser Pro Ser Gln Arg Val Thr Gly Ser  
 565 570 575  
 Gln Leu Ser Leu Arg Leu Gln Tyr Phe Gly Gln Ser Leu Ser Gly Gly  
 580 585 590

- 145 -

Gln Asp Leu Thr Gln Asp Gly Leu Val Asp Leu Ala Val Gly Ala Gln  
 595 600 605  
 Gly His Val Leu Leu Leu Arg Ser Leu Pro Leu Leu Lys Val Glu Leu  
 610 615 620  
 Ser Ile Arg Phe Ala Pro Met Glu Val Ala Lys Ala Val Tyr Gln Cys  
 625 630 635 640  
 Trp Glu Arg Thr Pro Thr Val Leu Glu Ala Gly Glu Ala Thr Val Cys  
 645 650 655  
 Leu Thr Val His Lys Gly Ser Pro Asp Leu Leu Gly Asn Val Gln Gly  
 660 665 670  
 Ser Val Arg Tyr Asp Leu Ala Leu Asp Pro Gly Arg Leu Ile Ser Arg  
 675 680 685  
 Ala Ile Phe Asp Glu Thr Lys Asn Cys Thr Leu Thr Gly Arg Lys Thr  
 690 695 700  
 Leu Gly Leu Gly Asp His Cys Glu Thr Val Lys Leu Leu Leu Pro Asp  
 705 710 715 720  
 Cys Val Glu Asp Ala Val Ser Pro Ile Ile Leu Arg Leu Asn Phe Ser  
 725 730 735  
 Leu Val Arg Asp Ser Ala Ser Pro Arg Asn Leu His Pro Val Leu Ala  
 740 745 750  
 Val Gly Ser Gln Asp His Ile Thr Ala Ser Leu Pro Phe Glu Lys Asn  
 755 760 765  
 Cys Lys Gln Glu Leu Leu Cys Glu Gly Asp Leu Gly Ile Ser Phe Asn  
 770 775 780  
 Phe Ser Gly Leu Gln Val Leu Val Val Gly Gly Ser Pro Glu Leu Thr  
 785 790 795 800  
 Val Thr Val Thr Val Trp Asn Glu Gly Glu Asp Ser Tyr Gly Thr Leu  
 805 810 815  
 Val Lys Phe Tyr Tyr Pro Ala Gly Leu Ser Tyr Arg Arg Val Thr Gly  
 820 825 830  
 Thr Gln Gln Pro His Gln Tyr Pro Leu Arg Leu Ala Cys Glu Ala Glu  
 835 840 845  
 Pro Ala Ala Gln Glu Asp Leu Arg Ser Ser Ser Cys Ser Ile Asn His  
 850 855 860  
 Pro Ile Phe Arg Glu Gly Ala Lys Thr Thr Phe Met Ile Thr Phe Asp  
 865 870 875 880  
 Val Ser Tyr Lys Ala Phe Leu Gly Asp Arg Leu Leu Leu Arg Ala Lys  
 885 890 895  
 Ala Ser Ser Glu Asn Asn Lys Pro Asp Thr Asn Lys Thr Ala Phe Gln  
 900 905 910

- 146 -

Leu Glu Leu Pro Val Lys Tyr Thr Val Tyr Thr Leu Ile Ser Arg Gln  
 915 920 925  
 Glu Asp Ser Thr Asn His Val Asn Phe Ser Ser Ser His Gly Gly Arg  
 930 935 940  
 Arg Gln Glu Ala Ala His Arg Tyr Arg Val Asn Asn Leu Ser Pro Leu  
 945 950 955 960  
 Lys Leu Ala Val Arg Val Asn Phe Trp Val Pro Val Leu Leu Asn Gly  
 965 970 975  
 Val Ala Val Trp Asp Val Thr Leu Ser Ser Pro Ala Gln Gly Val Ser  
 980 985 990  
 Cys Val Ser Gln Met Lys Pro Pro Gln Asn Pro Asp Phe Leu Thr Gln  
 995 1000 1005  
 Ile Gln Arg Arg Ser Val Leu Asp Cys Ser Ile Ala Asp Cys Leu His  
 1010 1015 1020  
 Phe Arg Cys Asp Ile Pro Ser Leu Asp Ile Gln Asp Glu Leu Asp Phe  
 1025 1030 1035 1040  
 Ile Leu Arg Gly Asn Leu Ser Phe Gly Trp Val Ser Gln Thr Leu Gln  
 1045 1050 1055  
 Glu Lys Val Leu Leu Val Ser Glu Ala Glu Ile Thr Phe Asp Thr Ser  
 1060 1065 1070  
 Val Tyr Ser Gln Leu Pro Gly Gln Glu Ala Phe Leu Arg Ala Gln Val  
 1075 1080 1085  
 Glu Thr Thr Leu Glu Glu Tyr Val Val Tyr Glu Pro Ile Phe Leu Val  
 1090 1095 1100  
 Ala Gly Ser Ser Val Gly Gly Leu Leu Leu Leu Ala Leu Ile Thr Val  
 1105 1110 1115 1120  
 Val Leu Tyr Lys Leu Gly Xaa Xaa Lys Arg Gln Tyr Lys Glu Met Leu  
 1125 1130 1135  
 Asp Gly Lys Ala Ala Asp Pro Val Thr Xaa Gly Gln Ala Asp Phe Gly  
 1140 1145 1150  
 Cys Glu Thr Pro Pro Tyr Leu Val Ser  
 1155 1160

## (2) INFORMATION FOR SEQ ID NO:56:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 20 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

- 147 -

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:56:

CCTGTCATGG GTCTAACCTG

20

(2) INFORMATION FOR SEQ ID NO:57:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 19 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:57:

AGGTTAGACC CATGACAGG

19

(2) INFORMATION FOR SEQ ID NO:58:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 20 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:58:

GGCCTTGCA G CTGGACAATG

20

(2) INFORMATION FOR SEQ ID NO:59:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 22 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:59:

CCAAAGCTGG CTGCATCCTC TC

22

(2) INFORMATION FOR SEQ ID NO:60:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 21 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

- 148 -

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:60:

CCGCCTGCCA CTGGCGTGTG C

21

(2) INFORMATION FOR SEQ ID NO:61:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 22 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:61:

CCCAGATGAA GGACTTCGTC AA

22

(2) INFORMATION FOR SEQ ID NO:62:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 20 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:62:

GCTGGGATCA TTCGCTATGC

20

(2) INFORMATION FOR SEQ ID NO:63:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 21 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:63:

CAATGGATGG ACCAGTTCTG G

21

(2) INFORMATION FOR SEQ ID NO:64:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 20 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear



- 149 -

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:64:

CAGATCGGCT CCTACTTTGG

20

(2) INFORMATION FOR SEQ ID NO:65:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 19 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:65:

CATGGAGCCT CGAGACAGG

19

(2) INFORMATION FOR SEQ ID NO:66:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 21 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:66:

CCACTGTCCT CGAAGCTGGA G

21

(2) INFORMATION FOR SEQ ID NO:67:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 26 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:67:

CTTCGTCCTG TGCTGGCTGT GGGCTC

26

(2) INFORMATION FOR SEQ ID NO:68:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 21 base pairs
  - (B) TYPE: nucleic acid

- 150 -

- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:68:

CGCCTGGCAT GTGAGGCTGA G

21

(2) INFORMATION FOR SEQ ID NO:69:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 21 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:69:

CCGTGATCAG TAGGCAGGAA G

21

(2) INFORMATION FOR SEQ ID NO:70:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 18 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:70:

GTCACAGAGG GAACCTCC

18

(2) INFORMATION FOR SEQ ID NO:71:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 23 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:71:

GCTCCTGAGT GAGGCTGAAA TCA

23

(2) INFORMATION FOR SEQ ID NO:72:

- 151 -

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 23 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

- (ii) MOLECULE TYPE: DNA

- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:72:

GAGATGCTGG ATCTACCATC TGC

23

- (2) INFORMATION FOR SEQ ID NO:73:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 22 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

- (ii) MOLECULE TYPE: DNA

- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:73:

CTGAGCTGGG AGATTTTAT GG

22

- (2) INFORMATION FOR SEQ ID NO:74:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 21 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

- (ii) MOLECULE TYPE: DNA

- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:74:

GTGGATCAGC ACTGAAATCT G

21

- (2) INFORMATION FOR SEQ ID NO:75:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 21 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

- (ii) MOLECULE TYPE: DNA

- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:75:

CGTTTGAAGA AGCCAAGCTT G

21

- 152 -

## (2) INFORMATION FOR SEQ ID NO:76:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 20 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:76:

CACAGCGGAG GTGCAGGCAG

20

## (2) INFORMATION FOR SEQ ID NO:77:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 18 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:77:

CTCACTGCTT GCGCTGGC

18

## (2) INFORMATION FOR SEQ ID NO:78:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 20 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:78:

CGGTAAGATA GCTCTGCTGG

20

## (2) INFORMATION FOR SEQ ID NO:79:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 20 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:79:

- 153 -

GAGCCCACAG CCAGCACAGG

20

## (2) INFORMATION FOR SEQ ID NO:80:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 21 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: DNA

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:80:

GATCCAACGC CAGATCATAC C

21

## (2) INFORMATION FOR SEQ ID NO:81:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 20 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: DNA

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:81:

CACGGCCAGG TCCACCAGGC

20

## (2) INFORMATION FOR SEQ ID NO:82:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 21 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: DNA

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:82:

CACGTCCCCT AGCACTGTCA G

21

## (2) INFORMATION FOR SEQ ID NO:83:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 22 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: DNA

- 154 -

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:83:

TTGACGAAGT CCTTCATCTG GG

22

(2) INFORMATION FOR SEQ ID NO:84:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 21 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:84:

GAACTGCAAG CTGGAGCCCA G

21

(2) INFORMATION FOR SEQ ID NO:85:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 21 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:85:

CTGGATGCTG CGAAGTGCTA C

21

(2) INFORMATION FOR SEQ ID NO:86:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 21 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:86:

GCCTTGGAGC TGGACGATGG C

21

(2) INFORMATION FOR SEQ ID NO:87:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 33 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

- 155 -

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:87:

GTAAGATCTC CAGAGTGTCC AAGACAAGAG ATG

33

(2) INFORMATION FOR SEQ ID NO:88:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 33 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:88:

CTTCTCGAGT GTGAGAGCTG AACTGAAACC TTC

33

(2) INFORMATION FOR SEQ ID NO:89:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 32 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:89:

CGCTGTGACG TCAGAGTTGA GTCCAAATAT GG

32

(2) INFORMATION FOR SEQ ID NO:90:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 21 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:90:

GGTGACACTA TAGAATAGGG C

21

(2) INFORMATION FOR SEQ ID NO:91:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 18 base pairs
- (B) TYPE: nucleic acid

- 156 -

(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:91:

AAGCAGGAGCTCCTGTGT

18

(2) INFORMATION FOR SEQ ID NO:92:

(i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 852 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(ix) FEATURE:

(A) NAME/KEY: CDS  
(B) LOCATION: 61..852

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:92:

TGATCTCCCT	CCAGGCCACT	GTTCCCTCTC	CACTTCCCCT	CACCGCTGCA	CTGCTCAGAG	60
ATG GCC CTT GGG GCT GTG GTC CTC CTT GGG GTC CTG GCT TCT TAC CAC	108					
Met Ala Leu Gly Ala Val Val Leu Leu Gly Val Leu Ala Ser Tyr His						
1 5 10 15						
GGA TTC AAC TTG GAC GTG ATG AGC GGT GAT CTT CCA GGA AGA CGC AGC	156					
Gly Phe Asn Leu Asp Val Met Ser Gly Asp Leu Pro Gly Arg Arg Ser						
20 25 30						
GGG CTT CGG GCA GAG CGT GAT GCA GTT TGG GGA TCT CGA CTC GTG GTG	204					
Gly Leu Arg Ala Glu Arg Asp Ala Val Trp Gly Ser Arg Leu Val Val						
35 40 45						
GGA GCC CCC CTG GCG GTG GTG TCG GCC AAC CAC ACA GGA CGG CTG TAC	252					
Gly Ala Pro Leu Ala Val Val Ser Ala Asn His Thr Gly Arg Leu Tyr						
50 55 60						
GAG TGT GCG CCT GCC TCC GGC ACC TGC ACG CCC ATT TTC CCA TTC ATG	300					
Glu Cys Ala Pro Ala Ser Gly Thr Cys Thr Pro Ile Phe Pro Phe Met						
65 70 75 80						
CCC CCC GAA GCC GTG AAC ATG TCC CTG GGC CTG TCC CTG GCA GCC TCC	348					
Pro Pro Glu Ala Val Asn Met Ser Leu Gly Leu Ser Leu Ala Ala Ser						
85 90 95						
CCC AAC CAT TCC CAG CTG CTG GCT TGT GGC CCG ACC GTG CAT AGA GCC	396					
Pro Asn His Ser Gln Leu Leu Ala Cys Gly Pro Thr Val His Arg Ala						
100 105 110						



- 157 -

TGC GGG GAG GAC GTG TAC GCC CAG GGT TTC TGT GTG CTG CTG GAT GCC	444
Cys Gly Glu Asp Val Tyr Ala Gln Gly Phe Cys Val Leu Leu Asp Ala	
115 120 125	
CAC GCA CAG CCC ATC GGG ACT GTG CCA GCT GCC CTG CCC GAG TGC CCA	492
His Ala Gln Pro Ile Gly Thr Val Pro Ala Ala Leu Pro Glu Cys Pro	
130 135 140	
GAT CAA GAG ATG GAC ATT GTC TTC CTG ATT GAC GGC TCT GGC AGC ATT	540
Asp Gln Glu Met Asp Ile Val Phe Leu Ile Asp Gly Ser Gly Ser Ile	
145 150 155 160	
AGC TCA AAT GAC TTC CGC AAG ATG AAG GAC TTT GTC AGA GCT GTG ATG	588
Ser Ser Asn Asp Phe Arg Lys Met Lys Asp Phe Val Arg Ala Val Met	
165 170 175	
GAC CAG TTC AAG GAC ACC AAC ACC CAG TTC TCG CTG ATG CAG TAC TCC	636
Asp Gln Phe Lys Asp Thr Asn Thr Gln Phe Ser Leu Met Gln Tyr Ser	
180 185 190	
AAT GTG CTG GTG ACA CAT TTC ACC TTC AGC AGC TTC CGG AAC AGC TCC	684
Asn Val Leu Val Thr His Phe Thr Phe Ser Ser Phe Arg Asn Ser Ser	
195 200 205	
AAT CCT CAG GCC CTA GTG GAG CCC ATT GTG CAG CTG ACA GGC CTC ACG	732
Asn Pro Gln Gly Leu Val Glu Pro Ile Val Gln Leu Thr Gly Leu Thr	
210 215 220	
TTC ACG GCC ACA GGG ATC CTG AAA GTG GTG ACA GAG CTG TTT CAA ACC	780
Phe Thr Ala Thr Gly Ile Leu Lys Val Val Thr Glu Leu Phe Gln Thr	
225 230 235 240	
AAG AAC GGG GCC CGC GAA AGT GCC AAG AAG ATC CTC ATC GTC ATC ACA	828
Lys Asn Gly Ala Arg Glu Ser Ala Lys Lys Ile Leu Ile Val Ile Thr	
245 250 255	
GAT GGG CAG AAG TAC AAA GCG GCA	852
Asp Gly Gln Lys Tyr Lys Ala Ala	
260	

## (2) INFORMATION FOR SEQ ID NO:93:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 264 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:93:

Met	Ala	Leu	Gly	Ala	Val	Val	Leu	Leu	Gly	Val	Leu	Ala	Ser	Tyr	His
1				5					10					15	
Gly	Phe	Asn	Leu	Asp	Val	Met	Ser	Gly	Asp	Leu	Pro	Gly	Arg	Arg	Ser
		20						25					30		
Gly	Leu	Arg	Ala	Glu	Arg	Asp	Ala	Val	Trp	Gly	Ser	Arg	Leu	Val	Val
		35					40					45			

Gly 50	Ala	Pro	Leu	Ala	Val 55	Ser	Ala	Asn	His	Thr 60	Gly	Arg	Leu	Tyr	
Glu 65	Cys	Ala	Pro	Ala	Ser 70	Gly	Thr	Cys	Thr	Pro 75	Ile	Phe	Pro	Phe 80	
Pro	Pro	Glu	Ala	Val 85	Asn	Met	Ser	Leu	Gly 90	Leu	Ser	Leu	Ala	Ala 95	
Pro	Asn	His	Ser 100	Gln	Leu	Leu	Ala	Cys 105	Gly	Pro	Thr	Val	His 110	Arg	Ala
Cys	Gly	Glu 115	Asp	Val	Tyr	Ala	Gln 120	Gly	Phe	Cys	Val	Leu 125	Leu	Asp	Ala
His 130	Ala	Gln	Pro	Ile	Gly	Thr 135	Val	Pro	Ala	Ala	Leu 140	Pro	Glu	Cys	Pro
Asp 145	Gln	Glu	Met	Asp	Ile 150	Val	Phe	Leu	Ile	Asp 155	Gly	Ser	Gly	Ser	Ile 160
Ser	Ser	Asn	Asp	Phe 165	Arg	Lys	Met	Lys	Asp 170	Phe	Val	Arg	Ala	Val 175	Met
Asp	Gln	Phe 180	Lys	Asp	Thr	Asn	Thr	Gln 185	Phe	Ser	Leu	Met	Gln 190	Tyr	Ser
Asn	Val	Leu 195	Val	Thr	His	Phe	Thr 200	Phe	Ser	Ser	Phe	Arg 205	Asn	Ser	Ser
Asn	Pro 210	Gln	Gly	Leu	Val	Glu 215	Pro	Ile	Val	Gln	Leu 220	Thr	Gly	Leu	Thr
Phe 225	Thr	Ala	Thr	Gly	Ile 230	Leu	Lys	Val	Val	Thr 235	Glu	Leu	Phe	Gln	Thr 240
Lys	Asn	Gly	Ala	Arg 245	Glu	Ser	Ala	Lys	Lys 250	Ile	Leu	Ile	Val	Ile 255	Thr
Asp	Gly	Gln 260	Lys	Tyr	Lys	Ala	Ala								

- 159 -

WHAT IS CLAIMED IS:

1. A purified and isolated  $\alpha_d$  polynucleotide consisting essentially of human  $\alpha_d$  protein coding sequence set out in SEQ ID NO: 1.
2. The polynucleotide of claim 1 which is a DNA molecule.
3. The DNA molecule of claim 2 which is a cDNA molecule.
4. The DNA molecule of claim 2 which is a genomic DNA molecule.
5. The DNA molecule of claim 2 which is a wholly or partially chemically synthesized DNA molecule.
6. A full length purified and isolated  $\alpha_d$ -encoding polynucleotide selected from the group consisting of:
  - a) the human DNA sequence set out in SEQ ID NO: 1, and
  - b) a DNA molecule which hybridizes under stringent conditions to the noncoding strand of the protein coding portion of the DNA of a).
7. A DNA molecule encoding the human  $\alpha_d$  amino acid sequence set out in SEQ ID NO: 2.
8. A DNA expression construct comprising a DNA molecule according to claim 2.
9. A host cell transformed with a DNA molecule according to claim 2.

- 160 -

10. A method for producing an  $\alpha_d$  polypeptide comprising growing a host cell according to claim 9 in a suitable medium and isolating  $\alpha_d$  polypeptide from said host cell or the medium of its growth.
11. Purified and isolated  $\alpha_d$  polypeptide consisting essentially of the human  $\alpha_d$  amino acid sequence set out in SEQ ID NO: 2.
12. A polypeptide capable of specifically binding to  $\alpha_d$ .
13. A polypeptide according to claim 12 which is an antibody.
14. An antibody according to claim 13 which is a monoclonal antibody.
15. An anti-idiotypic antibody specific for the monoclonal antibody of claim 14.
16. A hybridoma cell line producing the monoclonal antibody according to claim 14.
17. A purified and isolated  $\alpha_d$  extracellular domain polypeptide fragment comprising amino acids 17 to 1108 of the human  $\alpha_d$  amino acid sequence set out in SEQ ID NO: 2.
18. A purified and isolated  $\alpha_d$  I domain polypeptide fragment comprising amino acids 145 to 355 of the human  $\alpha_d$  amino acid sequence set out in SEQ ID NO: 2.

- 161 -

19. A fusion protein comprising  $\alpha_d$  extracellular domain polypeptide amino acids 17 to 1108 of SEQ ID NO: 2 and human immunoglobulin constant domain sequences.

20. A purified and isolated murine polynucleotide consisting essentially of the  $\alpha$  subunit protein coding sequence set out in SEQ ID NO: 45.

21. A method for isolating a polynucleotide encoding a protein that binds to  $\alpha_d$  comprising the steps of:

a) transforming or transfecting appropriate host cells with a DNA construct comprising a reporter gene under the control of a promoter regulated by a transcription factor having a DNA-binding domain and an activating domain;

b) expressing in said host cells a first hybrid DNA sequence encoding a first fusion of part or all of  $\alpha_d$  and either the DNA binding domain or the activating domain of said transcription factor;

c) expressing in said host cells a library of second hybrid DNA sequences encoding second fusions of part or all of putative  $\alpha_d$  binding proteins and the DNA binding domain or activating domain of said transcription factor which is not incorporated in said first fusion;

d) detecting binding of an  $\alpha_d$  binding protein to  $\alpha_d$  in a particular host cell by detecting the production of reporter gene product in said host cell; and

e) isolating second hybrid DNA sequences encoding  $\alpha_d$  binding protein from said particular host cell.

22. A method for identifying a compound capable of reacting specifically with  $\alpha_d$  and of modulating the interaction of binding partners  $\alpha_d$  and ICAM-R comprising the steps of:

- 162 -

a) immobilizing  $\alpha_d$  or a fragment thereof, or ICAM-R or a fragment thereof, on a solid support coated or impregnated with a fluorescent agent;

b) labelling the non-immobilized binding partner with a compound capable of exciting said fluorescent agent;

c) contacting said immobilized binding partner with said labelled binding partner in the presence and absence of a putative modulator compound capable of specifically reacting with  $\alpha_d$ ;

d) detecting light emission by said fluorescent agent; and

e) identifying modulating compounds as those compounds that affect the emission of light by said fluorescent agent in comparison to the emission of light by said fluorescent agent in the absence of said modulating compound.

23. A purified and isolated  $\alpha_d$  extracellular domain polypeptide fragment comprising about amino acid 127 to about amino acid 353 of the human  $\alpha_d$  amino acid sequence set out in SEQ ID NO: 2.

24. A fusion protein comprising the polypeptide fragment of claim 23 and human immunoglobulin constant region sequences.

25. A purified and isolated  $\alpha_d$  extracellular domain polypeptide fragment comprising about amino acid 17 to about amino acid 603 of the human  $\alpha_d$  amino acid sequence set out in SEQ ID NO: 2.

26. A fusion protein comprising the polypeptide fragment of claim 25 and human immunoglobulin constant region sequences.

$\alpha$ D	TF-GT--VLL	LSVLASYHGF	NLDVEEPTIF	QEDAGGFGOS	VVQFGGSRLV	47
CD11B	MA-LR--VLL	LTALTCHGF	NLDTENAMTF	QENARGFGOS	VVQLOGSRVV	47
CD11C	MTRTRAALL	FTALATSLGF	NLDTEELTAF	RVDSAGFGDS	VVQYANSWV	50
$\alpha$ D	VGAPLEVAA	NOTGRLYDCA	AATGMCQIP	LHIRPEAVNM	SLGLTLAAS	97
CD11B	VGAPQEI VAA	NORGSLYQCD	YSTGSCEPIR	LQVPVEAVNM	SLGLSLAATT	97
CD11C	VGAPOKIIAA	NOIGGLYQCG	YSTGACEPIG	LQVPPEAVNM	SLGLSLASTT	100
$\alpha$ D	NGSRLLACGP	TLHRVCGENS	YSKGSCLLLG	SR-WEIIQTV	PDATPECPHQ	146
CD11B	SPPQLLACGP	TVHQTCSNT	YVKGLCFLFG	SNLRQOPQKF	PEALRGCPQE	147
CD11C	SPSOLLACGP	TVHHECGRNM	YLTGLCFLLG	PT--QLTORL	PVSRQECPRQ	148
$\alpha$ D	EMDIVFLIDG	SGSIDQNDEN	OMKGFVQAVM	GQFEGTDTLF	ALMQYSNLLK	196
CD11B	DSDIAFLIDG	SGSIIPHDFR	RMKEFVSTVM	EQLKSKSTLF	SLMQYSEEFR	197
CD11C	EODIVFLIDG	SGSISRNFA	TMMNFVRAVI	SOFORPSTQF	SLMQFSNKFO	198
$\alpha$ D	IHFTFTQFRT	SPSQOQSLVDP	IVQLKGLTFT	ATGILTVVTQ	LFHHKNGARK	246
CD11B	IHFTFKEFON	NPNPRSLVKP	ITQLLGRHT	ATGIRKVVRE	LFNITNGARK	247
CD11C	THFTFEEFRR	TSNPLSLLAS	VHQLQGFTYT	ATAIONVVHR	LFHASYGARR	248
$\alpha$ D	SAKKILIVIT	DGOKYKDPLE	YSDVIPOAEK	AGIIRYAIGV	GHAFOGPTAR	296
CD11B	NAFKILVIT	DGEKFGDPLG	YEDVIPEADR	EGVIRYVIGV	GDAFRSEKSR	297
CD11C	DAIKILIVIT	DGKKEGDSLD	YKDVIPMADA	AGIIRYAIGV	GLAFQNRNSW	298

FIGURE 1A

αD	QELNTISSAP	PODHVFKVDN	FAALGSIQKQ	LOEKIYAVEG	TQSRASSSFQ	346
CD11B	QELNTIASKP	PRDHVFQVNN	FEALKTIONQ	LREKIFAIEG	TQTGSSSSFE	347
CD11C	KELNDIASKP	SQEHIFKVED	FDALKDIONQ	LKEKIFAIEG	TETISSSSFE	348
αD	HEMSQEGFST	ALTMDFGLG	AVGSFSWGG	AFLYPPNMSP	TFINMSQENV	396
CD11B	HEMSQEGFSA	AITSNGPLLS	TVGSYDWAGG	VFLYTSKEKS	TFINMTRVDS	397
CD11C	LEMAQEGFSA	VFTPDGPVLG	AVGSFTWGG	AFLYPPNMSP	TFINMSQENV	398
αD	DMRDSYLGYS	TELALWKGVQ	NVLGAPRYQ	HTGKAVIFTO	VSRQWRKKA	446
CD11B	DMNDAYLGYA	AAIILNRVQ	SLVLGAPRYQ	HIGLVAMFRQ	NTGMMESNAN	447
CD11C	DMRDSYLGYS	TELALWKGVQ	SLVLGAPRYQ	HIGKAVIFIO	VSRQWRMKA	448
αD	VTGTQIGSYF	GASLCSVDVD	SDGSTDLILI	GAPHYEQTR	GGQVSVCPPLP	496
CD11B	VKGTOIGAYF	GASLCSVDVD	SNGSTDLVLI	GAPHYEQTR	GGQVSVCPPLP	497
CD11C	VIGTOIGSYF	GASLCSVDVD	TDGSTDLVLI	GAPHYEQTR	GGQVSVCPPLP	498
αD	RGQRVQWQCD	AVLRGEQGHP	WGRFGAALT	LGDVNEDKLI	DVAIGAPGEE	546
CD11B	RGQRARWQCD	AVLYGEQGQP	WGRFGAALT	LGDVNGDKLT	DVAIGAPGEE	547
CD11C	RGWRRW-CD	AVLYGEQGHP	WGRFGAALT	LGDVNGDKLT	DVVIGAPGEE	547
αD	ENRGAVYLFH	GASESGISPS	HSQRIASSQL	SPRLQYFGQA	LSGGODLTOD	596
CD11B	DNRGAVYLFH	GTSGSGISPS	HSQRIAGSKL	SPRLQYFGQS	LSGGODLTMD	597
CD11C	ENRGAVYLFH	GVLGPSISPS	HSQRIAGSQL	SSRLQYFGQA	LSGGODLTOD	597

FIGURE 1B



- 163 -

27. A purified and isolated  $\alpha_d$  extracellular domain polypeptide fragment comprising about amino acid 17 to about amino acid 1029 of the human  $\alpha_d$  amino acid sequence set out in SEQ ID NO: 2.

28. A fusion protein comprising the polypeptide fragment of claim 27 and immunoglobulin constant region sequences.

29. A purified and isolated murine polynucleotide comprising the  $\alpha$  subunit protein coding sequence as set out in SEQ ID NO: 52.

30. A purified and isolated  $\alpha_d$  polypeptide consisting essentially of the murine  $\alpha_d$  amino acid sequence set out in SEQ ID NO: 53.

31. A purified and isolated rat polynucleotide comprising the  $\alpha$  subunit protein coding sequence as set out in SEQ ID NO: 54.

32. A purified and isolated  $\alpha_d$  polypeptide consisting essentially of the rat  $\alpha_d$  amino acid sequence set out in SEQ ID NO: 55.

33. A purified and isolated polypeptide fragment comprising extracellular domain sequences of the polypeptide of claim 32.

34. A polypeptide capable of specifically binding to the polypeptide of claim 32.

35. A polypeptide according to claim 34 which is an antibody.

36. An antibody according to claim 35 which is a monoclonal antibody.

- 164 -

37. A rodent that does not express a functional  $\alpha_d$  protein.

38. A rodent that expresses a variant  $\alpha_d$  protein.

αD	GLMDLAVGAR	GOVLLLRSLP	VLKVGAMRF	SPVEVAKAVY	RCWEEKPSAL	646
CD11B	GLVDLTVGAQ	GHVLLLR SOP	VLRVKAIM EF	NPREVARNVF	ECNDQVVK GK	647
CD11C	GLVDLAVGAR	GOVLLLRTRP	VLWVGSMQF	IPAEIPRS AF	ECREQVVS EQ	647
αD	EAGDATVCLT	IOKSSLDQL-	-GDIOSSVF	DLALDPGR LT	SRAIFNETKN	694
CD11B	EAGEVRVCLH	VOKSTRDR LR	EGQIOSVVTY	DLALDSGR PH	SRAVFN ETKN	697
CD11C	TLVQSNICLY	IDKRSKNLLG	SRDLOSSVTL	DLALAPGR LS	PRAIFQETKN	697
αD	PTLTTRKTLG	LGIHCETLKL	LLPDCVEDV V	SPIILHLNFS	LVREPIPS PQ	744
CD11B	STRRQTQVLG	LTQT CETLKL	QLPNCIEDPV	SPIVLR LNFS	LVGTPLSAFG	747
CD11C	RSLSRVRVLG	LKAHCENFNL	LLPSCVEDSV	IPIILRLNFT	LVGKPLLA FR	747
αD	NLRPVLAVGS	ODLFTASLPF	EKNCGODGLC	EGDLGVTL SF	SGLQTLTVGS	794
CD11B	NLRPVLAE DA	ORLFTALFPF	EKNCGNDNIC	ODDLSITFSF	MSLDCLVVG G	797
CD11C	NLRPMLAALA	ORYFTASLPF	EKNCGADHIC	ODNLGISFSF	PGLKSLLVGS	797
αD	SLELNIVITV	WNAGEDSYGT	VVSLYYPAGL	SHRRVSGAOK	OPHQSALRLA	844
CD11B	PREFNVTVTV	RNDGEDSYRT	QVTFFFPLDL	SYRKVSTLON	QRSORSWR LA	847
CD11C	NLELNAEVMV	WNDGEDSYGT	TITFSHPAGL	SYRYVAEGOK	QGQLRSLH LT	847
αD	CETVPTD--	EGLRSSRC SV	NHPIFHEGSN	GTFFIVTFDVS	Y-----KATLG	888
CD11B	CESASSTEVS	GALKSTSCSI	NHPIFPENSE	-----VTFNIT	FDVDSKASLG	893
CD11C	CCSA-PVGSQ	GTW-STSCRI	NHLIFRGGAQ	-----ITFLAT	FDVSPKAVGL	891

FIGURE 1C

αD	DRMLRASAS	SENNKASSK	ATFOLELPVK	YAVYTMISRQ	EESTKYFNFA	938
CD11B	NKLLLLKANVT	SENNMPRTNK	TEFOLELPVK	YAVYMVVTSH	GVSTKYLNFT	943
CD11C	DRLLLIANVS	SENNIPRTSK	TIFOLELPVK	YAVYIVVSSH	EQFTKYLNFS	941
αD	TS-DEKMKKE	AEHRYRVNNL	SORDLAISIN	FWVPVLLNGV	AVWDVVMEAP	987
CD11B	AS-ENTS-RV	MOHQYQVSNL	GORSPLISLV	FLVPVRLNQT	VIWDRPQVTF	991
CD11C	ESEEKES-HV	AMHRYQVNNL	GORDLPVSIN	FWVPVELNQE	AVWMDVEVSH	990
αD	SQSLP--CVS	ERKPPQHSDF	LTOISRSPML	DCSIADCLOF	RCDVPSFSVQ	1035
CD11B	SENLSSTCHT	KERLPSSHDF	LAELRKAPVV	NCSIAVCORI	QCDIPFFGIQ	1041
CD11C	PQNPSLRCS	EKIAPPASDF	LAHIQKNPVL	DCSIAGCLR	RCDVPSFSVQ	1040
αD	EELDFTLKGN	LSFGWVRETL	QKKVLVSVVA	EITFDTSVYS	QLPGQEAAMR	1085
CD11B	EEFNATLKGN	LSFDWYIKTS	HNHLLIVSTA	EILFNDSVFT	LLPGOGAFVR	1091
CD11C	EELDFTLKGN	LSFGWVRQIL	QKKVSVSVVA	EIIFDTSVYS	QLPGQEAAMR	1090
αD	AOMEMVLEED	EVYNAIPIIM	GSSVGALLLL	ALITATLYKL	GFFKRHYKEM	1135
CD11B	SQTETKVEPF	EVNPNPLPIV	GSSVGGLLLL	ALITAALYKL	GFFKRQYKDM	1141
CD11C	AQTITVLEKY	KVHNPIPLIV	GSSIGGLLLL	ALITAVLYKV	GFFKRQYKEM	1140
αD	LEDKPED---	-----TATFS	GDDFSCVAPN	VPLS		1161
CD11B	M---SEG---	-----GP--P	GAE-----PP	----		1153
CD11C	M---EEANGO	IAPENG--Q	TPS-----PP	SEK		1163

FIGURE 1D

# INTERNATIONAL SEARCH REPORT

In. ational application No.  
PCT/US94/14832

## A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) :Please See Extra Sheet.

US CL :Please See Extra Sheet.

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 435/6, 7.1, 69.1, 240.2, 252.3, 320.1; 530/350, 387.1, 387.2, 388.1, 388.22; 536/22.1, 23.1, 23.5

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

None

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

APS, BIOSIS, CAS ONLINE, STN

search terms: human beta2 integrin, leukointegrin, leu cam, leukocyte integrin#, alphad, integrin, alphas

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	CELL, Volume 72, issued 26 March 1993, Michishita et al., "A Novel Divalent Cation-Binding Site in the A Domain of the $\beta$ 2 Integrin CR3 (CD11b/CD18) Is Essential for Ligand Binding", pages 857-867, see entire document.	1-38
A	CELL, Volume 69, issued 03 April 1992, Hynes, "Integrins: Versatility, Modulation, and Signaling in Cell Adhesion", pages 11-25, see entire document.	1-38

☒ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

* Special categories of cited documents:	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"A" document defining the general state of the art which is not considered to be of particular relevance	"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"E" earlier document published on or after the international filing date	"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"&" document member of the same patent family
"O" document referring to an oral disclosure, use, exhibition or other means	
"P" document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search 11 APRIL 1995	Date of mailing of the international search report 17 APR 1995
Name and mailing address of the ISA/US Commissioner of Patents and Trademarks Box PCT Washington, D.C. 20231 Facsimile No. (703) 305-3230	Authorized officer Hyosuk Kim <i>Nahaytegr for</i> Telephone No. (703) 308-0196

Form PCT/ISA/210 (second sheet)(July 1992)\*

# INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US94/14832

## C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	THE JOURNAL OF IMMUNOLOGY, Volume 150, Number 2, issued 13 January 1993, Fleming et al., "Structural Analysis of the <i>CD11b</i> Gene and Phylogenetic Analysis of the $\alpha$ -Integrin Gene Family Demonstrate Remarkable Conservation of Genomic Organization and Suggest Early Diversification during Evolution", pages 480-490, see entire document.	1-38

Form PCT/ISA/210 (continuation of second sheet)(July 1992)\*

# INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US94/14832

## A. CLASSIFICATION OF SUBJECT MATTER:

IPC (6):

C07H 19/00, 21/00; C12N 5/00, 15/00, 1/20; C12P 21/06; C12Q 1/68, 1/00; C07K 1/00, 2/00, 4/00, 14/00, 16/00;  
A61K 35/14

## A. CLASSIFICATION OF SUBJECT MATTER:

US CL :

435/6, 7.1, 69.1, 240.2, 252.3, 320.1; 530/350, 387.1, 387.2, 388.1, 388.22; 536/22.1, 23.1, 23.5

